

RAT-STATS 2010 Companion Manual

Version 4

Department of Health & Human Services
Office of Inspector General
Office of Audit Services

RAT-STATS Companion Manual

TABLE OF CONTENTS

	Page
Random Numbers	1-1
Single-Stage Random Numbers	1-2
Sets of Two Numbers	1-3
Sets of Three Numbers	1-4
Sets of Four Numbers	1-5
Frames - Single Stage	1-6
Frames - Sets of Two	1-8
RHC Sample Selection	1-11
Procedure for Two-Stage RHC Sampling	1-11
Procedure for Three-Stage RHC Sampling	1-18
Summary of Input for RHC Sample Selection	1-20
Generating Spares for RHC Sampling	1-24
Comparison of RHC and Multistage SRS	1-29
Attribute Appraisals	2-1
Unrestricted Attribute Appraisal	2-2
Stratified Attribute Appraisal	2-6
Two-Stage Unrestricted	2-11
Three-Stage Unrestricted	2-19
RHC Two Stage	2-28
RHC Three Stage	2-36

Stratified Cluster	2-62
Stratified Multistage	2-68
Variable Appraisals	3-1
Unrestricted Variable Appraisal	3-2
Stratified Variable Appraisal	3-8
Using a Stratified Sample	3-8
Strata Formation	3-11
Two-Stage Unrestricted Variable Appraisal	3-17
Three-Stage Unrestricted Variable Appraisal	3-24
RHC Two Stage Variable Sampling	3-35
RHC Three Stage Variable Sampling	3-48
Stratified Cluster	3-77
Stratified Multistage	3-82
Post Stratification	3-91
Unknown Universe Size	3-99
Sample Size Determination	4-1
Variable Sample Size Determination	4-2
Unrestricted Using a Probe Sample	4-2
Unrestricted Using Estimated Error Rate	4-7
Stratified	4-11
Total Sample Size Unknown	4-11
Total Sample Size Known	4-16
Attribute Sample Size Determination	4-22

PREFACE

The purpose of this manual is to provide:

- an overview of each program in the Windows version of RAT-STATS,
- examples illustrating the application of the software,
- snapshots of data sets used by the programs,
- some discussion regarding the program output, and
- formulas used within the software.

The intent is for the auditor/specialist to use as much of this discussion as he/she finds helpful.

While the RAT-STATS Users Guide gives descriptions of program input and output, this Companion Manual should provide insight as to how to better use the software and exactly how the program derives the results. The formulas are provided so that OAS has a single source for all formulas in the event that a question is raised as to exactly how a particular result was obtained.

We hope you find that the manual makes the OAS software easier to understand and easier to apply. Please pass on any suggestions or corrections to Office of Inspector General, Office of Public Affairs at paffairs@oig.hhs.gov.

RANDOM NUMBERS

Whatever statistical sampling design you end up using (including stratified and/or multistage), at some point in the data collection you will need one or more random samples. The next section, dealing with Unrestricted Random Sampling, will examine the mechanics and estimation procedures using such a sample in detail, but first it is necessary to discuss procedures for generating a random sample. A number of programs exist for such purposes; namely:

- Single-Stage Random Numbers
- Sets of Two Numbers
- Sets of Three Numbers
- Sets of Four Numbers
- Frames - Single Stage
- Frames - Sets of Two
- RHC Sample Selection

Single-Stage Random Numbers

This program generates an unduplicated quantity of random numbers. Random values (the output from this program) can be output in sequential order, random order, or a mixture of both. For values in sequential order, you will see the values printed in sequential order, beginning with the smallest selected item number and proceeding to the largest item number. For values in random order, the items are printed in the order in which they were selected by the program.

Example 1. A universe contains 1,000 payments and a simple random sample of 10 payments (with four spares) is needed. What items should be selected?

Solution: Using this program and a seed value of 12345, the sampled payments are those numbered as follows:

9, 236, 337, 340, 346, 497, 556, 641, 658, and 884

The four spares are payments 404, 624, 927, and 947.

Sets of Two Numbers

This program will generate unduplicated pairs of random numbers. This is useful when sample items are selected through a two-step process (e.g., page number and line number).

Random values (the output from this program) can be output in sequential order, random order, or a mixture of both. Values in sequential order will be printed in sequential order, beginning with the smallest selected item number and proceeding to the largest item number. For values in random order, the items are printed in the order in which they were selected by the program.

Example 2. Items are selected from a computer printout that had pages numbered 1 through 658 and had 66 lines on each page. A simple random sample of 10 items (with four spares) is needed. Which items should be selected?

Solution: Using this program and a seed value of 12345, the sampled payments are:

PAGE:	224	258	266	327	366	400	422	433	561	610
-------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

ITEM:	23	37	42	1	16	7	23	59	40	63
-------	----	----	----	---	----	---	----	----	----	----

The spares are: PAGE: 579 109 188 330

ITEM:	61	54	50	49
-------	----	----	----	----

Sets of Three Numbers

This program will generate unduplicated sets of three random numbers. This should be used when sample items are selected through a three-step process (e.g., month, page, and line number).

Random values (the output from this program) can be output in sequential order, random order, or a mixture of both. Values in sequential order will be printed in sequential order, beginning with the smallest selected item number and proceeding to the largest item number. For values in random order, the items are printed in the order in which they were selected by the program.

Example 3. Same as Example 2, where the pages are numbered 1 through 658 for each month. Here the universe consists of 1 year's worth (12 months) of computer printouts. We need four sample items and two spares.

Solution: Using this program and a seed value of 12345, the sampled items are:

MONTH:	3	5	6	8
--------	---	---	---	---

PAGE:	224	266	6	582
-------	-----	-----	---	-----

ITEM:	23	42	37	43
-------	----	----	----	----

The two spares are:	MONTH:	12	8
---------------------	--------	----	---

	PAGE:	623	258
--	-------	-----	-----

	ITEM:	57	37
--	-------	----	----

Sets of Four Numbers

This program will generate unduplicated sets of four random numbers. This should be used when sample items are selected through a four-step process (e.g., year, month, page, and line number).

Random values (the output from this program) can be output in sequential order, random order, or a mixture of both. For values in sequential order, you will see the values printed in sequential order, beginning with the smallest selected item number and proceeding to the largest item number. For values in random order, the items are printed in the order in which they were selected by the program.

Example 4. Same as Example 3, where the pages are numbered 1 through 658 for each month and year (total of 5 years). We need three sample items and two spares.

Solution: Using this program and a seed value of 12345, the sampled items are:

YEAR: 2 3 4

MONTH: 5 1 5

PAGE: 433 366 266

ITEM: 59 16 42

The two spares are: YEAR: 5 2

MONTH: 12 7

PAGE: 561 400

ITEM: 40 7

Frames - Single Stage

This program will generate an unduplicated set of random numbers which is useful when the universe of sampling items either (1) contains gaps of numbers or (2) the numbering system repeats within the universe. For instance, the universe of items consists of two frames, numbered 1 through 1,050 and 8,405 through 9,565.

Random values (the output from this program) can be output in sequential order, random order, or a mixture of both. For values in sequential order, you will see the values printed in sequential order, beginning with the smallest selected item number in the first frame (if any) and proceeding to the largest item number in the last frame (if any). For values in random order, the items are printed in the order in which they were selected by the program.

Example 5. A universe of items that refer to payment of a particular medical procedure are numbered as follows:

1 - 1,050 (frame 1)

and 8,405 - 9,565 (frame 2)

A sample of five items is needed. Three of these items should be in sequential order and the remaining two in random order.

Solution: Using this program and a seed number of 12345, the three sample items in sequential order are:

<u>FRAME</u>	<u>ITEM NUMBER</u>
1	20
2	8,452
2	8,584

The two items in random order are

<u>FRAME</u>	<u>ITEM NUMBER</u>
1	520
1	752

Explanation: For this example there are 2,211 items in the frame since there are 1,050 items in the first frame and $(9,565 - 8,405 + 1)$, i.e., 1,161 items, in the second frame. For the sequential items, three values between 1 and 2,211 are generated. These values are 20, 1,098, and 1,230. Since 1,098 is outside the first frame, it is in the second frame; in particular, its location in the second frame would be

$$[8,405 + (1,098 - 1,050) - 1]$$

that is, item number 8,452. Similarly, the value of 1,230 points to item number 8,584.

Similarly, for the two items in random order, the program generated two values between 1 and 2,211. These values were 520 and 752. Since both values are less than 1,050, these locations are items 520 and 752 in the first frame.

Frames - Sets of Two

This program is a combination of two programs, Frames - Single Stage and Sets of Two Numbers. The program will generate an unduplicated set of random numbers which should be used when (1) pairs of numbers are used to locate sample items (as in Sets of Two Numbers) and (2) the universe has gaps or the numbering system repeats (as in Frames - Single Stage).

Random values (the output from this program) can be output in sequential order, random order, or a mixture of both. For values in sequential order, you will see the values printed in sequential order, beginning with the smallest selected item number in the first frame (if any) and proceeding to the largest item number in the last frame (if any). For values in random order, the items are printed in the order in which they were selected by the program.

Example 6. A universe of transactions consists of three sets of computer pages, numbered as follows:

<u>FRAME</u>	<u>RANGE (Page Numbers)</u>
1	1 - 100
2	1 - 456
3	45 - 832

In addition, within each frame there are an equal number of line items per page. The range within each frame is:

<u>FRAME</u>	<u>RANGE (Number of Lines)</u>
1	1 - 66
2	1 - 66
3	1 - 66

A sample of three items in sequential order and two items in random order is needed.

Solution: Using this program and a seed value of 12345, the three sample items in sequential order are:

<u>FRAME</u>	<u>PAGE NO.</u>	<u>LINE</u>
1	12	45
3	156	21
3	236	14

The two items in random order are:

<u>FRAME</u>	<u>PAGE NO.</u>	<u>LINE</u>
2	216	64
2	357	36

Explanation: For this example, the frame consists of

$$(100)(66) + (456)(66) + (832 - 45 + 1)(66)$$

$$= 6,600 + 30,096 + 52,008 = 88,704 \text{ items.}$$

Random numbers between 1 and 88,704 are generated. Values between 1 and 6,600 will be in the first frame; values between 6,601 and $6,600 + 30,096 = 36,696$ are in the second frame; and values between 36,697 and 88,704 will come from the third frame.

The three random values (not the spares) generated are 771, 44,043, and 49,316.

To find the value corresponding to 771:

1. Find $(771/66) + 1 = 12.682$. The integer part of this is 12. So, this value is on page number (subframe) 12 of frame 1.
2. The decimal part of this number is .682. Multiply this by 66 (the number of lines per page in the first frame for this example) and round to the nearest integer. This is 45. This item is on line 45 of page 12.

To find the sample value corresponding to 44,043: This value is larger than 36,696 so it is in the third frame.

1. Find $[(44,043 - 36,696)/66] + 1 = 156.318$. Here, 45 is the low number (input) for frame 3.

The integer part of this value is 156. So, this value is on page number (subframe) 156 of frame 2.

2. The decimal part of this number is .318. Multiply this by 66 (the number of lines per page in the second frame for this example) and round to the nearest integer. This is 21. This item is on line 21 of page 156.

Finally, consider the third randomly generated value of 49,316. This is 36,696, so it is in the third frame.

1. Find $[(49,316 - 36,696)/66] + 45 = 236.212$. The integer part of 236.212 is 236. So, this value is on page number 236 of frame 3.
2. The decimal part of this number is .212. Multiply this by 66 (the number of lines per page) and round to the nearest integer. This is 14. This item is on line 14 of page 236.

RHC Sample Selection

The RHC selection/appraisal procedure is named after three statisticians -- J.N.K. Rao, H.O. Hartley, and William Cochran -- and was originally proposed in 1962. This procedure is essentially the same as single-stage SRS sampling except that the size of each primary unit (cluster) is used to select the sample. It provides a method of sample selection that allows sampling without replacement (the usual procedure) while “maintaining the flavor” of using probability proportional to size. It can be used to select primary units (P.U.s) in a two-stage design or primary and secondary units (S.U.s) in a three-stage design.

Comment: Strictly speaking, you cannot use pure probability proportional to size (pps) sampling when sampling without replacement. To understand why, consider a situation in which a population contains 10 primary units with seven “large” P.U.s and three “small” ones. If the sample size is eight, then one of the small P.U.s must be selected, regardless of its small size. The RHC procedure is not pure pps sampling, but comes very close while allowing the auditor to sample without replacement.

Procedure for Two-Stage RHC Sampling

Suppose that you have N P.U.s and you want a sample of n P.U.s. The procedure is to:

1. Randomly put (partition) the N P.U.s into n groups (no attention to size here).
2. Within each of the n groups, select one P.U. using pps.

Example 7. $N = 15$, $n = 3$

1. Generate 3 groups, each containing 5 P.U.s

One possibility: Generate 15 random numbers between 0 and 1. Suppose the smallest value is in location 8, next largest in location 5, next largest in location 13, next largest in location 12, and the next largest in location 2. The first group consists of P.U.s 2, 5, 8, 12, and 13.

Continue, to get the remaining two groups.

2. Suppose P.U. #7 is put into group 3.

Size of group 3 is 1,000 beds and size of P.U. #7 is 100 beds.

P.U. #7 will be selected from group 3 with probability $100/1000 = .1$.

Example 8. In a particular region of the U.S. there are $N = 90$ universities with government research grants. Because these universities are so widespread, it was decided to use a sample of $n = 10$ universities. Rather than audit all grants at a selected university, it was decided (based on available resources) to audit roughly 20% of the grants at each selected university. We know that there are a total of $M = 4,500$ grants in all 90 universities.

Size: As a measure of the size for each university, use the total grant dollars.

Each row of the data file will contain:

University ID, number of grants, total grant dollars

i.e.,

ID of P.U., number of S.U.s (universe) in this P.U., size of P.U.

There are 90 rows of data (one for each P.U.) contained in this data set, named UNIVRHC.TXT.

Output: The 10 universities to use in the sample (see last page of computer output) are:

UNIV78, UNIV42, UNIV49, UNIV5, UNIV19, UNIV38, UNIV62, UNIV28,
UNIV60, and UNIV75

This program will create an output file specified by the user (OutRHCsummary.txt for this illustration) that is used as one of the input files by the RHC appraisal program. Dataset

UNIVRHC.TXT, the program output, and output file OutRHCsummary.txt are contained in the following pages.

Dataset UNIVRHC.TXT

			< - - - continued - - - >			< - - - continued - - - >		
			(1)	(2)	(3)			
UNIV1	42	8	UNIV31	52	11	UNIV61	66	13
UNIV2	21	4	UNIV32	66	14	UNIV62	77	18
UNIV3	63	13	UNIV33	25	5	UNIV63	31	7
UNIV4	74	16	UNIV34	60	12	UNIV64	46	9
UNIV5	51	11	UNIV35	19	4	UNIV65	32	7
UNIV6	43	9	UNIV36	24	5	UNIV66	68	14
UNIV7	57	11	UNIV37	44	9	UNIV67	41	9
UNIV8	49	10	UNIV38	76	17	UNIV68	28	6
UNIV9	63	13	UNIV39	41	9	UNIV69	66	14
UNIV10	18	4	UNIV40	77	18	UNIV70	31	7
UNIV11	64	13	UNIV41	37	8	UNIV71	27	6
UNIV12	56	11	UNIV42	63	12	UNIV72	33	7
UNIV13	19	4	UNIV43	52	11	UNIV73	23	4
UNIV14	44	9	UNIV44	76	17	UNIV74	71	15
UNIV15	20	4	UNIV45	51	10	UNIV75	75	16
UNIV16	34	7	UNIV46	23	4	UNIV76	47	10
UNIV17	25	6	UNIV47	24	5	UNIV77	50	10
UNIV18	38	9	UNIV48	68	15	UNIV78	37	7
UNIV19	72	16	UNIV49	34	7	UNIV79	77	18
UNIV20	46	10	UNIV50	49	10	UNIV80	49	10
UNIV21	44	9	UNIV51	55	11	UNIV81	76	17
UNIV22	64	13	UNIV52	38	9	UNIV82	66	14
UNIV23	45	9	UNIV53	72	16	UNIV83	28	6
UNIV24	55	11	UNIV54	51	10	UNIV84	77	17
UNIV25	29	7	UNIV55	71	15	UNIV85	27	6
UNIV26	36	7	UNIV56	59	12	UNIV86	75	17
UNIV27	40	9	UNIV57	23	4	UNIV87	71	15
UNIV28	78	18	UNIV58	57	11	UNIV88	59	12
UNIV29	49	10	UNIV59	53	11	UNIV89	71	15
UNIV30	60	12	UNIV60	64	13	UNIV90	72	16

Columns: (1) primary unit ID
 (2) number of grants
 (3) grant dollar amount (x \$100,000) ◀ This is the size of the university.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 12:52
 NAME OF INPUT FILE: C:\TEMP\UNIVRHC.TXT

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV51                          11                      55
UNIV44                          17                      76
UNIV32                          14                      66
UNIV78 <-- selected              7                      37
UNIV79                          18                      77
UNIV2                           4                      21
UNIV52                          9                      38
UNIV33                          5                      25
UNIV47                          5                      24

GROUP TOTALS:    9                      90                      419

```

```

***** GROUP 2 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV6                           9                      43
UNIV42 <-- selected             12                      63
UNIV65                          7                      32
UNIV40                          18                      77
UNIV45                          10                      51
UNIV1                           8                      42
UNIV80                          10                      49
UNIV36                          5                      24
UNIV70                          7                      31

GROUP TOTALS:    9                      86                      412

```

```

***** GROUP 3 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV46                          4                      23
UNIV7                           11                      57
UNIV90                          16                      72
UNIV49 <-- selected              7                      34
UNIV21                          9                      44
UNIV4                          16                      74
UNIV54                          10                      51
UNIV61                          13                      66
UNIV77                          10                      50

GROUP TOTALS:    9                      96                      471

```

```

***** GROUP 4 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV73                          4                      23
UNIV50                          10                      49
UNIV58                          11                      57
UNIV57                          4                      23
UNIV82                          14                      66
UNIV23                          9                      45
UNIV5 <-- selected              11                      51
UNIV64                          9                      46

```

UNIV34	12	60
GROUP TOTALS: 9	84	420

***** GROUP 5 *****

PRIMARY UNIT IDENTIFICATION	PRIMARY UNIT SIZE	SECONDARY UNIVERSE
=====	=====	=====
UNIV84	17	77
UNIV35	4	19
UNIV16	7	34
UNIV81	17	76
UNIV27	9	40
UNIV85	6	27
UNIV19 <-- selected	16	72
UNIV68	6	28
UNIV26	7	36
GROUP TOTALS: 9	89	409

***** GROUP 6 *****

PRIMARY UNIT IDENTIFICATION	PRIMARY UNIT SIZE	SECONDARY UNIVERSE
=====	=====	=====
UNIV37	9	44
UNIV83	6	28
UNIV63	7	31
UNIV14	9	44
UNIV43	11	52
UNIV31	11	52
UNIV15	4	20
UNIV48	15	68
UNIV38 <-- selected	17	76
GROUP TOTALS: 9	89	415

***** GROUP 7 *****

PRIMARY UNIT IDENTIFICATION	PRIMARY UNIT SIZE	SECONDARY UNIVERSE
=====	=====	=====
UNIV69	14	66
UNIV18	9	38
UNIV25	7	29
UNIV59	11	53
UNIV30	12	60
UNIV10	4	18
UNIV24	11	55
UNIV62 <-- selected	18	77
UNIV17	6	25
GROUP TOTALS: 9	92	421

```

***** GROUP 8 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV28 <-- selected              18                  78
UNIV41                           8                   37
UNIV89                          15                  71
UNIV66                          14                  68
UNIV11                          13                  64
UNIV86                          17                  75
UNIV56                          12                  59
UNIV12                          11                  56
UNIV72                           7                   33

GROUP TOTALS:    9                  115                  541

```

```

***** GROUP 9 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV71                           6                   27
UNIV8                          10                  49
UNIV67                          9                   41
UNIV3                          13                  63
UNIV60 <-- selected             13                  64
UNIV76                          10                  47
UNIV74                          15                  71
UNIV9                           13                  63
UNIV20                          10                  46

GROUP TOTALS:    9                  99                   471

```

```

***** GROUP 10 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV22                          13                  64
UNIV39                           9                   41
UNIV88                          12                  59
UNIV55                          15                  71
UNIV29                          10                  49
UNIV75 <-- selected             16                  75
UNIV87                          15                  71
UNIV13                           4                   19
UNIV53                          16                  72

GROUP TOTALS:    9                  110                  521

```

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

GENERATION OF PRIMARY UNIT SAMPLE

Time: 12:52

NAME OF OUTPUT FILE: C:\TEMP\OutRHCsummary.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF PRIMARY UNITS IN THE POPULATION: 90

NUMBER OF PRIMARY UNITS SAMPLED: 10

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
UNIV78	37	7	90	9
UNIV42	63	12	86	9
UNIV49	34	7	96	9
UNIV5	51	11	84	9
UNIV19	72	16	89	9
UNIV38	76	17	89	9
UNIV62	77	18	92	9
UNIV28	78	18	115	9
UNIV60	64	13	99	9
UNIV75	75	16	110	9

NOTE: In practice, it is recommended that you not set the two seed values unless you are trying to duplicate prior results.

Output file OutRHCsummary.txt

(1)	(2)	(3)	(4)	(5)
UNIV78	37	7	90	9
UNIV42	63	12	86	9
UNIV49	34	7	96	9
UNIV5	51	11	84	9
UNIV19	72	16	89	9
UNIV38	76	17	89	9
UNIV62	77	18	92	9
UNIV28	78	18	115	9
UNIV60	64	13	99	9
UNIV75	75	16	110	9

Columns: (1) selected primary unit
 (2) number of grants (secondary units)
 (3) grant dollar amount (x \$100,000) ◀ This is the size of the primary unit.
 (4) size of the group containing this primary unit
 (5) number of universities (primary units) in this group

Procedure for Three-Stage RHC Sampling

1. A sample of primary units (clusters) is obtained as in the two-stage procedure. The size of the primary units is considered for this sample, where pps sampling is used for each group of primary units.
2. A sample of secondary units is obtained within each chosen primary unit by partitioning the primary unit into random groups of secondary units. The numbers of S.U.s in each group are chosen to be as nearly equal as possible. Using pps sampling, and the size of each secondary unit, one secondary unit is chosen from each of the secondary groups.
3. A random sample of third-stage units is obtained for each of the chosen secondary units. No attention is paid to “size” here. This is a random sample.

Example 9. The previous example was expanded to include geographical regions.

Primary units: 12 regions (select four)

Secondary units: Universities (select 10 from each region)

Third stage units: Grants (audit 20% from each university)

Selection of Primary Units

A file must be constructed containing (for each region) (1) the number of secondary units (universities) in this region and (2) the size of this region (total grant dollars). This file is GRANTSPU.TXT. The selected regions are 4, 6, 8, and 10 using seed values of 100 and 200.

File GRANTSPU.TXT

(1)	(2)	(3)
REGION1	117	1250
REGION2	63	610
REGION3	91	720
REGION4	123	1320
REGION5	107	1160
REGION6	116	1240
REGION7	102	960
REGION8	118	1300
REGION9	122	1320
REGION10	85	640
REGION11	94	930
REGION12	62	550

Columns: (1) region ID
 (2) number of universities (secondary units)
 (3) size (total grant amount x \$100,000)

Selection of Secondary Units

The three-stage RHC sample selection procedure requires the user to only obtain information for each **selected** primary unit (i.e., regions 4, 6, 8, and 10 here). The information in each of these four files consists of the size of each secondary unit (university, here) and the number of third-stage units in the universe for each secondary unit. Each of these files should resemble file UNIVRHC.TXT contained in the previous two-stage RHC discussion. Consequently, for each sampled P.U., each line of the corresponding file should contain:

university ID, number of grants at this university, total grant dollars

i.e.,

secondary unit ID, no. of third-stage units, size of S.U.

After running the **RHC Sample Selection** program on each of these four regions, the following universities were selected:

REGION	UNIVERSITIES
4	85, 46, 7, 82, 30, 34, 27, 66, 65, 80
6	113, 43, 78, 104, 89, 112, 30, 65, 3, 99
8	112, 6, 77, 93, 75, 111, 62, 115, 70, 99
10	78, 43, 7, 73, 55, 33, 10, 59, 64, 39

Selection of Third-Stage Units

Suppose that approximately 20% of the grants at each selected university are to be audited. Each of these 40 samples (4 regions x 10 universities) is obtained randomly using the **Single-Stage Random Numbers** program.

- NOTES:** (1) The previous five program runs (one at the primary level and four at the secondary level) created five output files. Using a word processor, these files can be joined to form one of the input files (the one containing primary/secondary unit information) for the three-stage RHC appraisal program which calculates the confidence interval.
- (2) This example is examined in more detail in the three-stage RHC appraisal section.

Summary of Input for RHC Sample Selection

RHC Two-Stage

The user must know:

1. The number of P.U.s in the universe and the sample.
2. The size of all P.U.s in the universe.

- Procedure:**
1. The user can set the number of S.U.s in each universe P.U. equal to one if these are difficult to determine. This is the middle column in file UNIVRHC.TXT used in the previous illustration.
 2. Next, run the **RHC Two-Stage Sample Selection** program. Store the output in a text file.

3. Using a word processor or spreadsheet, change the number of S.U.s for each sampled P.U. from one to the correct value.

RHC Three-Stage

The user must know:

1. The number of P.U.s in the universe and the sample.
2. The size of all P.U.s in the universe.
3. The number of S.U.s to be sampled within each P.U.
4. For each sampled P.U.,
 - a. The number of S.U.s in the universe within this P.U.
 - b. The size of all S.U.s within this P.U.

- Procedure:**
1. The user can set the number of S.U.s in each universe P.U. equal to 1 if these are difficult to determine.
 2. Run the **RHC Two-Stage Sample Selection** program. Store the output in a text file.
 3. Using a word processor or spreadsheet, change the number of S.U.s for each sampled P.U. from 1 to the correct value.
 4. For each sampled P.U., build a data file where each row consists of
 - a. S.U. ID
 - b. Number of third-stage units for this S.U. (OK to use a value of 1 here and correct later).
 - c. Size of this S.U.
 5. For each sampled P.U., use the data set in step 4 as input to the **RHC Two-Stage Sample Selection** program. Store the output in a text file.
 6. Using a word processor or spreadsheet, change the number of third-stage units for each sampled S.U. from 1 to the correct value.
 7. Merge the results from step 2 and each sampled P.U. into one file. See PUSURHC3.TXT (below) for an example. The values in the second column (123, 54, 44, . . .) can be set to one and later changed to the correct values. This is one of the input files to the **RHC Three-Stage Appraisal** program.

NOTE: Although this procedure allows for substituting 1s for the number of second- and third-stage units in the original pass, the required size information must be known.

8. Build the data file (file PUSURHC3.TXT for this illustration) containing the sampled third-stage units. This is the other input file required by the RHC three-stage appraisal program. Using a word processor or spreadsheet, the column of sample sizes (highlighted) was added to the files created by the five RHC Sample Selection programs.

File PUSURHC3.TXT

REGION4	123	10	1320	3410	3
UNIV85	54	11	11	125	12
UNIV46	44	9	9	131	12
UNIV7	77	15	17	119	12
UNIV82	52	10	11	129	12
UNIV30	54	11	11	141	12
UNIV34	50	10	10	140	12
UNIV27	76	15	16	138	12
UNIV66	76	15	16	128	13
UNIV65	62	12	14	125	13
UNIV80	70	14	15	155	13
REGION6	116	10	1240	3100	3
UNIV113	33	7	8	108	11
UNIV43	39	8	7	105	11
UNIV78	63	13	12	104	11
UNIV104	25	5	9	96	11
UNIV89	35	7	7	124	12
UNIV112	27	5	10	108	12
UNIV30	58	12	11	95	12
UNIV65	57	11	10	109	12
UNIV3	56	11	11	115	12
UNIV99	80	16	14	113	12
REGION8	118	10	1300	3170	3
UNIV112	75	15	16	125	11
UNIV6	34	7	8	127	11
UNIV7	51	10	11	120	12
UNIV93	54	11	11	136	12
UNIV75	52	10	11	126	12
UNIV111	84	17	17	134	12
UNIV62	64	13	14	123	12
UNIV115	59	12	13	137	12
UNIV70	65	13	14	143	12
UNIV99	60	12	13	129	12
REGION10	85	10	640	2320	3
UNIV78	39	8	7	62	8
UNIV43	42	8	7	68	8

File PUSURHC3.TXT - continued

UNIV7	56	11	9	54	8
UNIV73	27	5	5	63	8
UNIV55	78	16	12	70	8
UNIV33	65	13	10	77	9
UNIV10	60	12	9	76	9
UNIV59	52	10	8	71	9
UNIV64	50	10	8	73	9
UNIV39	38	8	6	68	9

Generating Spares for RHC Sampling

One question that arises here is what to do if one or more of the selected primary units is unattainable or unusable for some reason. There is a method of generating spares without having to start the sample selection process all over again, once the nonusable primary units have been identified.

A similar approach can be used in a three-stage plan if one or more secondary units are nonusable within a selected primary unit. The following example will illustrate how to recover when one or more primary units are nonusable with a two-stage RHC sampling plan.

Example 10. Population: $N = 90$ P.U.s (universities) Sample: $n = 5$ P.U.s

The final section of the output using the RHC sample selection program is shown below; it turns out that universities 51 (in group 5) and 69 (in group 4) could not be used.

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV47	24	5	191	18
UNIV64	46	9	175	18
UNIV52	38	9	185	18
UNIV69	66	14	203	18
UNIV51 — can't use	55	11	196	18

The corresponding output file created from the first pass is shown below.

UNIV47	24	5	191	18
UNIV64	46	9	175	18
UNIV52	38	9	185	18
UNIV69	66	14	203	18
UNIV51	55	11	196	18

The section of the output containing the contents of group 4 follows.

```

***** GROUP      4 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====            SIZE            UNIVERSE
=====
UNIV15                      4                      20
UNIV81                      17                     76
UNIV38                      17                     76
UNIV59                      11                     53
UNIV13                      4                      19
UNIV76                      10                     47
UNIV20                      10                     46
UNIV66                      14                     68
UNIV14                      9                      44
UNIV29                      10                     49
UNIV55                      15                     71
UNIV26                      7                      36
UNIV17                      6                      25
UNIV87                      15                     71
UNIV22                      13                     64
UNIV50                      10                     49
UNIV84                      17                     77
UNIV69 <-- Selected, can't use      14                     66

GROUP TOTALS:    18                      203                      957

```

Remove UNIV69 from the population and this group. Construct a data file (same format as UNIVRHC.TXT) using this group only. This file (TEMP1.TXT) is shown below.

NOTE: When constructing this file, notice that columns 2 and 3 above (i.e., PRIMARY UNIT SIZE and SECONDARY UNIVERSE) need to be switched. This was done correctly in TEMP1.TXT.

```

UNIV15    20    4
UNIV81    76    17
UNIV38    76    17
UNIV59    53    11
UNIV13    19    4
UNIV76    47    10
UNIV20    46    10
UNIV66    68    14
UNIV14    44    9
UNIV29    49    10
UNIV55    71    15
UNIV26    36    7
UNIV17    25    6
UNIV87    71    15
UNIV22    64    13
UNIV50    49    10
UNIV84    77    17

```

File TEMP1.TXT

Next, run the **RHC Sample Selection** program. Your input file is TEMP1.TXT and your sample size is 1. This generates another P.U. (university) from this group. The output from this program is shown below. UNIV22 was selected.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 13:17
 NAME OF INPUT FILE: C:\TEMP\TEMP1.txt

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                SIZE                UNIVERSE
=====                =====
UNIV81                      17                      76
UNIV76                      10                      47
UNIV15                       4                      20
UNIV20                      10                      46
UNIV59                      11                      53
UNIV13                       4                      19
UNIV50                      10                      49
UNIV87                      15                      71
UNIV22 <--- Selected          13                      64
UNIV29                      10                      49
UNIV84                      17                      77
UNIV55                      15                      71
UNIV26                       7                      36
UNIV66                      14                      68
UNIV38                      17                      76
UNIV14                       9                      44
UNIV17                       6                      25

GROUP TOTALS: 17                      189                      891
  
```

NAME OF OUTPUT FILE: C:\TEMP\OUTTEMP1.TXT

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF PRIMARY UNITS IN THE POPULATION: 17

NUMBER OF PRIMARY UNITS SAMPLED: 1

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV22	64	13	189	17

Next, repeat this for group 5. Remove UNIV51 from the population and this group. Construct data file TEMP2.TXT.

NOTE: As before, be sure to switch columns 2 and 3 when building this file.

UNIV63	31	7
UNIV18	38	9
UNIV58	57	11
UNIV31	52	11
UNIV56	59	12
UNIV90	72	16
UNIV65	32	7
UNIV12	56	11
UNIV16	34	7
UNIV2	21	4
UNIV79	77	18
UNIV74	71	15
UNIV8	49	10
UNIV86	75	17
UNIV53	72	16
UNIV23	45	9
UNIV33	25	5

File TEMP2.TXT

Again, run the **RHC Sample Selection** program. The input file is TEMP2.TXT and the sample size is 1. This generates another P.U. from this group. The output from this program is shown below. UNIV86 was selected.

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 5/22/2010      GENERATION OF PRIMARY UNIT SAMPLE      Time: 13:24
NAME OF INPUT FILE: C:\TEMP\Temp2.txt

```

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====      SIZE      UNIVERSE
UNIV18                      9                      38
UNIV90                     16                      72
UNIV63                      7                      31
UNIV65                      7                      32
UNIV31                     11                      52
UNIV56                     12                      59
UNIV23                      9                      45
UNIV86  <--- Selected         17                      75
UNIV53                     16                      72
UNIV2                      4                      21
UNIV33                      5                      25

```

< OUTPUT -- continued >

UNIV79	18	77
UNIV74	15	71
UNIV12	11	56
UNIV58	11	57
UNIV16	7	34
UNIV8	10	49

GROUP TOTALS: 17	185	866
------------------	-----	-----

NAME OF OUTPUT FILE: C:\TEMP\OUTTEMP2.TXT

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF PRIMARY UNITS IN THE POPULATION: 17

NUMBER OF PRIMARY UNITS SAMPLED: 1

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
UNIV86	75	17	185	17

Finally, be sure to update the original output file shown earlier to reflect the two new selected universities. This is shown below. This file is one of the input files to the **RHC Two-Stage Appraisal** program.

Final output file (input file to RHC appraisal program)

UNIV47	24	5	191	18
UNIV64	46	9	175	18
UNIV52	38	9	185	18
UNIV22	64	13	189	17
UNIV86	75	17	185	17

Discussion: RHC Three-Stage sampling

A similar procedure can be used to generate “spare” secondary units. For example, if one of the secondary units within a selected primary unit is nonusable, another secondary unit can be selected from this group using the procedure outlined above.

Discussion: Final-stage units

At the second stage for RHC Two-Stage sampling and the third stage for RHC Three-Stage sampling, a random sample of units is obtained. Spares for this stage can be obtained in the usual manner using the single-stage random number generator software (Single Stage Random Numbers).

Comparison of RHC and Multistage SRS

In general, you can expect greater precision with the RHC procedure, provided there is a significant correlation between the second and third columns (Number of Units and Size of Unit) of each file using the RHC sample selection procedure. To illustrate, consider the file containing the primary unit information used in the three-stage RHC illustration.

(1)	(2)	(3)	Columns:	(1) unit ID (2) number of units (3) size of unit
REGION1	117	1250		
REGION2	63	610		
REGION3	91	720		
.				
.				
.				
REGION10	85	640		
REGION11	94	930		
REGION12	62	550		

For this example, the correlation between Size of Unit and Number of Units is .958, and we would expect a two-stage RHC procedure to work quite well. For a three-stage procedure, this correlation rule must also apply within each of the sampled primary units, at the secondary unit level.

The benefits of RHC sampling include:

- increased precision if the above correlation rule is satisfied;
- maintaining the flavor of pps sampling, since pps sampling is used to select a unit from each random group;
- relatively simple and straightforward computations;
- unbiased and stable point estimate of the universe total (\hat{T}). This implies that when sampling indefinitely, \hat{T} , on the average, is equal to the actual universe total, T, and \hat{T} will exhibit relatively small variation.
- a staple point estimate of the variance of \hat{T} , producing more reliable confidence intervals. This implies that when sampling indefinitely, the lower confidence limits will exhibit relatively small variation.

ATTRIBUTE APPRAISALS

An attribute appraisal is carried out to estimate a particular universe proportion (p) and its corresponding sampling error. This proportion is typically an error rate (proportion of the universe in error) but, more generally, it is the proportion of the universe items that meet (or do not meet) a specified set of criteria. Also of interest may be the total number of items in the universe (Np) that meet the criteria.

In an attribute sample, each sample item is either a yes response (met the criteria) or no response (did not meet the criteria). This version of RAT-STATS contains eight modules that can be used to appraise an attribute sample. These sampling strategies are listed below and described in the sections to follow.

- Unrestricted
- Stratified
- Two-Stage Unrestricted
- Three-Stage Unrestricted
- RHC Two Stage
- RHC Three Stage
- Stratified Cluster
- Stratified Multistage

Unrestricted Attribute Appraisal

An **unrestricted** sample is the same as a **simple random sample**. Consequently, every sample of size n has the same chance of being selected. For an unrestricted sample, a sample of size n is randomly obtained and the number of sample elements meeting the criteria (say, x) is recorded.

Example 1. An unrestricted sample of 400 documents was obtained and examined to determine if they had the proper approval signature. In the sample, 82 of the items did not contain the proper signature (were in error). The sample error rate is then $82/400 = .205$ (i.e., 20.5%). This is the estimate of p , the error rate for the entire universe. If the universe size is $N = 10,000$, then the estimated number of universe items in error is $(10,000)(.205) = 2,050$ items.

Using the RAT-STATS software, the corresponding 90% confidence interval for the total number of universe items in error is from 1,729 to 2,403. The 90% confidence interval for the universe error rate (p) is from 17.29% to 24.03%. Notice that the (point) estimate of 20.5% is between 17.29% and 24.03% but it is not in the center of this interval. The center of the 90% confidence interval is $(17.29 + 24.03)/2 = 20.66\%$. The reason for this result is that this estimation procedure is based on the exact hypergeometric distribution, rather than the normal approximation. The resulting 95% confidence interval for p is 16.73% to 24.70% and for Np (the total number of errors in the universe) is from 1,673 to 2,470.

Discussion. Consider the 90% confidence interval. Define

$$\text{TAIL} = (1 - .90) / 2 = .05$$

The 90% confidence interval for Np is, say, k_1 to k_2 . There were $x = 82$ sample items in error, so (referring to the Formulas section below) k_1 is the smallest value of k for which the probability of observing 82 or fewer errors is $> .05$, where $.05$ is the value of TAIL. This value of k is $k_1 = 1,729$. The corresponding error rate is $1,729/10,000 = .1729$ (i.e., 17.29%). To find the upper limit of the 90% confidence interval, the program determines the largest value of k (say, k_2) for which the probability of observing $x = 82$ or more errors is $> \text{TAIL} = .05$. This is $k_2 = 2,403$ with a corresponding error rate of $2,403/10,000 = .2403$ (i.e., 24.03%). A similar argument applies to the 95% confidence interval, where now the value of TAIL is $.025$.

- NOTES:**
1. Using these definitions of k_1 and k_2 for a 90% confidence interval, the user can be assured that the actual confidence level is at least 90%. This also applies to 80% and 95% confidence intervals.
 2. In the event that no items having the characteristic(s) of interest are found in the sample, the user has the option of having the program determine both confidence limits or only the upper confidence limits.
 3. In the event that the number of items having the characteristic(s) of interest in the sample is the same as the sample size, the user has the option of having the program determine both confidence limits or only the lower confidence limits.
 4. The universe size (N) is declared to be a long integer in the RAT-STATS program. Consequently, the largest allowable universe size is $N = 2^{31} - 1 = 2,147,483,647$.

FORMULAS

To determine a 90% confidence interval for the total number of universe items in error, define

$$\text{TAIL} = (1 - .90)/2 = .05.$$

Upper Limit: Let k_2 = largest value of k for which

$$\sum_{i=0}^x \frac{\binom{k}{i} \binom{N-k}{n-i}}{\binom{N}{n}} > .05$$

where N = universe size

n = sample size

k = total number of universe items in error

x = number of sample items in error

Lower Limit: Let k_1 = smallest value of k for which

$$\sum_{i=x}^n \frac{\binom{k}{i} \binom{N-k}{n-i}}{\binom{N}{n}} > .05$$

The resulting 90% confidence interval for the total number of universe items in error is from k_1 to k_2 and the corresponding 90% confidence interval for the error rate (p) is k_1/N to k_2/N .

For a 95% confidence interval, use the same two equations, where .05 is replaced with

TAIL = .025. For an 80% confidence interval, the value of TAIL is .10.

The procedure used to derive this confidence interval can be found in the following article.

John P. Buonaccorsi (1987), "A Note on Confidence Intervals for Proportions in Finite Populations," *The American Statistician*, Vol. 41, No. 3, 215-218.

Standard Errors

For universe proportion: Standard Error = $\sqrt{\frac{\hat{p}(1-\hat{p})}{n-1} \left(1 - \frac{n}{N}\right)}$ where $\hat{p} = x/n$.

For universe total: Standard Error = $N \cdot \sqrt{\frac{\hat{p}(1-\hat{p})}{n-1} \left(1 - \frac{n}{N}\right)}$.

NOTE: RAT-STATS does not use the preceding standard errors when deriving a confidence interval for the universe proportion and universe total. Other software packages use this standard error to derive an approximate confidence interval based on the normal distribution. RAT-STATS derives an exact confidence interval based on the hypergeometric distribution.

Stratified Attribute Appraisal

In a stratified attribute sampling plan, the universe is divided into two or more nonoverlapping categories (strata). As with an unrestricted sample, the intent is to make a statistical estimate for a universe proportion (p) or a universe total (Np) that meets a specified set of criteria. This plan involves obtaining a random sample from each of the strata. The program will request the number of universe items in each stratum and these values must be known. The program will develop estimates for each stratum as well as for the entire universe.

NOTE: In the discussion to follow, we will refer to the proportion, p , as the “error rate.”

Example 2. A universe of 2,500 Medicare claims is stratified into inpatient (Stratum 1) and outpatient (Stratum 2) claims. The universe sizes are $N_1 = 1,000$ inpatient claims and $N_2 = 1,500$ outpatient claims. Of interest is the proportion, p , of claims in error (containing improper charges).

A random sample of $n_1 = 100$ inpatient claims revealed $x_1 = 2$ errors and a random sample of $n_2 = 100$ outpatient claims uncovered $x_2 = 6$ errors.

NOTE: Both random samples were obtained using the **Single-Stage Random Numbers** program whereby 100 random numbers between 1 and 1,000 were obtained for stratum 1 and 100 random numbers between 1 and 1,500 were obtained for stratum 2.

The following output was obtained from the stratified attribute appraisal program.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/15/2010 STRATIFIED ATTRIBUTE APPRAISAL Time: 10:55
 AUDIT/REVIEW: Attribute - Stratified

STRATUM =====	SAMPLE =====	*ITEMS** =====	**RATIO* =====	*UNIVERSE* =====	PROJ. ITEMS IN UNIVERSE =====
1	100	2	2.000%	1,000	20
2	100	6	6.000%	1,500	90
COMBINED	200	8	4.400%	2,500	110
STANDARD ERROR:			1.483%	37	

STRATUM =====	PRECISION AT 80% CL =====	PRECISION AT 90% CL =====	PRECISION AT 95% CL =====
1	1.711%	2.196%	2.616%
2	2.955%	3.793%	4.519%
COMBINED	1.901%	2.439%	2.907%
LOWER LIMIT - QUANTITY	62	49	37
PERCENT	2.499%	1.961%	1.493%
UPPER LIMIT - QUANTITY	158	171	183
PERCENT	6.301%	6.839%	7.307%

Discussion. The strata sample error rates are 2% and 6%. The projected number of inpatient claims in error is $(.02)(1,000) = 20$ and the projected number for the outpatient stratum is $(.06)(1,500) = 90$. Consequently, the projected value for the universe is $20 + 90 = 110$ (highlighted) with a corresponding error rate of $(110/2,500) \times 100\% = 4.4\%$ (highlighted).

A look at the inpatient stratum: The estimated error rate is 2%. The corresponding precision at the 90% confidence level is 2.196% (highlighted). The term “precision” refers to the amount that is added and subtracted to the point estimate (2%, here) in deriving a confidence interval.

Consequently, the 90% confidence interval for the proportion of inpatient claims in error is $2\% \pm 2.196\%$; that is, -0.196% to 4.196% . Since the lower limit is negative, it may be set equal to zero. Similarly, the 95% confidence interval for the proportion of inpatient claims in error is $2\% \pm 2.616\%$ (highlighted); that is, 0% to 4.616% , once again setting the lower limit equal to zero.

NOTE: These confidence intervals are not actually contained in the program output.

A look at the outpatient stratum: The estimated error rate is 6% . Continuing the discussion from the inpatient stratum, the 90% confidence interval for the proportion of outpatient claims in error is $6\% \pm 3.793\%$ (highlighted); that is, 2.207% to 9.793% . The corresponding 95% confidence interval is $6\% \pm 4.519\%$ (highlighted); that is, 1.481% to 10.519% . As before, these confidence intervals are not actually contained in the program output.

A look at the overall precision: The precision at the 90% level is 2.439% (highlighted) and so the resulting 90% confidence interval for the universe proportion of claims in error is $4.4\% \pm 2.439\%$; that is, 1.961% to 6.839% . Multiplying these two values by 2,500 (and dividing by 100), the corresponding 90% confidence interval for the total number of universe claims in error is 49 to 171. Notice that these values are rounded to the nearest integer. Using the precision at the 95% confidence level (i.e., 2.907%), the 95% confidence interval in the previous output can be obtained.

FORMULAS

The estimated proportion for stratum i is \hat{p}_i where $\hat{p}_i = x_i / n_i$ and where x_i is the number of sample elements in stratum i in error and n_i is the number of sample items from stratum i . The value of Ratio is $\hat{p}_i \times 100\%$. The Projected Items in Universe for stratum i is $(\hat{p}_i)(N_i)$ where N_i is the number of universe items in stratum i . The PRECISION AT 90% CL (CL stands for confidence level) is 1.644853626951 times the standard error of \hat{p}_i ; that is

$$1.644853626951 \sqrt{\frac{N_i - n_i}{N_i} \cdot \frac{\hat{p}_i(1 - \hat{p}_i)}{n_i - 1}}$$

To obtain the Precision at 95% (80%) CL value for the i -th stratum, replace 1.644853626951 with 1.959963984540 (1.281551565545). The estimated standard error of \hat{p}_i is

$$SE(\hat{p}_i) = \sqrt{\frac{N_i - n_i}{N_i} \cdot \frac{\hat{p}_i(1 - \hat{p}_i)}{n_i - 1}}$$

Overall estimates: The estimate of the universe proportion (error rate) is \hat{p} (under the Ratio heading), where

$$\hat{p} = \sum_{i=1}^L \left(\frac{N_i}{N} \right) \hat{p}_i$$

the summation is over all of the L strata and $N = \sum N_i$ is the total universe size. The estimated standard error of \hat{p} is

$$SE(\hat{p}) = \sqrt{\sum_{i=1}^L \left(\frac{N_i}{N} \right)^2 [SE(\hat{p}_i)]^2}$$

The Precision at 90% CL value is $1.644853626951 \cdot SE(\hat{p})$.

The Precision at 95% CL value is obtained by replacing 1.644853626951 by 1.959963984540 in the above formula and for an 80% confidence interval, 1.644853626951 is replaced by 1.281551565545.

The resulting confidence intervals for the universe proportion (error rate) are

$$\hat{p} \pm (\text{PRECISION})$$

To obtain the confidence intervals for the universe total, multiply both ends of the confidence interval for the error rate by the universe size, N, and round to the nearest integer.

Two-Stage Unrestricted

This is a special case of **multistage sampling**. Multistage sampling is a very cost-effective sampling procedure when (1) obtaining a frame that lists all elements in the universe is very costly or impossible, or (2) the cost of obtaining observations increases as the distance separating the elements increases. Put another way, multistage sampling is cost effective when it is more costly to get to the sampling unit than it is to audit the sampling unit. The goal of multistage sampling is to get the most precise results per unit of examination cost.

General Comments

1. This is a very convenient sampling procedure for many situations because you don't have to visit all the locations.

2. For a two-stage procedure, the universe can be broken down into “subgroups.”

Example: 1st Stage: Carriers (P.U.s)

2nd Stage: Hospitals (S.U.s)

So, the procedure is to first obtain a random sample of P.U.s. These are called clusters.

Then, obtain a random sample of S.U.s within each selected P.U.. Notice that at the first stage, clusters are the sampling unit (sampling units are not always individual people, records, etc.).

3. You can estimate cost overpayments for the entire universe with multistage sampling; it is very useful for large, widespread universes.

Example 3. In a particular region of the U.S. there are $N = 90$ universities with government research grants. Because these universities are so widespread, it was decided to use a two-stage sample using 10 universities. Rather than audit all grants at a selected university, it was decided

(based on available resources) to audit roughly 20% of the grants at each selected university to estimate the proportion of grants containing charges after the scheduled completion of the grant. The following data were obtained, where a_i (p_i) is the number (proportion) of grants in the sample from the i -th university containing such charges, m_i is the number of audited (sampled) grants at the i -th university, and M_i is the total number of grants in the audit universe at the i -th university.

Univ.	M_i	m_i	a_i	p_i	$M_i p_i$
1	50	10	4	.400	20.00
2	65	13	5	.385	25.00
3	45	9	2	.222	10.00
4	48	10	3	.300	14.40
5	52	10	5	.500	26.00
6	58	12	3	.250	14.50
7	42	8	3	.375	15.75
8	66	13	4	.308	20.31
9	40	8	2	.250	10.00
10	<u>56</u>	<u>11</u>	4	.364	<u>20.36</u>
	522	104			176.32

Define M to be the total number of secondary units (grants) in the universe. In practice, M may be known or unknown. If M is unknown, the **Two-Stage Unrestricted** program estimates the universe proportion using a ratio estimator. No estimate of the universe total (total number of grants containing improper charges) is available. This is illustrated in the computer output to follow where M is unknown.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/30/2010 TWO STAGE UNRESTRICTED ATTRIBUTE APPRAISAL Time: 14:13
 AUDIT/REVIEW: Example
 DATA FILE: C:\Temp\DATA2STG.TXT

PRIMARY UNIT =====	UNIVERSE =====	SAMPLE SIZE =====	SAMPLE ITEMS WITH CHARACTERISTIC(S) =====	RATIO =====
1	50	10	4	40.00%
2	45	9	2	22.22%
3	52	10	5	50.00%
4	42	8	3	37.50%
5	40	8	2	25.00%
6	65	13	5	38.46%
7	48	10	3	30.00%
8	58	12	3	25.00%
9	66	13	4	30.77%
10	56	11	4	36.36%
TOTALS	522	104	35	
TOTAL PRIMARY UNITS IN THE UNIVERSE				90
OVERALL RATIO				33.78%
STANDARD ERROR				2.85%
CONFIDENCE LEVEL				
		80 PERCENT	90 PERCENT	95 PERCENT
LOWER LIMIT FOR PROPORTION		30.13%	29.09%	28.19%
UPPER LIMIT FOR PROPORTION		37.43%	38.47%	39.36%

Discussion. The estimate of the universe proportion, p , is

$$\hat{p}_r = \frac{\text{(estimated number of grants containing improper charges in the sampled universities)}}{\text{(number of grants in the sampled universities)}}$$

$$= [(50)(.4000) + (45)(.2222) + \cdots + (56)(.3636)] / (50 + 45 + \cdots + 56)$$

$$= 176.32 / 522 = .3378 \text{ (that is, 33.78\%)}$$

The estimated variance of \hat{p}_r is

$$v(\hat{p}_r) = (\text{Standard Error})^2 = (.0285)^2 = .000812.$$

NOTE: There is a formula for $v(\hat{p}_r)$ in the formula section.

The corresponding approximate 95% confidence interval for the universe proportion is

$$.3378 \pm (\text{Precision at 95\% Confidence Level})$$

The Precision at 95% Confidence Level value is the same as 1.959963984540

times the (Standard Error). So, the resulting 95% confidence interval can also be written

$$.3378 \pm (1.959963984540)(\text{Standard Error})$$

$$.3378 \pm (1.959963984540)(.0285)$$

$$.3378 \pm .0559$$

$$.2819 \text{ to } .3937 \text{ (28.19\% to 39.36\%)}$$

NOTE 1: When the value of M is unknown in the formula for $v(\hat{p}_r)$ (the case here), it is acceptable to replace this value with the average value of M for the sample (as was done in this illustration). This value is $522/10 = 52.2$. This is an advantage of using this estimator, since it does not require knowledge of M. If **M is known**, the user has two choices: (1) use the above ratio estimator, where now M is known or (2) use an unbiased estimator of p, illustrated in Example 4.

NOTE 2: If the value of M is known, the RAT-STATS software uses the unbiased estimator, illustrated in Example 4.

Example 4. Suppose that in Example 3, it is known that there is a total number of $M = 4,500$ grants (secondary units) in all 90 universities. As a result, \overline{M} is known and is equal to $M/N = 4,500/90 = 50$. The following output is obtained. Notice that estimated (projected) totals for each sampled university (primary unit) and for the entire universe are provided.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/30/2010 TWO STAGE UNRESTRICTED ATTRIBUTE APPRAISAL Time: 13:52
 AUDIT/REVIEW: Example
 DATA FILE: C:\Temp\DATA2STG.TXT

PRIMARY UNIT =====	UNIVERSE =====	SAMPLE SIZE =====	SAMPLE ITEMS WITH CHARACTERISTIC(S) =====	RATIO =====	PROJECTED =====
1	50	10	4	40.00%	20
2	45	9	2	22.22%	10
3	52	10	5	50.00%	26
4	42	8	3	37.50%	16
5	40	8	2	25.00%	10
6	65	13	5	38.46%	25
7	48	10	3	30.00%	14
8	58	12	3	25.00%	15
9	66	13	4	30.77%	20
10	56	11	4	36.36%	20
TOTALS	522	104	35		
OVERALL TOTALS					
90	4,500			35.26%	1,587
STANDARD ERROR				3.67%	165
CONFIDENCE LEVEL		80 PERCENT	90 PERCENT	95 PERCENT	
LOWER LIMIT FOR PROPORTION		30.56%	29.22%	28.06%	
UPPER LIMIT FOR PROPORTION		39.97%	41.31%	42.47%	
LOWER LIMIT FOR TOTAL		1,375	1,315	1,263	
UPPER LIMIT FOR TOTAL		1,799	1,859	1,911	

Discussion. An unbiased estimator of the universe proportion, p , is $\left(\frac{N}{M}\right)\left(\frac{A}{B}\right)$

where A = estimated number of grants containing improper charges in the sampled universities

B = number of sampled universities

and A/B = the projected average number of grants containing improper charges for the sampled universities

$$\begin{aligned}\text{So, } A/B &= [(50)(.4000) + (65)(.3846) + \cdots + (56)(.3636)] / 10 \\ &= 176.32 / 10 = 17.632 \text{ grants}\end{aligned}$$

The projected number of grants containing improper charges for the universe is $N \cdot (A/B)$; that is, $(90)(17.632) = 1586.88$ grants. Since there are 4500 grants in the universe, then the estimated proportion of grants containing improper charges is $1586.88/4,500 = .3526$ (35.26%).

The corresponding approximate 95% confidence interval for the universe proportion is

$$.3526 \pm (\text{Precision at 95\% Confidence Level}),$$

which is the same as $.3526 \pm 1.959963984540(\text{Standard Error})$. So, the resulting 95% confidence interval is

$$.3526 \pm (1.959963984540)(\text{Standard Error})$$

$$.3526 \pm (1.959963984540)(.0367)$$

$$.3526 \pm .0720$$

$$.2806 \text{ to } .4247 \text{ (28.06\% to 42.47\%)}.$$

The corresponding 95% confidence interval for the total number of grants in the universe containing improper charges is

$$1587 \pm 324; \text{ that is, } 1263 \text{ to } 1911 \text{ grants.}$$

NOTE: Formulas for \hat{p}_u and the corresponding confidence interval are contained in the formula section.

FORMULAS

Case 1: When the total number of secondary units in the universe (M) is unknown, the ratio estimator for the universe proportion is used. This estimator will be called \hat{p}_r . Define:

M_i = number of secondary units in the universe for the i -th sampled primary unit, m_i of which are sampled

\hat{p}_i = proportion of secondary units having the attribute of interest in the i-th sampled primary unit

n = number of sampled primary units

N = number of primary units in the universe (must be known)

M = number of secondary units in the universe (may be known or unknown)

\bar{M} = average number of secondary units per primary units in the universe. This is equal to M/N if M is known. It can be estimated using \bar{m} if M is unknown, where \bar{m} is the average number of secondary units in the sampled primary units.

The estimate of the universe proportion having the attribute of interest is

$$\hat{p}_r = \frac{\sum_{i=1}^n M_i \hat{p}_i}{\sum_{i=1}^n M_i}$$

The estimated variance of \hat{p}_r is

$$v(\hat{p}_r) = \left(\frac{N-n}{N} \right) \left(\frac{1}{n\bar{M}^2} \right) \left(\frac{\sum_{i=1}^n M_i^2 (\hat{p}_i - \hat{p}_r)^2}{n-1} \right) + \frac{1}{nN\bar{M}^2} \sum_{i=1}^n M_i^2 \left(\frac{M_i - m_i}{M_i} \right) \frac{\hat{p}_i (1 - \hat{p}_i)}{m_i - 1}$$

NOTE: The standard error of \hat{p}_r is the square root of $v(\hat{p}_r)$.

Case 2: When the total number of secondary units in the universe (M) is known, an unbiased estimator for the universe proportion is used. This estimator will be called \hat{p}_u .

$$\hat{p}_u = \frac{N}{M} \frac{\sum_{i=1}^n M_i \hat{p}_i}{n}$$

The estimated variance of \hat{p}_u is

$$v(\hat{p}_u) = \left(\frac{N-n}{N} \right) \left(\frac{1}{n\bar{M}^2} \right) \left(\frac{\sum_{i=1}^n (M_i \hat{p}_i - \bar{M} \hat{p}_u)^2}{n-1} \right) + \frac{1}{nN\bar{M}^2} \sum_{i=1}^n M_i^2 \left(\frac{M_i - m_i}{M_i} \right) \frac{\hat{p}_i(1 - \hat{p}_i)}{m_i - 1}$$

NOTE 1: The standard error of \hat{p}_u is the square root of $v(\hat{p}_u)$.

NOTE 2: When estimating the total number of secondary units in the universe having the attribute of interest, both \hat{p}_u and the standard error of \hat{p}_u are multiplied by M.

The Precision at the 95% Confidence Level value for the universe total is (1.959963984540)(M)(standard error of \hat{p}_u). For the Precision at the 90% Confidence Level value, replace 1.959963984540 with 1.644853626951 and for the Precision at the 80% Confidence Level value, replace 1.959963984540 with 1.281551565545.

Three-Stage Unrestricted

Example 5. The situation discussed in Example 4 was extended the following year to a three stage procedure by defining:

Stage 1: REGION (select 4 out of 12 regions)

Stage 2: UNIVERSITY (select 10 from each selected region)

Stage 3: GRANT (select approximately 20% of all grants at each university)

Using the random number module (**Single-Stage Random Numbers**), regions 5, 7, 8, and 10 were selected as the sampled primary units. Next, 10 universities (secondary units) were randomly selected (again using program **Single- Stage Random Numbers**) from the available universities in each of the four selected regions. The following data were obtained, where M_i is the number of grants in the universe for each university, m_i is the number of audited grants at each university (chosen to be roughly 20% of M_i), and a_i is the number of grants in the sample from the i -th university containing charges after the scheduled completion of the grant (in error).

REGION 5 (contains 90 secondary units, 10 to be sampled)

Univ.	M_i	m_i	a_i
1	47	9	3
2	51	10	2
3	45	9	4
4	46	9	1
5	46	9	3
6	50	10	1
7	50	10	4
8	57	11	3
9	54	11	4
10	64	13	2

REGION 7 (contains 110 secondary units, 10 to be sampled)

Univ.	M_i	m_i	a_i
1	53	11	2
2	59	12	5
3	52	10	1
4	67	13	3
5	59	12	1
6	73	15	6
7	51	10	3
8	75	15	2
9	66	13	1
10	58	12	4

REGION 8 (contains 85 secondary units, 10 to be sampled)

Univ.	M_i	m_i	a_i
1	45	9	3
2	39	8	2
3	43	9	4
4	34	7	1
5	54	11	2
6	54	11	3
7	34	7	1
8	59	12	1
9	49	10	4
10	43	9	2

REGION 10 (contains 120 secondary units, 10 to be sampled)

Univ.	M_i	m_i	a_i
1	59	12	2
2	68	14	6
3	57	11	3
4	72	14	6
5	70	14	1
6	73	15	2
7	83	17	5
8	89	18	4
9	73	15	3
10	77	15	2

The resulting data set is called DATA3ST.TXT and is shown on the next page. The corresponding computer output using the **Three-Stage Unrestricted** program immediately follows. For this illustration, the total number of third-stage units in the universe (S) is unknown.

--- Data set DATA3ST.TXT ---

REGION 5 90 10 ————— **There are 90 secondary units (universities) in this primary unit (region); 10 were audited.**

UNIV1	47	9	3
UNIV2	51	10	2
UNIV3	45	9	4
UNIV4	46	9	1
UNIV5	46	9	3
UNIV6	50	10	1
UNIV7	50	10	4
UNIV8	57	11	3
UNIV9	54	11	4
UNIV10	64	13	2

REGION 7 110 10

UNIV1 53 11 2 ————— **In UNIV1, there were 53 grants (third-stage units). Eleven of these grants were sampled and two contained improper charges.**

UNIV2	59	12	5
UNIV3	52	10	1
UNIV4	67	13	3
UNIV5	59	12	1
UNIV6	73	15	6
UNIV7	51	10	3
UNIV8	75	15	2
UNIV9	66	13	1
UNIV10	58	12	4

REGION 8 85 10

UNIV1	45	9	3
UNIV2	39	8	2
UNIV3	43	9	4
UNIV4	34	7	1
UNIV5	54	11	2
UNIV6	54	11	3
UNIV7	34	7	1
UNIV8	59	12	1
UNIV9	49	10	4
UNIV10	43	9	2

REGION 10 120 10

UNIV1	59	12	2
UNIV2	68	14	6
UNIV3	57	11	3
UNIV4	72	14	6
UNIV5	70	14	1
UNIV6	73	15	2
UNIV7	83	17	5
UNIV8	89	18	4
UNIV9	73	15	3
UNIV10	77	15	2

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES

Date: 5/30/2010 THREE STAGE ATTRIBUTE APPRAISAL Time: 15:18
AUDIT/REVIEW: Example
NAME OF INPUT FILE: C:\Temp\DATA3ST.TXT

FIRST STAGE SECOND STAGE =====	NEXT STAGE UNIVERSE =====	SAMPLE SIZE =====	MEETING CRITERIA =====	RATIO =====
REGION 5	90	10		
UNIV1	47	9	3	33.33%
UNIV3	51	10	2	20.00%
UNIV3	45	9	4	44.44%
UNIV4	46	9	1	11.11%
UNIV5	46	9	3	33.33%
UNIV6	50	10	1	10.00%
UNIV7	50	10	4	40.00%
UNIV8	57	11	3	27.27%
UNIV9	54	11	4	36.36%
UNIV10	64	13	2	15.38%
TOTALS	510	101	27	
REGION 7	110	10		
UNIV1	53	11	2	18.18%
UNIV2	59	12	5	41.67%
UNIV3	52	10	1	10.00%
UNIV4	67	13	3	23.08%
UNIV5	59	12	1	8.33%
UNIV6	73	15	6	40.00%
UNIV7	51	10	3	30.00%
UNIV8	75	15	2	13.33%
UNIV9	66	13	1	7.69%
UNIV10	58	12	4	33.33%
TOTALS	613	123	28	
REGION 8	85	10		
UNIV1	45	9	3	33.33%
UNIV2	39	8	2	25.00%
UNIV3	43	9	4	44.44%
UNIV4	34	7	1	14.29%
UNIV5	54	11	2	18.18%
UNIV6	54	11	3	27.27%
UNIV7	34	7	1	14.29%
UNIV8	59	12	1	8.33%
UNIV9	49	10	4	40.00%
UNIV10	43	9	2	22.22%
TOTALS	454	93	23	
REGION 10	120	10		
UNIV1	59	12	2	16.67%
UNIV2	68	14	6	42.86%
UNIV3	57	11	3	27.27%
UNIV4	72	14	6	42.86%
UNIV5	70	14	1	7.14%
UNIV6	73	15	2	13.33%
UNIV7	83	17	5	29.41%
UNIV8	89	18	4	22.22%
UNIV9	73	15	3	20.00%

UNIV10	77	15	2	13.33%
TOTALS	721	145	34	
OVERALL TOTALS		UNIVERSE	SAMPLED	
=====		=====	=====	
FIRST STAGE		12	4	
SECOND STAGE		405{ }	40	
THIRD STAGE		2,298{ }	462	
SAMPLED ITEMS MEETING CRITERIA			112	
{ } UNIVERSE SIZES FOR THE SECOND AND THIRD STAGES REPRESENT THE UNIVERSES FOR THE SAMPLED PRIOR STAGE.				
OVERALL POINT ESTIMATE OF THE PROPORTION			24.06%	
OVERALL STANDARD ERROR (PROPORTION)			1.35%	
OVERALL POINT ESTIMATE OF UNIVERSE TOTAL			17,210	
OVERALL STANDARD ERROR (TOTAL)			2,415	
CONFIDENCE LEVEL	80 PERCENT	90 PERCENT	95 PERCENT	
LOWER LIMIT FOR PROPORTION	22.33%	21.84%	21.42%	
UPPER LIMIT FOR PROPORTION	25.78%	26.27%	26.70%	
LOWER LIMIT FOR TOTAL	14,115	13,238	12,477	
UPPER LIMIT FOR TOTAL	20,304	21,182	21,942	

Highlighted values:

- (1) 33.33% is 3/9
- (2) 405 = 90 (in Region 5) + 110 (in Region 7) + 85 (in Region 8) + 120 (in Region 10)
- (3) 2,298 = (47 + ... + 64) in Region 5 + (53 + ... + 58) in Region 7 + (45 + ... + 43) in Region 8 + (59 + ... + 77) in Region 10.

Discussion. Based on the preceding output, the following results were obtained:

- (1) Estimate of the proportion of grants in the universe containing improper charges is .2406 (24.06%). This uses Equation 1 in the Formulas and Definitions section.
- (2) The 95% confidence interval for the universe proportion is from .2142 to .2670.
- (3) Estimate of the total number of grants in the universe containing improper charges is 17,210. This uses Equation 5 in the Formulas and Definitions section.
- (4) The 95% confidence interval for the total number of grants containing improper charges is from 12,477 to 21,942.

The standard error of the proportion estimate is .0135 (1.35%) and the 95% confidence interval is $.2406 \pm (1.959963984540)(.0135)$; that is, .2142 to .2670. The value of .0135 is found by taking the square root of the value obtained using Equation 2. For the universe total, the standard error is 2,415 (using the square root of Equation 6) and the 95% confidence interval is $17,210 \pm (1.959963984540)(2,415)$; that is, 12,477 to 21,942.

Is the total number of third-stage units in the universe known or unknown?

Let S = the total number of third-stage units in the universe. Two cases will be considered:

Case 1: S is unknown.

Case 2: S is known.

For Case 1:

To estimate the proportion, p , use the ratio (biased) estimator (\hat{p}_r). To estimate the number in the population (T) having the attribute of interest, use the unbiased estimator (\hat{T}_u).

For Case 2:

To estimate the proportion, p , use the unbiased estimator (\hat{p}_u). To estimate the number in the population (T) having the attribute of interest, use the unbiased estimator (\hat{T}_u).

NOTE: In the preceding example, the unbiased estimator \hat{T}_u was used where $\hat{T}_u = 17,210$. Here S was unknown and so the proportion estimator (\hat{p}_r) is from case 1 (Equation 1 in the Formulas and Definitions section). The standard error of \hat{p}_r (.0135) is the square root of the value obtained from Equation 2.

FORMULAS

Definitions

- S = total number of third-stage units in the universe
 N = number of primary units in the universe
 n = number of primary units in the sample
 M_i = number of secondary units (universe) in i -th primary unit
 m_i = number of secondary units (sample) in i -th primary unit
 B_{ij} = number of third-stage units (universe) in j -th secondary unit within i -th primary unit
 b_{ij} = number of third-stage units (sample) in j -th secondary unit within i -th primary unit
 \hat{p}_{ij} = proportion of b_{ij} sampled third-stage units in error

Formulas for \hat{p}_r (Case 1) and \hat{p}_u (Case 2)

The ratio estimator \hat{p}_r (Equation 1)

$$\hat{p}_r = \frac{\sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij} \hat{p}_{ij}}{\sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij}}$$

Estimated variance of \hat{p}_r (Equation 2)

$$\begin{aligned}
 v(\hat{p}_r) = & \frac{N-n}{n(n-1)N\bar{S}^2} \sum_{i=1}^n \left[\left(\hat{T}_i - \frac{\hat{T}_u}{N} \right) - \hat{R} \left(\hat{B}_i - \frac{\hat{B}_u}{N} \right) \right]^2 \\
 & + \frac{1}{nN\bar{S}^2} \sum_{i=1}^n \frac{M_i(M_i - m_i)}{m_i(m_i - 1)} \sum_{j=1}^{m_i} \left[\left(\hat{T}_{ij} - \frac{\hat{T}_i}{M_i} \right) - \hat{R} \left(B_{ij} - \frac{\hat{B}_i}{M_i} \right) \right]^2 \\
 & + \frac{1}{nN\bar{S}^2} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} \frac{B_{ij}(B_{ij} - b_{ij})}{b_{ij} - 1} \hat{p}_{ij}(1 - \hat{p}_{ij})
 \end{aligned}$$

$$\text{where } \hat{T}_u = \frac{N}{n} \sum_{i=1}^n \hat{T}_i$$

$$\hat{T}_i = \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij} \hat{p}_{ij}$$

$$\hat{T}_{ij} = B_{ij} \hat{p}_{ij}$$

$$\hat{B}_u = \frac{N}{n} \sum_{i=1}^n \hat{B}_i$$

$$\hat{B}_i = \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij}$$

$$\bar{S} = \frac{S}{N}$$

$$\hat{R} = \frac{\hat{T}_u}{\hat{B}_u}$$

Notes: (1) In Equation 2, $\hat{R} = \hat{p}_r$

(2) To estimate \bar{S} , use the sample estimate \bar{s} where

$$\bar{s} = \frac{\hat{B}_u}{N} = \frac{1}{n} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij}$$

The unbiased estimator \hat{p}_u (Equation 3)

$$\hat{p}_u = \frac{N}{nS} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij} \hat{p}_{ij}$$

Estimated variance of \hat{p}_u (Equation 4)

This variance can most easily be determined by setting $\hat{R} = 0$ in equation 2. Consequently,

$$\begin{aligned} v(\hat{p}_u) = & \frac{N-n}{n(n-1)N\bar{S}^2} \sum_{i=1}^n \left(\hat{T}_i - \frac{\hat{T}_u}{N} \right)^2 \\ & + \frac{1}{nN\bar{S}^2} \sum_{i=1}^n \frac{M_i(M_i - m_i)}{m_i(m_i - 1)} \sum_{j=1}^{m_i} \left(\hat{T}_{ij} - \frac{\hat{T}_i}{M_i} \right)^2 \\ & + \frac{1}{nN\bar{S}^2} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} \frac{B_{ij}(B_{ij} - b_{ij})}{b_{ij} - 1} \hat{p}_{ij}(1 - \hat{p}_{ij}) \end{aligned}$$

The unbiased estimator \hat{T}_u (Equation 5)

Since $\hat{T}_u = S \cdot \hat{p}_u$ then

$$\hat{T}_u = \frac{N}{n} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} B_{ij} \hat{p}_{ij}$$

NOTE: The value of S is not needed here.

Estimated variance of \hat{T}_u (Equation 6)

Since $v(\hat{T}_u) = S^2 \cdot v(\hat{p}_u)$, then

$$\begin{aligned} v(\hat{T}_u) = & \frac{N(N-n)}{n(n-1)} \sum_{i=1}^n \left(\hat{T}_i - \frac{\hat{T}_u}{N} \right)^2 \\ & + \frac{N}{n} \sum_{i=1}^n \frac{M_i(M_i - m_i)}{m_i(m_i - 1)} \sum_{j=1}^{m_i} \left(\hat{T}_{ij} - \frac{\hat{T}_i}{M_i} \right)^2 \\ & + \frac{N}{n} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} \frac{B_{ij}(B_{ij} - b_{ij})}{b_{ij} - 1} \hat{p}_{ij}(1 - \hat{p}_{ij}) \end{aligned}$$

NOTE: The value of S is not needed here.

RHC Two Stage

For a discussion on the motivation behind the RHC sampling procedure, refer to the RHC SAMPLE SELECTION section on page 1-11, contained in the RANDOM NUMBERS section of this manual. It provides a method of sample selection that allows sampling without replacement while “maintaining the flavor” of sampling using probability proportional to size. When the P.U.s are selected, the size of each P.U. is considered rather than obtaining a simple random sample of P.U.s.

The size of each P.U. is rather arbitrary and can be the number of people, dollars, beds (for hospitals), and so forth. In general, you can expect improved precision using the RHC procedure if there is a high correlation between the size of each P.U. and the number of S.U.s within each P.U.. In other words, P.U.s having a larger size should contain a larger number of S.U.s.

The P.U.s are selected using the RHC SAMPLE SELECTION program. A random sample is then obtained for each selected P.U. and the number of S.U.s having the attribute of interest (e.g., in error) is recorded.

Example 6. An audit was carried out for state-supported university grants in a particular region. The universe consisted of all charge vouchers recorded for these grants. Because these universities are so widespread, it was decided to employ a two-stage sample using three of the 27 state-supported universities. Rather than audit all the vouchers at a selected university, it was decided (based on available resources) to audit 250 vouchers at each selected university to estimate the proportion of vouchers containing improper charges. The universities (P.U.s) were

to be selected using the RHC procedure where the “size” of each university was the total grant dollars awarded to that university.

The following file (RHC2STAGE.TXT) was constructed:

(1)	(2)	(3)
UNIV1	14928	8
UNIV2	12454	4
UNIV3	17404	13
UNIV4	18700	16
UNIV5	15989	11
UNIV6	15046	9
UNIV7	16696	11
UNIV8	15754	10
UNIV9	17404	13
UNIV10	12100	4
UNIV11	17522	13
UNIV12	16578	11
UNIV13	12218	4
UNIV14	15164	9
UNIV15	12336	4
UNIV16	13986	7
UNIV17	12925	6
UNIV18	14457	9
UNIV19	18464	16
UNIV20	15400	10
UNIV21	15164	9
UNIV22	17522	13
UNIV23	15282	9
UNIV24	16461	11
UNIV25	13396	7
UNIV26	14222	7
UNIV27	14693	9

Data file RHC2STAGE.TXT

Columns: (1) unit ID
 (2) number of vouchers
 (3) size of university (dollar amount of grants x \$10,000)

Using the RHC SAMPLE SELECTION program, the following output is produced:

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/21/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 14:29
 NAME OF INPUT FILE: C:\TEMP\RHC2STAGE.txt

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV27                          9                14,693
UNIV2                           4                12,454
UNIV6                           9                15,046
UNIV1                           8                14,928
UNIV7 <-- Selected              11               16,696
UNIV21                          9                15,164
UNIV4                          16               18,700
UNIV5                          11               15,989
UNIV16                          7                13,986

GROUP TOTALS:    9                84                137,656

```

```

***** GROUP 2 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV19                          16               18,464
UNIV20 <-- Selected              10               15,400
UNIV14                          9                15,164
UNIV26                          7                14,222
UNIV15                          4                12,336
UNIV18                          9                14,457
UNIV24                          11               16,461
UNIV10                          4                12,100
UNIV23                          9                15,282

GROUP TOTALS:    9                79                133,886

```

```

***** GROUP 3 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV17                          6                12,925
UNIV11                          13               17,522
UNIV12                          11               16,578
UNIV8 <-- Selected              10               15,754
UNIV3                          13               17,404
UNIV9                          13               17,404
UNIV22                          13               17,522
UNIV25                          7                13,396
UNIV13                          4                12,218

GROUP TOTALS:    9                90                140,723

```


DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/21/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 14:29
 NAME OF OUTPUT FILE: C:\TEMP\OutRHCsummary.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF PRIMARY UNITS IN THE POPULATION: 27
 NUMBER OF PRIMARY UNITS SAMPLED: 3

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
UNIV7	16,696	11	84	9
UNIV20	15,400	10	79	9
UNIV8	15,754	10	90	9

The selected universities are UNIV7, UNIV8, and UNIV20. A sample of 250 vouchers is obtained at each university with the following results:

University	Number of sampled vouchers	Number of vouchers in error
UNIV7	250	8
UNIV20	250	12
UNIV8	250	5

This information is recorded in the data file (RHC2DATA.TXT) required by the appraisal program (TWO-STAGE RHC) and is shown below:

1	250	8	<-- Data file RHC2DATA.TXT
2	250	12	
3	250	5	

The final portion of the preceding output was stored by the sample selection program in file RHC2PU.TXT. This file is shown below:

Primary unit file RHC2PU.TXT

UNIV7	16696	11	84	9
UNIV20	15400	10	79	9
UNIV8	15754	10	90	9

Using these two files, the following output is generated by the TWO-STAGE RHC program:

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/21/2010 TWO STAGE RHC ATTRIBUTE APPRAISAL Time: 14:40
 AUDIT/REVIEW: Example

NAME OF DATA FILE: C:\TEMP\RHC2DATA.TXT
 NAME OF PRIMARY UNIT FILE: C:\TEMP\RHC2PU.TXT
 OUTPUT FILE: C:\TEMP\OutRHC2attr.txt

PRIMARY UNIT =====	SAMPLE SIZE =====	== ATTRIBUTE == SAMPLE TOTAL =====
1	250	8
2	250	12
3	250	5
TOTALS	750	25

P.U. NBR	PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====	=====
1	UNIV7	16,696	11	84	9
2	UNIV20	15,400	10	79	9
3	UNIV8	15,754	10	90	9
TOTALS:		47,850	31	253	27

P.U. NBR	SAMPLE SIZE	SAMPLE MEAN	SECONDARY UNIVERSE	SIZES RATIO	POINT ESTIMATE
=====	=====	=====	=====	=====	=====
1	250	.03	16,696	7.636	4,079.90
2	250	.05	15,400	7.900	5,839.68
3	250	.02	15,754	9.000	2,835.72
TOTALS:	750		47,850		12,755.30

--- VARIANCE COMPONENTS ---

P.U. NBR	WITHIN VARIANCE	BETWEEN VARIANCE	TOTAL VARIANCE
=====	=====	=====	=====
1	260,846.68	32,187.19	293,033.87
2	338,251.15	4,907,290.25	5,245,541.41
3	173,034.95	3,618,097.64	3,791,132.59
TOTALS:	772,132.78	8,557,575.09	9,329,707.87

PRIMARY UNITS SAMPLED: 3
 PRIMARY UNITS NOT SAMPLED: 24
 PRIMARY UNITS IN POPULATION: 27

PROJECTED QUANTITY IN UNIVERSE: 12,755
 STANDARD ERROR: 3,054

CONFIDENCE LEVEL	80 PERCENT	90 PERCENT	95 PERCENT
LOWER LIMIT	8,841	7,731	6,769
UPPER LIMIT	16,670	17,779	18,742
PRECISION AMOUNT	3,914	5,024	5,987
PRECISION PERCENT	30.69%	39.39%	46.93%
Z-VALUE USED	1.281551565545	1.644853626951	1.959963984540

Final results: The point estimate of the total number of vouchers in error is 12,755 with a corresponding standard error of 3,054. The resulting 90% confidence interval is from 7,731 to 17,779 vouchers. Notice that this a very wide confidence interval. The PRECISION PERCENT is 39.39%, obtained by multiplying the standard error by 1.644853626951 and dividing by the point estimate (expressed as a percentage); that is $(100)(1.644853626951)(3054)/12755$. In general, this can be reduced by sampling a larger number of P.U.s.

Discussion

For this example, $\hat{p}_1 = 8/250 = .032$, $\hat{p}_2 = 12/250 = .048$, and $\hat{p}_3 = 5/250 = .020$. Referring to the Formula section on the next page, the estimate for the total number of vouchers containing improper charges is:

$$\begin{aligned}\hat{T} &= (84/11)(16696)(.032) + (79/10)(15400)(.048) + (90/10)(15754)(.02) \\ &= 4,079.90 + 5,839.68 + 2,835.72 = 12,755 \text{ (rounded)}\end{aligned}$$

To determine the variance of \hat{T} , the first component of this variance is the “within variance” equal to $V_2 = 772,132.78$. This accounts for the variation within the primary units (universities). The larger variance component is the variation between the primary units measured by $V_1 = 8,557,575.09$. The total variance is $V_1 + V_2 = 9,329,707.87$ and the estimated standard error of \hat{T} is $\sqrt{9,329,707.87} = 3,054$.

The 95% confidence interval for T is $12,755 \pm (1.959963984540)(3,054)$; that is 6,769 to 18,742.

FORMULAS

Definitions

1. P.U. stands for primary unit and S.U. is secondary unit
2. A_i = size of i-th P.U.
3. S_i = (size of i-th P.U.)/(size of entire population) = A_i /(size of entire population)
4. B_i = total size for i-th group
5. π_i = (total size for i-th group)/(size of entire population) = B_i /(size of entire population)
6. N = number of P.U.s in the population
7. N_i = number of P.U.s in the i-th group
8. n = number of P.U.s in the sample
9. M_i = number of S.U.s in the i-th sampled P.U. (population)
10. m_i = number of S.U.s in the i-th sampled P.U. (sample)

Estimator of population total (T)

$$\hat{T} = \sum_{i=1}^n \left(\frac{B_i}{A_i} \right) M_i \hat{p}_i$$

where \hat{p}_i = proportion of m_i sampled S.U.s having the attribute of interest.

Estimated variance of \hat{T}

$$v(\hat{T}) = V_1 + V_2 \quad \text{where}$$

$$V_1 = \frac{\sum_{i=1}^n N_i^2 - N}{N^2 - \sum_{i=1}^n N_i^2} \sum_{i=1}^n \pi_i \left(\frac{M_i \hat{p}_i}{S_i} - \hat{T} \right)^2$$

and

$$V_2 = \sum_{i=1}^n \pi_i \frac{M_i}{S_i} (M_i - m_i) \frac{\hat{p}_i(1 - \hat{p}_i)}{m_i - 1}$$

NOTE: The estimated standard error of \hat{T} is $\sqrt{v(\hat{T})}$.

Approximate 95% confidence interval for the population total (T)

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval, replace 1.959963984540 with 1.281551565545.

RHC Three Stage

The RHC sampling procedure can be used for a three-stage design.

The steps for such a procedure are the following:

1. A sample of primary units (clusters) is obtained as in the one- and two-stage procedures. The size of the primary units is considered for this sample, where pps sampling is used for each group of primary units.
2. A sample of secondary units is obtained within each chosen primary unit by partitioning the primary unit into random groups. The group sizes are chosen to be as nearly equal as possible. Using pps sampling, and the size of each secondary unit, one secondary unit is chosen from each of the secondary groups.
3. A random sample of third-stage units is obtained for each of the chosen secondary units. No attention is paid to “size” here. This is a random sample.

Prior to running the appraisal program, the user must run the RAT-STATS RHC SAMPLE SELECTION program.

Example 7. The situation discussed in Example 5 in the THREE-STAGE UNRESTRICTED section will be appraised using the RHC methodology. For this example, the stages are:

Stage 1: REGION (select 4 out of 12 regions)

Stage 2: UNIVERSITY (select 10 from each selected region)

Stage 3: GRANT (select approximately 20% of all grants at each university)

Selection of Primary Units

A file must be constructed containing (for each region) (1) the number of secondary units (universities) in this region and (2) the size of this region (total dollars of grants). This file is GRANTSPU.TXT. The selected regions are 4, 6, 8, and 10 and the output file created by the RHC SAMPLE SELECTION program is GRANTSPUOUT.TXT.

Data set GRANTSPU.TXT

(1)	(2)	(3)
REGION1	117	1250
REGION2	63	610
REGION3	91	720
REGION4	123	1320
REGION5	107	1160
REGION6	116	1240
REGION7	102	960
REGION8	118	1300
REGION9	122	1320
REGION10	85	640
REGION11	94	930
REGION12	62	550

NOTE: It is okay to set the number of S.U.s [column (2)] equal to one in this file. The actual number of S.U.s must be known for the selected P.U.s. The correct number of S.U.s must then be inserted into file GRANTSPUOUT.TXT (the highlighted values).

Columns: (1) unit ID
 (2) number of universities (S.U.s)
 (3) size (total grant dollar amount x \$100,000)

--- Data set GRANTSPUOUT.TXT ---

REGION6	116	1240	3100	3
REGION4	123	1320	3410	3
REGION8	118	1300	3170	3
REGION10	85	640	2320	3

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 15:09
 NAME OF INPUT FILE: C:\TEMP\GRANTSPU.TXT

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
===== SIZE ===== UNIVERSE =====
REGION2                          610                63
REGION6 <-- Selected             1,240              116
REGION1                          1,250              117

GROUP TOTALS: 3                3,100              296
  
```

```

***** GROUP 2 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
===== SIZE ===== UNIVERSE =====
REGION4 <-- Selected             1,320              123
REGION5                          1,160              107
REGION11                         930                94

GROUP TOTALS: 3                3,410              324
  
```

```

***** GROUP 3 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
===== SIZE ===== UNIVERSE =====
REGION12                         550                62
REGION8 <-- Selected             1,300              118
REGION9                          1,320              122

GROUP TOTALS: 3                3,170              302
  
```

```

***** GROUP 4 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
===== SIZE ===== UNIVERSE =====
REGION3                          720                91
REGION7                          960              102
REGION10 <-- Selected            640                85

GROUP TOTALS: 3                2,320              278
  
```

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

In practice, do not set these seed values.

NUMBER OF PRIMARY UNITS IN THE POPULATION: 12
 NUMBER OF PRIMARY UNITS SAMPLED: 4

< Program output - continued >

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
REGION6	116	1,240	3,100	3
REGION4	123	1,320	3,410	3
REGION8	118	1,300	3,170	3
REGION10	85	640	2,320	3

NOTE: This is file GRANTSPUOUT.TXT

Selection of Secondary Units

The input for three-stage RHC program can be greatly simplified if you only obtain information for each **selected** primary unit (i.e., regions 4, 6, 8, and 10 here). The information consists of the size of each secondary unit (here, university) and the number of third-stage units in the universe for each secondary unit (it is acceptable to set these equal to one and change later). This input is shown in files REGION4.TXT, REGION6.TXT, REGION8.TXT, and REGION10.TXT. Each line in the files contains the number of third-stage units (grants) in the universe and the size of that secondary unit (total grant \$ x \$100,000), in that order.

After each of these four files is the computer output using the RHC SAMPLE SELECTION program. A sample of 10 universities is selected for each region. The results are:

REGION	UNIVERSITIES
4	85, 46, 7, 82, 30, 34, 27, 66, 65, 80
6	113, 43, 78, 104, 89, 112, 30, 65, 3, 99
8	112, 6, 77, 93, 75, 111, 62, 115, 70, 99
10	78, 43, 7, 73, 55, 33, 10, 59, 64, 39

The previous five program runs (one at the primary level and four at the secondary level) created five output files. Using a word processor or spreadsheet, these files can be joined to form one of the input files (the one containing primary/secondary unit information) for the three-stage RHC program which calculates the confidence interval. The file for this example is PUSURHC3.TXT.

Data set REGION4.TXT

< - - - continued - - - >

< - - - continued - - - >

(1) (2) (3)

UNIV1 52 11
 UNIV2 37 9
 UNIV3 38 9
 UNIV4 20 5
 UNIV5 69 15
 UNIV6 69 15
 UNIV7 77 17
 UNIV8 32 7
 UNIV9 49 10
 UNIV10 73 15
 UNIV11 21 5
 UNIV12 62 13
 UNIV13 55 11
 UNIV14 59 12
 UNIV15 55 11
 UNIV16 36 8
 UNIV17 51 11
 UNIV18 26 7
 UNIV19 25 6
 UNIV20 73 15
 UNIV21 71 15
 UNIV22 47 10
 UNIV23 34 8
 UNIV24 25 6
 UNIV25 39 9
 UNIV26 49 10
 UNIV27 76 16
 UNIV28 21 5
 UNIV29 33 8
 UNIV30 54 11
 UNIV31 45 10
 UNIV32 74 16
 UNIV33 69 14
 UNIV34 50 10
 UNIV35 29 7
 UNIV36 56 12
 UNIV37 64 14
 UNIV38 66 14
 UNIV39 63 14
 UNIV40 57 12
 UNIV41 71 15
 UNIV42 45 10
 UNIV43 21 5
 UNIV44 46 10
 UNIV45 48 10
 UNIV46 44 9
 UNIV47 71 15
 UNIV48 67 14
 UNIV49 23 6
 UNIV50 54 11

UNIV51 62 13
 UNIV52 52 11
 UNIV53 56 11
 UNIV54 70 15
 UNIV55 41 9
 UNIV56 65 14
 UNIV57 76 16
 UNIV58 30 7
 UNIV59 75 16
 UNIV60 27 7
 UNIV61 36 8
 UNIV62 61 13
 UNIV63 58 12
 UNIV64 61 13
 UNIV65 62 14
 UNIV66 76 16
 UNIV67 71 15
 UNIV68 34 8
 UNIV69 62 13
 UNIV70 23 6
 UNIV71 28 7
 UNIV72 46 10
 UNIV73 62 14
 UNIV74 67 14
 UNIV75 25 6
 UNIV76 24 6
 UNIV77 57 12
 UNIV78 44 10
 UNIV79 73 16
 UNIV80 70 15
 UNIV81 45 10
 UNIV82 52 11
 UNIV83 34 8
 UNIV84 59 12
 UNIV85 54 11
 UNIV86 31 7
 UNIV87 69 14
 UNIV88 22 6
 UNIV89 47 10
 UNIV90 57 12
 UNIV91 31 7
 UNIV92 73 15
 UNIV93 52 11
 UNIV94 22 6
 UNIV95 22 6
 UNIV96 29 7
 UNIV97 56 12
 UNIV98 74 16
 UNIV99 43 9
 UNIV100 57 12

UNIV101 34 8
 UNIV102 28 7
 UNIV103 73 15
 UNIV104 65 14
 UNIV105 68 14
 UNIV106 28 7
 UNIV107 55 11
 UNIV108 37 9
 UNIV109 54 11
 UNIV110 47 10
 UNIV111 44 9
 UNIV112 24 6
 UNIV113 50 10
 UNIV114 52 11
 UNIV115 66 14
 UNIV116 50 10
 UNIV117 66 14
 UNIV118 34 8
 UNIV119 73 16
 UNIV120 37 8
 UNIV121 42 9
 UNIV122 59 12
 UNIV123 45 11

**NOTE: This file has
123 lines.**

Columns: (1) unit ID (2) number of grants (3) size of university (grant amount x \$100,000)

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 14:21
 NAME OF INPUT FILE: C:\TEMP\REGION4.TXT

GROUPS OF SECONDARY UNITS

```

***** GROUP 1 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV57                             16                  76
UNIV48                             14                  67
UNIV35                              7                  29
UNIV107                            11                  55
UNIV85 <-- Selected                 11                  54
UNIV103                             15                  73
UNIV86                              7                  31
UNIV2                               9                  37
UNIV81                             10                  45
UNIV58                              7                  30
UNIV36                             12                  56
UNIV49                              6                  23

GROUP TOTALS: 12                      125              576

```

```

***** GROUP 2 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV52                             11                  52
UNIV6                               15                  69
UNIV46 <-- Selected                  9                  44
UNIV69                             13                  62
UNIV108                             9                  37
UNIV44                             10                  46
UNIV50                             11                  54
UNIV121                             9                  42
UNIV1                               11                  52
UNIV43                              5                  21
UNIV87                             14                  69
UNIV39                             14                  63

GROUP TOTALS: 12                      131              611

```

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

```

***** GROUP 10 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV53                             11                  56
UNIV24                              6                  25
UNIV42                             10                  45
UNIV120                             8                  37
UNIV105                             14                  68
UNIV97                              12                  56
UNIV119                             16                  73
UNIV32                             16                  74
UNIV80 <-- Selected                 15                  70

```

UNIV96	7	29
UNIV13	11	55
UNIV62	13	61
UNIV59	16	75

GROUP TOTALS: 13	155	724
------------------	-----	-----

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/3/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 14:21

NAME OF OUTPUT FILE: C:\TEMP\OutRegion4.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 123

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV85	54	11	125	12
UNIV46	44	9	131	12
UNIV7	77	17	119	12
UNIV82	52	11	129	12
UNIV30	54	11	141	12
UNIV34	50	10	140	12
UNIV27	76	16	138	12
UNIV66	76	16	128	13
UNIV65	62	14	125	13
UNIV80	70	15	155	13

<-- continued -->

UNIV1 56 10
 UNIV2 27 5
 UNIV3 56 11
 UNIV4 23 5
 UNIV5 72 13
 UNIV6 24 5
 UNIV7 61 11
 UNIV8 65 12
 UNIV9 68 13
 UNIV10 40 8
 UNIV11 64 12
 UNIV12 66 13
 UNIV13 80 14
 UNIV14 53 9
 UNIV15 36 7
 UNIV16 53 10
 UNIV17 47 9
 UNIV18 73 14
 UNIV19 41 8
 UNIV20 58 11
 UNIV21 45 9
 UNIV22 43 8
 UNIV23 56 10
 UNIV24 35 7
 UNIV25 34 7
 UNIV26 65 13
 UNIV27 78 14
 UNIV28 35 7
 UNIV29 31 6
 UNIV30 58 11
 UNIV31 29 6
 UNIV32 76 14
 UNIV33 57 10
 UNIV34 42 8
 UNIV35 69 13
 UNIV36 58 11
 UNIV37 31 6
 UNIV38 33 6
 UNIV39 40 8
 UNIV40 51 9
 UNIV41 60 11
 UNIV42 78 14
 UNIV43 39 7
 UNIV44 46 9
 UNIV45 58 11
 UNIV46 59 11
 UNIV47 53 10
 UNIV48 57 10
 UNIV49 28 6
 UNIV50 63 12
 UNIV51 31 6
 UNIV52 60 11
 UNIV53 30 6
 UNIV54 30 6
 UNIV55 40 8
 UNIV56 26 5
 UNIV57 24 5
 UNIV58 44 8

UNIV59 67 13
 UNIV60 56 10
 UNIV61 33 7
 UNIV62 40 8
 UNIV63 68 13
 UNIV64 70 13
 UNIV65 57 10
 UNIV66 40 7
 UNIV67 54 10
 UNIV68 65 12
 UNIV69 62 12
 UNIV70 28 5
 UNIV71 56 10
 UNIV72 41 8
 UNIV73 31 6
 UNIV74 31 6
 UNIV75 46 9
 UNIV76 38 7
 UNIV77 62 12
 UNIV78 63 12
 UNIV79 50 9
 UNIV80 53 9
 UNIV81 39 7
 UNIV82 39 7
 UNIV83 39 7
 UNIV84 25 5
 UNIV85 67 13
 UNIV86 47 9
 UNIV87 54 10
 UNIV88 50 9
 UNIV89 35 7
 UNIV90 66 13
 UNIV91 65 12
 UNIV92 71 13
 UNIV93 29 6
 UNIV94 74 14
 UNIV95 66 13
 UNIV96 71 13
 UNIV97 43 8
 UNIV98 62 11
 UNIV99 80 14
 UNIV100 57 11
 UNIV101 22 5
 UNIV102 33 6
 UNIV103 78 5
 UNIV104 25 9
 UNIV105 76 8
 UNIV106 39 5
 UNIV107 48 5
 UNIV108 54 7
 UNIV109 63 12
 UNIV110 28 8
 UNIV111 69 8
 UNIV112 27 10
 UNIV113 33 8
 UNIV114 52 7
 UNIV115 33 15
 UNIV116 23 10

Data set REGION6.TXT

**NOTE: This file has
116 lines.**

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 13:57
 NAME OF INPUT FILE: C:\TEMP\REGION6.txt

GROUPS OF SECONDARY UNITS

```

***** GROUP 1 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE                UNIVERSE
=====                          =====
UNIV52                             11                   60
UNIV45                             11                   58
UNIV32                             14                   76
UNIV86                              9                   47
UNIV113 <-- Selected                8                   33
UNIV85                             13                   67
UNIV109                             12                   63
UNIV87                             10                   54
UNIV2                               5                   27
UNIV80                              9                   53
UNIV53                              6                   30

GROUP TOTALS: 11                      108                  568

```

```

***** GROUP 2 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE                UNIVERSE
=====                          =====
UNIV33                             10                   57
UNIV48                             10                   57
UNIV6                               5                   24
UNIV43 <-- Selected                7                   39
UNIV68                             12                   65
UNIV41                             11                   60
UNIV46                             11                   59
UNIV1                              10                   56
UNIV40                              9                   51
UNIV88                              9                   50
UNIV36                             11                   58

GROUP TOTALS: 11                      105                  576

```

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

```

***** GROUP 10 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE                UNIVERSE
=====                          =====
UNIV20                             11                   58
UNIV22                              8                   43
UNIV39                              8                   40
UNIV111                             8                   69
UNIV100                             11                   57
UNIV29                              6                   31
UNIV105                             8                   76
UNIV79                              9                   50
UNIV99 <-- Selected                14                   80
UNIV13                             14                   80
UNIV60                             10                   56

```

UNIV54

6

30

GROUP TOTALS: 12

113

670

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/3/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 13:57

NAME OF OUTPUT FILE: C:\TEMP\OutRegion6.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 116

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
UNIV113	33	8	108	11
UNIV43	39	7	105	11
UNIV78	63	12	104	11
UNIV104	25	9	96	11
UNIV89	35	7	124	12
UNIV112	27	10	108	12
UNIV30	58	11	95	12
UNIV65	57	10	109	12
UNIV3	56	11	115	12
UNIV99	80	14	113	12

Data set REGION8.TXT

< - - - continued - - - >

< - - - continued - - - >

UNIV1 72 15
 UNIV2 44 10
 UNIV3 43 10
 UNIV4 55 12
 UNIV5 27 7
 UNIV6 34 8
 UNIV7 51 11
 UNIV8 42 10
 UNIV9 54 12
 UNIV10 25 6
 UNIV11 82 17
 UNIV12 65 14
 UNIV13 33 8
 UNIV14 48 10
 UNIV15 32 8
 UNIV16 82 17
 UNIV17 35 8
 UNIV18 54 12
 UNIV19 34 8
 UNIV20 62 14
 UNIV21 26 6
 UNIV22 31 7
 UNIV23 58 13
 UNIV24 61 13
 UNIV25 61 14
 UNIV26 54 12
 UNIV27 53 11
 UNIV28 56 12
 UNIV29 57 12
 UNIV30 26 6
 UNIV31 25 5
 UNIV32 37 9
 UNIV33 79 16
 UNIV34 60 13
 UNIV35 57 12
 UNIV36 27 7
 UNIV37 31 7
 UNIV38 75 15
 UNIV39 26 6
 UNIV40 36 9
 UNIV41 36 9
 UNIV42 49 10
 UNIV43 83 17
 UNIV44 71 15
 UNIV45 31 7
 UNIV46 42 10
 UNIV47 62 14
 UNIV48 54 11
 UNIV49 31 7
 UNIV50 80 16

UNIV51 77 16
 UNIV52 36 9
 UNIV53 75 16
 UNIV54 68 15
 UNIV55 34 8
 UNIV56 55 12
 UNIV57 42 10
 UNIV58 36 9
 UNIV59 36 9
 UNIV60 66 15
 UNIV61 61 13
 UNIV62 64 14
 UNIV63 72 15
 UNIV64 65 14
 UNIV65 58 13
 UNIV66 49 11
 UNIV67 30 7
 UNIV68 75 16
 UNIV69 33 8
 UNIV70 65 14
 UNIV71 55 12
 UNIV72 38 9
 UNIV73 36 9
 UNIV74 60 13
 UNIV75 52 11
 UNIV76 65 14
 UNIV77 49 10
 UNIV78 27 7
 UNIV79 48 10
 UNIV80 36 9
 UNIV81 66 15
 UNIV82 62 14
 UNIV83 70 15
 UNIV84 68 15
 UNIV85 53 11
 UNIV86 38 9
 UNIV87 35 8
 UNIV88 36 9
 UNIV89 26 6
 UNIV90 26 6
 UNIV91 51 11
 UNIV92 25 5
 UNIV93 54 11
 UNIV94 56 12
 UNIV95 81 17
 UNIV96 73 15
 UNIV97 44 10
 UNIV98 50 11
 UNIV99 60 13
 UNIV100 31 7

UNIV101 24 5
 UNIV102 26 6
 UNIV103 40 10
 UNIV104 77 16
 UNIV105 27 6
 UNIV106 65 15
 UNIV107 61 13
 UNIV108 36 9
 UNIV109 26 6
 UNIV110 38 9
 UNIV111 84 17
 UNIV112 75 16
 UNIV113 26 6
 UNIV114 45 10
 UNIV115 59 13
 UNIV116 59 13
 UNIV117 57 12
 UNIV118 58 12

NOTE: This file has 118 lines.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 14:03
 NAME OF INPUT FILE: C:\TEMP\REGION8.TXT

GROUPS OF SECONDARY UNITS

```

***** GROUP 1 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE                UNIVERSE
=====
UNIV54                             15                  68
UNIV46                             10                  42
UNIV33                             16                  79
UNIV86                              9                  38
UNIV112 <-- Selected               16                  75
UNIV85                             11                  53
UNIV108                             9                  36
UNIV87                              8                  35
UNIV2                              10                  44
UNIV55                              8                  34
UNIV34                             13                  60

GROUP TOTALS: 11                  125                  564

```

```

***** GROUP 2 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE                UNIVERSE
=====
UNIV47                             14                  62
UNIV50                             16                  80
UNIV6 <-- Selected                  8                  34
UNIV44                             15                  71
UNIV68                             16                  75
UNIV42                             10                  49
UNIV48                             11                  54
UNIV1                              15                  72
UNIV41                              9                  36
UNIV89                              6                  26
UNIV37                              7                  31

GROUP TOTALS: 11                  127                  590

```

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

```

***** GROUP 10 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE                UNIVERSE
=====
UNIV21                              6                  26
UNIV23                             13                  58
UNIV40                              9                  36
UNIV110                             9                  38
UNIV100                             7                  31
UNIV30                              6                  26
UNIV104                             16                  77
UNIV81                              15                  66
UNIV99 <-- Selected                13                  60
UNIV13                              8                  33
UNIV60                             15                  66

```

UNIV56 12 55

GROUP TOTALS: 12 129 572

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/3/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 14:03

NAME OF OUTPUT FILE: C:\TEMP\OutRegion8.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 118

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV112	75	16	125	11
UNIV6	34	8	127	11
UNIV7	51	11	120	12
UNIV93	54	11	136	12
UNIV75	52	11	126	12
UNIV111	84	17	134	12
UNIV62	64	14	123	12
UNIV115	59	13	137	12
UNIV70	65	14	143	12
UNIV99	60	13	129	12

Data set
REGION10.TXT

<--continued -->

UNIV1 34 6
 UNIV2 32 5
 UNIV3 69 10
 UNIV4 23 4
 UNIV5 60 9
 UNIV6 72 11
 UNIV7 56 9
 UNIV8 28 5
 UNIV9 38 6
 UNIV10 60 9
 UNIV11 58 9
 UNIV12 37 6
 UNIV13 70 10
 UNIV14 37 6
 UNIV15 81 12
 UNIV16 53 9
 UNIV17 63 10
 UNIV18 32 5
 UNIV19 33 5
 UNIV20 37 6
 UNIV21 77 11
 UNIV22 52 8
 UNIV23 63 10
 UNIV24 41 7
 UNIV25 45 8
 UNIV26 34 6
 UNIV27 61 10
 UNIV28 70 10
 UNIV29 34 5
 UNIV30 22 4
 UNIV31 66 10
 UNIV32 69 10
 UNIV33 65 10
 UNIV34 26 4
 UNIV35 43 7
 UNIV36 65 10
 UNIV37 80 12
 UNIV38 74 11
 UNIV39 38 6
 UNIV40 43 7
 UNIV41 47 8
 UNIV42 59 9
 UNIV43 42 7
 UNIV44 54 9
 UNIV45 73 11

UNIV46 78 12
 UNIV47 72 11
 UNIV48 30 5
 UNIV49 47 8
 UNIV50 52 8
 UNIV51 24 4
 UNIV52 26 4
 UNIV53 22 4
 UNIV54 57 9
 UNIV55 78 12
 UNIV56 62 10
 UNIV57 57 9
 UNIV58 68 10
 UNIV59 52 8
 UNIV60 54 9
 UNIV61 41 7
 UNIV62 61 10
 UNIV63 79 12
 UNIV64 50 8
 UNIV65 54 9
 UNIV66 53 9
 UNIV67 40 7
 UNIV68 44 7
 UNIV69 39 7
 UNIV70 72 11
 UNIV71 76 11
 UNIV72 34 5
 UNIV73 27 5
 UNIV74 40 7
 UNIV75 41 7
 UNIV76 25 4
 UNIV77 41 7
 UNIV78 39 7
 UNIV79 58 9
 UNIV80 71 11
 UNIV81 37 6
 UNIV82 30 5
 UNIV83 78 12
 UNIV84 59 9
 UNIV85 29 5

Note: This file has 85 lines.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 13:49
 NAME OF INPUT FILE: C:\TEMP\REGION10.TXT

GROUPS OF SECONDARY UNITS

```

***** GROUP 1 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV44                             9                  54
UNIV32                             10                 69
UNIV77                             7                  41
UNIV78 <-- Selected                 7                  39
UNIV2                              5                  32
UNIV50                             8                  52
UNIV34                             4                  26
UNIV46                             12                 78

GROUP TOTALS:  8                  62                 391

```

```

***** GROUP 2 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV6                              11                 72
UNIV43 <-- Selected                 7                  42
UNIV62                             10                 61
UNIV41                             8                  47
UNIV1                              6                  34
UNIV40                             7                  43
UNIV79                             9                  58
UNIV36                             10                 65

GROUP TOTALS:  8                  68                 422

```

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

```

***** GROUP 10 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV71                             11                 76
UNIV9                              6                  38
UNIV21                             11                 77
UNIV23                             10                 63
UNIV39 <-- Selected                 6                  38
UNIV29                             5                  34
UNIV72                             5                  34
UNIV13                             10                 70
UNIV51                             4                  24

GROUP TOTALS:  9                  68                 454

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 13:49
 NAME OF OUTPUT FILE: C:\TEMP\OutRegion10.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 85
 NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV78	39	7	62	8
UNIV43	42	7	68	8
UNIV7	56	9	54	8
UNIV73	27	5	63	8
UNIV55	78	12	70	8
UNIV33	65	10	77	9
UNIV10	60	9	76	9
UNIV59	52	8	71	9
UNIV64	50	8	73	9
UNIV39	38	6	68	9

--- Data set PUSURHC3.TXT ---

REGION4	123	1320	3410	3	10
UNIV85	54	11	125	12	
UNIV46	44	9	131	12	
UNIV7	77	17	119	12	
UNIV82	52	11	129	12	
UNIV30	54	11	141	12	
UNIV34	50	10	140	12	
UNIV27	76	16	138	12	
UNIV66	76	16	128	13	
UNIV65	62	14	125	13	
UNIV80	70	15	155	13	
REGION6	116	1240	3100	3	10
UNIV113	33	8	108	11	
UNIV43	39	7	105	11	
UNIV78	63	12	104	11	
UNIV104	25	9	96	11	
UNIV89	35	7	124	12	
UNIV112	27	10	108	12	
UNIV30	58	11	95	12	
UNIV65	57	10	109	12	
UNIV3	56	11	115	12	
UNIV99	80	14	113	12	
REGION8	118	1300	3170	3	10
UNIV112	75	16	125	11	
UNIV6	34	8	127	11	
UNIV7	51	11	120	12	
UNIV93	54	11	136	12	
UNIV75	52	11	126	12	
UNIV111	84	17	134	12	
UNIV62	64	14	123	12	
UNIV115	59	13	137	12	
UNIV70	65	14	143	12	
UNIV99	60	13	129	12	
REGION10	85	640	2320	3	10
UNIV78	39	7	62	8	
UNIV43	42	7	68	8	
UNIV7	56	9	54	8	
UNIV73	27	5	63	8	
UNIV55	78	12	70	8	
UNIV33	65	10	77	9	
UNIV10	60	9	76	9	
UNIV59	52	8	71	9	
UNIV64	50	8	73	9	
UNIV39	38	6	68	9	

NOTE: This is the data file constructed using the RHC SAMPLE SELECTION program to select the primary units (regions) and, within each selected primary unit, the 10 secondary units (universities). The four lines beginning with REGIONx are from the output file created during the primary unit selection (GRANTSPUOUT.TXT). A value of 10 (the number of selected universities for that region) is added to the end of each of these lines. The 10 lines after each REGIONx line consist of the output file created when selecting the universities from each region (OUTREGION4.TXT, . . . , OUTREGION10.TXT).

Selection of Third-Stage Units

Since approximately 20% of the grants at each university in the sample are to be audited, the following sample sizes are determined:

Region 4:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV85	54	11
	UNIV46	44	9
	UNIV7	77	15
	UNIV82	52	10
	UNIV30	54	11
	UNIV34	50	10
	UNIV27	76	15
	UNIV66	76	15
	UNIV65	62	12
	UNIV80	70	<u>14</u>
			122

Region 6:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV113	33	7
	UNIV43	39	8
	UNIV78	63	13
	UNIV104	25	5
	UNIV89	35	7
	UNIV112	27	5
	UNIV30	58	12
	UNIV65	57	11
	UNIV3	56	11
	UNIV99	80	<u>16</u>
			95

Region 8:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV112	75	15
	UNIV6	34	7
	UNIV7	51	10
	UNIV93	54	11
	UNIV75	52	10
	UNIV111	84	17
	UNIV62	64	13
	UNIV115	59	12
	UNIV70	65	13
	UNIV99	60	<u>12</u>
			120

Region 10:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV78	39	8
	UNIV43	42	8
	UNIV7	56	11
	UNIV73	27	5
	UNIV55	78	16
	UNIV33	65	13
	UNIV10	60	12
	UNIV59	52	10
	UNIV64	50	10
	UNIV39	38	<u>8</u>
			101

The data file containing the errors for these 438 audited grants is RHC3DATA.TXT. Each line contains (1) a counter, (2) the number of sampled (audited) secondary units (grants), and (3) the number of grants containing improper charges (in error).

File RHC3DATA.TXT

< - - - continued - - - >

```

1.1   11   2
1.2    9   4
1.3   15   3
1.4   10   2
1.5   11   5
1.6   10   2
1.7   15   2
1.8   15   4
1.9   12   1
1.10  14   3
2.1    7   2
2.2    8   4
2.3   13   5
2.4    5   1
2.5    7   3
2.6    5   2
2.7   12   3
2.8   11   2
2.9   11   4
2.10  16   3

```

```

3.1   15   1
3.2    7   3
3.3   10   2
3.4   11   4
3.5   10   3
3.6   17   6
3.7   13   0
3.8   12   1
3.9   13   2
3.10  12   3
4.1    8   5
4.2    8   1
4.3   11   5
4.4    5   3
4.5   16   3
4.6   13   4
4.7   12   0
4.8   10   2
4.9   10   3
4.10   8   3

```

**Note: This file
has 40 lines.**

(*)

(*) To illustrate, the fifth university in the fourth sampled P.U. (Region 10) had 16 grants (third-stage units) audited and three of them contained improper charges.

Finally, the three-stage RHC program is run to generate a confidence interval for the universe total using input files PUSURHC3.TXT and RHC3DATA.TXT. The output from this program is shown at the end of this section.

Summary of results

The estimate for the number of grants in error for the universe (all 12 regions) is the OVERALL POINT ESTIMATE of 15,861 with a corresponding estimated OVERALL STANDARD ERROR of 2,039 grants.

NOTE: This estimate does not require knowing the number of grants in the universe. If this value is known, you can convert the point estimate into a proportion. For example, if the total number of grants in all 12 regions is 59,200, then the point estimate for the proportion of grants in error is $15,861/59,200 = .268$ (26.8%) with a corresponding standard error of $2,039/59,200 = .034$ (3.4%).

The program also provides estimates for the number of grants in error for each sampled P.U. (region) and for each of the groups of S.U.s (universities) within each sampled region. For example, the estimated number of grants in error for Region 4 is 1,531 and the estimated number of grants in error for the group of nine universities containing UNIV85 is 112. The SIZES RATIO refers to the ratio of the size of the group containing this university to the size of this university. To illustrate, UNIV85 in Region 4 has a size of 11 and is in a group of size 125 (look at file REGION4.TXT). The SIZES RATIO here is $125/11 = 11.3636$.

The 95% confidence interval for the number of grants in error is 11,865 to 19,857. If the total number of third-stage units in the universe is known (say, 59,200), this interval can be converted into an interval for the proportion of grants in error by dividing both limits by this value (here, .200 to .335).

The PRECISION AMOUNT is the amount added and subtracted to the point estimate (15,861) to obtain the corresponding confidence interval. In the 95% confidence interval, the lower limit of 11,865 is obtained by subtracting the precision amount of 3,996 from 15,861. The

PRECISION PERCENT is the precision amount divided by the point estimate, expressed as a percentage.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/3/2010 THREE STAGE RHC ATTRIBUTE APPRAISAL Time: 10:04
 AUDIT/REVIEW: RHC 3-Stage

DATA FILE USED: C:\TEMP\RHC3DATA.txt
 PRIMARY/SECONDARY UNIVERSE FILE USED: C:\TEMP\PUSURHC3.txt
 OUTPUT FILE: C:\TEMP\OutRHC3.txt

**** SAMPLED UNITS ****	THIRD STAGE	*** ATTRIBUTE ***	
PRIMARY / SECONDARY IDENTIFICATION	UNIVERSE	SAMPLE SIZE	NO. WITH ATTRIBUTE
=====	=====	=====	=====
REGION4			
UNIV85	54	11	2
UNIV46	44	9	4
UNIV7	77	15	3
UNIV82	52	10	2
UNIV30	54	11	5
UNIV34	50	10	2
UNIV27	76	15	2
UNIV66	76	15	4
UNIV65	62	12	1
UNIV80	70	14	3
REGION6			
UNIV113	33	7	2
UNIV43	39	8	4
UNIV78	63	13	5
UNIV104	25	5	1
UNIV89	35	7	3
UNIV112	27	5	2
UNIV30	58	12	3
UNIV65	57	11	2
UNIV3	56	11	4
UNIV99	80	16	3
REGION8			
UNIV112	75	15	1
UNIV6	34	7	3
UNIV7	51	10	2
UNIV93	54	11	4
UNIV75	52	10	3
UNIV111	84	17	6
UNIV62	64	13	0
UNIV115	59	12	1
UNIV70	65	13	2
UNIV99	60	12	3
REGION10			
UNIV78	39	8	5
UNIV43	42	8	1
UNIV7	56	11	5
UNIV73	27	5	3
UNIV55	78	16	3
UNIV33	65	13	4
UNIV10	60	12	0
UNIV59	52	10	2

UNIV64	50	10	3
UNIV39	38	8	3
TOTALS	2,270	438	111

--- POINT ESTIMATES ---

*** ATTRIBUTE ***

**** SAMPLED UNITS ****		SIZES		POINT
PRIMARY / SECONDARY IDENTIFICATION	SAMPLE MEAN	RATIO		ESTIMATE
=====				
REGION4				
UNIV85	NOTE: 112 is the estimate	0.18	11.3636	112
UNIV46	for the <u>group</u> containing	0.44	14.5556	285
UNIV7	UNIV85 (not just UNIV85).	0.20	7.0000	108
UNIV82	This group contained 12	0.20	11.7273	122
UNIV30	universities shown earlier	0.45	12.8182	315
UNIV34	in the output using	0.20	14.0000	140
UNIV27	data set REGION4.TXT.	0.13	8.6250	87
UNIV66		0.27	8.0000	162
UNIV65		0.08	8.9286	46
UNIV80		0.21	10.3333	155
TOTAL			Estimate for Region 4 →	1,531
REGION6				
UNIV113		0.29	13.5000	127
UNIV43		0.50	15.0000	293
UNIV78		0.38	8.6667	210
UNIV104		0.20	10.6667	53
UNIV89		0.43	17.7143	266
UNIV112		0.40	10.8000	117
UNIV30		0.25	8.6364	125
UNIV65		0.18	10.9000	113
UNIV3		0.36	10.4545	213
UNIV99		0.19	8.0714	121
TOTAL				1,638
REGION8				
UNIV112		0.07	7.8125	39
UNIV6		0.43	15.8750	231
UNIV7		0.20	10.9091	111
UNIV93		0.36	12.3636	243
UNIV75		0.30	11.4545	179
UNIV111		0.35	7.8824	234
UNIV62		0.00	8.7857	0
UNIV115		0.08	10.5385	52
UNIV70		0.15	10.2143	102
UNIV99		0.25	9.9231	149
TOTAL				1,340
REGION10				
UNIV78		0.63	8.8571	216
UNIV43		0.13	9.7143	51
UNIV7		0.45	6.0000	153
UNIV73		0.60	12.6000	204
UNIV55		0.19	5.8333	85
UNIV33		0.31	7.7000	154

UNIV10	0.00	8.4444	0
UNIV59	0.20	8.8750	92
UNIV64	0.30	9.1250	137
UNIV39	0.38	11.3333	162

TOTAL			1,254
-------	--	--	-------

--- VARIANCE COMPONENTS FOR PRIMARY UNITS ---

**** SAMPLED UNITS **** PRIMARY UNIT IDENTIFICATION	WITHIN VARIANCE	BETWEEN VARIANCE	TOTAL VARIANCE
=====	=====	=====	=====
REGION4	4,722	59,965	64,687
REGION6	4,734	59,651	64,385
REGION8	4,102	68,169	72,272
REGION10	3,351	47,934	51,285
	(Values of V₄)	(Values of V₃)	

--- COMBINED VARIANCE COMPONENTS ---

STAGE 1	STAGES 2 AND 3	TOTAL VARIANCE
=====	=====	=====
3,466,892	690,212	4,157,104
(Value of V₁)	(Value of V₂)	

*** ATTRIBUTE ***

--- SUMMARY OF APPRAISAL RESULTS ---

PRIMARY UNITS SAMPLED	4
PRIMARY UNITS NOT SAMPLED	8
TOTAL PRIMARY UNITS	12

PROJECTED QUANTITY IN UNIVERSE	15,861
STANDARD ERROR	2,039

CONFIDENCE LEVEL	80 PERCENT	90 PERCENT	95 PERCENT
LOWER LIMIT	13,248	12,508	11,865
UPPER LIMIT	18,474	19,215	19,857
PRECISION AMOUNT	2,613	3,354	3,996
PRECISION PERCENT	16.47%	21.14%	25.19%
Z-VALUE USED	1.281551565545	1.644853626951	1.959963984540

FORMULAS

Definitions

1. S_i = (size of i-th P.U.)/(size of entire population)
2. $\pi_i = \sum S_i$ over the i-th group of P.U.s
3. S_{ij} = (size of j-th S.U. in the i-th sampled P.U.)/(size of i-th sampled P.U.)
(**Note:** denominator of S_{ij} = numerator of S_i)
4. $\pi_{ij} = \sum S_{ij}$ over the j-th group in i-th sampled P.U.
5. N = number of P.U.s (population)
6. n = number of P.U.s (sample)
7. M_i = number of S.U.s in i-th sampled P.U. (population)
8. m_i = number of S.U.s in i-th sampled P.U. (sample)
9. K_{ij} = number of records for j-th sampled S.U. in i-th sampled P.U. (population)
10. k_{ij} = number of records for j-th sampled S.U. in i-th sampled P.U. (sample)

Estimator of population total (T)

$$\hat{T} = \sum_{i=1}^n \pi_i \left(\frac{\hat{T}_i}{S_i} \right)$$

where \hat{T}_i = estimator of total for i-th sampled P.U.

$$\sum_{j=1}^{m_i} \pi_{ij} \left(\frac{\hat{T}_{ij}}{S_{ij}} \right) \quad (\text{equation 1})$$

and \hat{T}_{ij} = estimator of population total for j-th sampled S.U. in i-th sampled P.U.

$$= K_{ij} \hat{p}_{ij}$$

with \hat{p}_{ij} = proportion of k_{ij} records having the attribute of interest

NOTE: It can be shown that \hat{T} is an unbiased estimator of T.

Estimated variance of \hat{T}

$v(\hat{T}) = V_1 + V_2$ where

$$V_1 = \frac{\sum_{i=1}^n N_i^2 - N}{N^2 - \sum_{i=1}^n N_i^2} \sum_{i=1}^n \pi_i \left(\frac{\hat{T}_i}{S_i} - \hat{T} \right)^2 \quad (\text{equation 2})$$

and

$$V_2 = \sum_{i=1}^n \left(\frac{\pi_i}{S_i} \right) v(\hat{T}_i) \quad (\text{equation 3})$$

and N_i = number of P.U.s in the i-th group after the random split into n groups.

$v(\hat{T}_i)$ is obtained by applying the two stage RHC procedure within the i-th sampled P.U.; i.e., the i-th P.U. is viewed as the entire population. Consequently, $v(\hat{T}_i) = V_{3,i} + V_{4,i}$, where

$$V_{3,i} = \frac{\sum_{j=1}^{m_i} M_{ij}^2 - M_i}{M_i^2 - \sum_{j=1}^{m_i} M_{ij}^2} \sum_{j=1}^{m_i} \pi_{ij} \left(\frac{K_{ij} \hat{p}_{ij}}{S_{ij}} - \hat{T}_i \right)^2$$

and

$$V_{4,i} = \sum_{j=1}^{m_i} \pi_{ij} \frac{K_{ij}}{S_{ij}} (K_{ij} - k_{ij}) \frac{\hat{p}_{ij}(1 - \hat{p}_{ij})}{k_{ij} - 1}$$

and where (1) M_{ij} = the number of S.U.s in the j-th random group, i-th sampled P.U.

(2) \hat{p}_{ij} = proportion of the k_{ij} items having the attribute of interest for the j-th sampled S.U. within the i-th sampled P.U.

- Comments**
1. V_1 is essentially the same expression obtained for the single-stage RHC procedure and will be referred to as the “between unit” variation.
 2. V_2 is the contribution of the second- and third-stage variation and is obtained by treating each sampled P.U. as the population to be sampled using two (additional) stages.
 3. The estimated standard error of \hat{T} is $\sqrt{v(\hat{T})}$.

Approximate 95% confidence interval for the population total (T)

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval, replace 1.959963984540 with 1.281551565545.

Stratified Cluster

With this procedure, you first stratify, then obtain a cluster (single-stage) sample within each stratum. This is motivated by the discussion in the RAT-STATS User's Guide.

Example 8. In a large section of the U.S., an audit was conducted for 583 universities with health-related research grants. Two strata were defined:

Stratum 1: state-supported universities and

Stratum 2: private universities.

It was decided to estimate the proportion of contracts containing charges after the scheduled completion of the contract using the same two strata. The strata sizes were $N_1 = 415$ and $N_2 = 168$. Within each stratum, a **single-stage** cluster sample was obtained with $n_1 = 25$ universities selected from Stratum 1 (state supported universities) and $n_2 = 10$ universities from Stratum 2 (private universities). The total number of grants in the universe for Stratum 1 is 2,500 and for Stratum 2, the total number is 1,000 grants. Consequently, there are 3,500 grants in the entire universe.

NOTE 1: These sample sizes are not adequate according to OAS policy and are used here for illustration purposes only.

NOTE 2: This procedure does require knowledge of the total number of elements in the universe for each stratum.

The following data were obtained, where a_j is the number of grants containing charges after the scheduled completion of the grant for the j -th university, M_j is the number of grants (universe) for this university (all of which are audited), and p_j is a_j/M_j .

Summary Using Computer Output and Corresponding Formulas:

Stratum 1: $\Sigma a_j = 38$, $\Sigma M_j = 151$, $\hat{p}_1 = 38/151 = .2517$

The projected number in the universe for Stratum 1 is

$$\hat{T}_1 = (2500)(.2517) = 629$$

Stratum 2: $\Sigma a_j = 19$, $\Sigma M_j = 49$, $\hat{p}_2 = 19/49 = .3878$

The projected number in the universe for Stratum 2 is

$$\hat{T}_2 = (1000)(.3878) = 388$$

The estimate of the total number of grants in the universe with charges after the scheduled grant completion is

$$\hat{T} = 629 + 388 = 1017$$

The estimated proportion of grants with such charges is

$$\hat{p} = 1017 / 3500 = .2905$$

The estimated standard error for \hat{T} (using the square root of Equation 5) is 46.

The estimated standard error for \hat{p} (using the square root of Equation 6) is .0132.

Stratum 1 -- State Universities

Univ.	M _j	a _j	p _j	Univ.	M _j	a _j	p _j
1	8	2	.25	14	10	3	.30
2	12	3	.25	15	9	1	.11
3	4	2	.50	16	3	1	.33
4	5	1	.20	17	6	2	.33
5	6	1	.17	18	5	1	.20
6	6	2	.33	19	5	1	.20
7	7	2	.29	20	4	1	.25
8	5	2	.40	21	6	1	.17
9	8	2	.25	22	8	1	.12
10	3	1	.33	23	7	2	.29
11	2	0	.00	24	3	1	.33
12	6	2	.33	25	8	2	.25
13	5	1	.20				

Stratum 2 -- Private Universities

Univ.	M _j	a _j	p _j	Univ.	M _j	a _j	p _j
1	2	1	.50	6	8	3	.37
2	5	2	.40	7	6	2	.33
3	7	2	.29	8	10	4	.40
4	4	2	.50	9	3	1	.33
5	3	1	.33	10	1	1	1.00

These results are combined into data set DATACLUS.TXT (39 lines)

Data set

```

STATE UNIVERSITIES  415  25
2500
UNIV1      8  2
UNIV2     12  3
UNIV3      4  2
UNIV4      5  1
UNIV5      6  1
UNIV6      6  2
UNIV7      7  2
UNIV8      5  2
UNIV9      8  2
UNIV10     3  1
UNIV11     2  0
UNIV12     6  2
UNIV13     5  1
UNIV14    10  3
UNIV15     9  1
UNIV16     3  1
UNIV17     6  2

```

```

UNIV18      5      1
UNIV19      5      1
UNIV20      4      1
UNIV21      6      1
UNIV22      8      1
UNIV23      7      2
UNIV24      3      1
UNIV25      8      2
PRIVATE UNIVERSITIES  168
10  1000
UNIV1       2      1
UNIV2       5      2
UNIV3       7      2
UNIV4       4      2
UNIV5       3      1
UNIV6       8      3
UNIV7       6      2
UNIV8      10      4
UNIV9       3      1
UNIV10      1      1

```

The following computer printout is produced:

```

                                DEPARTMENT OF HEALTH & HUMAN SERVICES
                                OIG - OFFICE OF AUDIT SERVICES
Date: 4/20/2010                STRATIFIED CLUSTER ATTRIBUTE APPRAISAL                Time: 15:36
                                AUDIT/REVIEW: Attribute - Stratified Cluster
                                NAME OF INPUT FILE: C:\Temp\DATACLUS.TXT

```

STRATUM IDENTIFICATION CLUSTER IDENTIFICATION	SAMPLE UNIVERSE	SAMPLE SIZE	MEETING CRITERIA	PERCENT	PROJECTED QUANTITY
=====	=====	=====	=====	=====	=====
STATE UNIVERSITIES	415	25			
UNIV1	8	8	2		
UNIV2	12	12	3		
UNIV3	4	4	2		
UNIV4	5	5	1		
UNIV5	6	6	1		
UNIV6	6	6	2		
UNIV7	7	7	2		
UNIV8	5	5	2		
UNIV9	8	8	2		
UNIV10	3	3	1		
UNIV11	2	2	0		
UNIV12	6	6	2		
UNIV13	5	5	1		
UNIV14	10	10	3		
UNIV15	9	9	1		
UNIV16	3	3	1		
UNIV17	6	6	2		
UNIV18	5	5	1		
UNIV19	5	5	1		
UNIV20	4	4	1		
UNIV21	6	6	1		
UNIV22	8	8	1		

UNIV23	7	7	2		
UNIV24	3	3	1		
UNIV25	8	8	2		
STRATUM TOTALS	2,500	151	38	25.17%	629
PRIVATE UNIVERSITIES	168	10			
UNIV1	2	2	1		
UNIV2	5	5	2		
UNIV3	7	7	2		
UNIV4	4	4	2		
UNIV5	3	3	1		
UNIV6	8	8	3		
UNIV7	6	6	2		
UNIV8	10	10	4		
UNIV9	3	3	1		
UNIV10	1	1	1		
STRATUM TOTALS	1,000	49	19	38.78%	388
STRATA TOTALS	583	35			
CLUSTER UNIT TOTALS	3,500	200	57		
OVERALL TOTALS				29.05%	1,017
OVERALL STANDARD ERROR				1.32%	46
CONFIDENCE LEVEL	80 PERCENT	90 PERCENT	95 PERCENT		
LOWER LIMIT FOR PROPORTION	27.36%	26.88%	26.47%		
UPPER LIMIT FOR PROPORTION	30.74%	31.22%	31.64%		
LOWER LIMIT FOR TOTAL	958	941	926		
UPPER LIMIT FOR TOTAL	1,076	1,093	1,107		

FORMULAS

1. Estimated proportion in stratum h that possess the attribute of interest

$$\hat{p}_h = \frac{\sum_{j=1}^{n_h} a_{j,h}}{\sum_{j=1}^{n_h} M_{j,h}}$$

where $a_{j,h}$ is the number of elements in the j-th secondary unit in stratum h possessing the attribute of interest, $M_{j,h}$ is the number of secondary units in the j-th primary unit in stratum h, and n_h is the number of sample items in stratum h.

2. Estimated total number of elements in stratum h that possess the attribute of interest

$$\hat{T}_h = M_h \hat{p}_h$$

where M_h = number of secondary units in the universe for stratum h (**must be known**)

3. Estimated universe total having the attribute of interest

$$\hat{T} = \sum_{h=1}^L \hat{T}_h \quad \text{summed over the } L \text{ strata}$$

4. Estimated universe proportion having the attribute of interest is $\hat{p} = \hat{T} / M$ where M is the total number of secondary units in the universe and $M = \sum M_h$ (summed over the L strata).

5. Estimated variance of \hat{T}

$$v(\hat{T}) = \sum_{h=1}^L \frac{N_h(N_h - n_h)}{n_h(n_h - 1)} \sum_{j=1}^{n_h} (a_{j,h} - \hat{p}_h M_{j,h})^2$$

where N_h is the number of universe items in stratum h.

6. Estimated variance of \hat{p}

$$v(\hat{p}) = v(\hat{T}) / M^2$$

7. Approximate 95% confidence interval for T

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

8. Approximate 95% confidence interval for p

$$\hat{p} \pm 1.959963984540 \sqrt{v(\hat{p})}$$

NOTE: For the Precision at the 90% Confidence Level, replace 1.959963984540 with 1.644853626951 and for the Precision at the 80% Confidence Level, replace 1.959963984540 with 1.281551565545.

Stratified Multistage

As with the stratified cluster procedure, you must first stratify the universe. Rather than take a cluster (single-stage) sample within each stratum, you will obtain a multistage (two-stage or three-stage) sample within each stratum. These multistage samples may be random (using the **Two-Stage Unrestricted** or **Three-Stage Unrestricted** programs) or may be obtained using the RHC procedure and the **RHC Two-Stage** or **RHC Three-Stage** programs.

Unlike the **Stratified Cluster** program, this program requires that you first run the appropriate multistage program on each stratum and record the results. The output results are then used as input to the **Stratified Multistage** program. You may store the results from each stratum (point estimate, standard error, universe size) in a file or simply input these values interactively.

NOTE: The “universe size” refers to the number of units at the most detailed level of the multistage sample. For example, if you are obtaining a three-stage sample within each stratum, then the “universe size” refers to the total number of third-stage units within this stratum.

Example 9. This example is similar to Example 8 in the Stratified Cluster section. The universe consisting of university grants is stratified by defining

Stratum 1: state-supported universities (5,600 grants) and
and Stratum 2: private universities (3,500 grants)

Because these universities are so widespread, it was decided to employ a two-stage sample using 20 state-supported universities and 10 private universities. Rather than audit all grants at a selected university, it was decided (based on available resources) to audit roughly 20% of the grants at each selected university to estimate the proportion of grants containing charges after the

scheduled completion of the grant. The following data were obtained, where a_i is the number of grants in the sample from the i -th university containing such charges, m_i is the number of audited (sampled) grants at the i -th university, and M_i is the total number of grants in the audit universe at the i -th university.

State-supported universities

Univ.	M_i	m_i	a_i
1	41	8	2
2	62	12	3
3	21	4	2
4	23	5	1
5	31	6	1
6	32	6	2
7	33	7	2
8	27	5	2
9	41	8	2
10	16	3	1
11	9	2	0
12	31	6	2
13	27	5	1
14	49	10	3
15	46	9	1
16	15	3	1
17	30	6	2
18	24	5	1
19	23	5	1
20	21	4	1

These values are stored in data set MULSTAT1.TXT.

Private universities

Univ.	M_i	m_i	a_i
1	11	2	1
2	25	5	2
3	34	7	2
4	18	4	2
5	16	3	1
6	40	8	3
7	31	6	2
8	50	10	4
9	14	3	1
10	12	2	1

These values are stored in data set MULSTAT2.TXT.

The following two computer outputs are obtained using the **Two-Stage Unrestricted** program.

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 4/25/2010 TWO STAGE UNRESTRICTED ATTRIBUTE APPRAISAL Time: 15:02
AUDIT/REVIEW: Stratum1
DATA FILE: C:\Temp\MULSTAT1.TXT

PRIMARY UNIT =====	UNIVERSE =====	SAMPLE SIZE =====	SAMPLE ITEMS WITH CHARACTERISTIC(S) =====	RATIO =====	PROJECTED =====
1	41	8	2	25.00%	10
2	62	12	3	25.00%	16
3	21	4	2	50.00%	11
4	23	5	1	20.00%	5
5	31	6	1	16.67%	5
6	32	6	2	33.33%	11
7	33	7	2	28.57%	9
8	27	5	2	40.00%	11
9	41	8	2	25.00%	10
10	16	3	1	33.33%	5
11	9	2	0	0.00%	0
12	31	6	2	33.33%	10
13	27	5	1	20.00%	5
14	49	10	3	30.00%	15
15	46	9	1	11.11%	5
16	15	3	1	33.33%	5
17	30	6	2	33.33%	10
18	24	5	1	20.00%	5
19	23	5	1	20.00%	5
20	21	4	1	25.00%	5
TOTALS	602	119	31		
OVERALL TOTALS					
120	5,600			16.90%	946
STANDARD ERROR				1.99%	111
CONFIDENCE LEVEL		80 PERCENT	90 PERCENT	95 PERCENT	
LOWER LIMIT FOR PROPORTION		14.35%	13.63%	13.00%	
UPPER LIMIT FOR PROPORTION		19.44%	20.17%	20.79%	
LOWER LIMIT FOR TOTAL		803	763	728	
UPPER LIMIT FOR TOTAL		1,089	1,129	1,164	

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/25/2010 TWO STAGE UNRESTRICTED ATTRIBUTE APPRAISAL Time: 15:13
 AUDIT/REVIEW: Stratum 2
 DATA FILE: C:\Temp\MULSTAT2.TXT

PRIMARY UNIT =====	UNIVERSE =====	SAMPLE SIZE =====	SAMPLE ITEMS WITH CHARACTERISTIC(S) =====	RATIO =====	PROJECTED =====
1	11	2	1	50.00%	6
2	25	5	2	40.00%	10
3	34	7	2	28.57%	10
4	18	4	2	50.00%	9
5	16	3	1	33.33%	5
6	40	8	3	37.50%	15
7	31	6	2	33.33%	10
8	50	10	4	40.00%	20
9	14	3	1	33.33%	5
10	12	2	1	50.00%	6
TOTALS	251	50	19		
OVERALL TOTALS					
80	3,500			21.84%	764
STANDARD ERROR				3.57%	125
CONFIDENCE LEVEL		80 PERCENT	90 PERCENT	95 PERCENT	
LOWER LIMIT FOR PROPORTION		17.27%	15.97%	14.85%	
UPPER LIMIT FOR PROPORTION		26.41%	27.70%	28.83%	
LOWER LIMIT FOR TOTAL		604	559	520	
UPPER LIMIT FOR TOTAL		924	970	1,009	

The values used as input to the **Stratified Multistage** program are highlighted in the preceding computer output. The following computer screen illustrates how to enter these values:

Attribute - Stratified Multistage

Name of Audit/Review

Number of strata for this appraisal Have you created a data file for this appraisal? ☐ Yes ☒ No

NOTE: Stratum ratios and standard errors are entered as percentages (e.g., enter 5 for 5% -- not .05).

Stratum	Ratio	Standard Error	Universe Size
1	16.9	1.99	5600
2	21.84	3.57	3500

Click here after entering or editing data.

HELP

Main Menu

EXIT

OUTPUT TO

☐ Text File and Screen

☐ Printer and Screen

☐ Text File, Printer, and Screen

☒ Screen Only

The following output is obtained from the **Stratified Multistage** program.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/25/2010 STRATIFIED MULTISTAGE ATTRIBUTE APPRAISAL Time: 16:08
 AUDIT/REVIEW: Illustration

THE ESTIMATORS ARE BASED ON THE FOLLOWING ENTRIES:

STRATUM	ATTRIBUTE	STANDARD ERROR	UNIVERSE SIZE
1	16.90%	1.99%	5,600
2	21.84%	3.57%	3,500

= = = = = RESULTS = = = = =

ESTIMATED PERCENTAGE: 18.80%
 ESTIMATED TOTAL: 1,711

STANDARD ERROR (PERCENTAGE): 1.84%
 STANDARD ERROR (TOTAL): 167

CONFIDENCE LEVEL	80 PERCENT	90 PERCENT	95 PERCENT
LOWER LIMIT FOR PROPORTION	16.44%	15.77%	15.19%
UPPER LIMIT FOR PROPORTION	21.16%	21.83%	22.41%

LOWER LIMIT FOR TOTAL	1,496	1,435	1,383
UPPER LIMIT FOR TOTAL	1,925	1,986	2,039

Final results: The point estimate for the percentage of grants containing improper charges is 18.8% (standard error of 1.84%) and the 90% confidence interval for this proportion is from 15.77% to 21.83%. The 90% confidence interval for the universe total is from 1,435 to 1,986.

Discussion. Using Equation 1 in the Formulas section,

$$\hat{p} = (5600/9100)(.169) + (3500/9100)(.2184) = .188 \text{ (18.8\%)}$$

and

$$v(\hat{p}) = (5600/9100)^2(.0199)^2 + (3500/9100)^2(.0357)^2 = .000339$$

The estimated standard error of \hat{p} is $\sqrt{v(\hat{p})} = .0184 \text{ (1.84\%)}$.

The corresponding 90% confidence interval is

$$18.8 \pm 1.644853626951(1.84); \text{ that is } 15.77\% \text{ to } 21.83\%$$

The estimate of the universe total and corresponding confidence interval are obtained by multiplying the previous results by the total universe size = 5,600 + 3,500 = 9,100.

FORMULAS

1. Estimated universe proportion having the attribute of interest

$$\hat{p} = \sum_{i=1}^L \left(\frac{M_i}{M} \right) \hat{p}_i$$

where L = number of strata

M_i = universe size for the most detailed level of the multistage sample

M = total universe size = $\sum m_i$

\hat{p}_i = estimated proportion for the i-th stratum

2. Estimated variance of \hat{p}

$$v(\hat{p}) = \sum_{i=1}^L \left(\frac{M_i}{M} \right)^2 (\text{standard error of } \hat{p}_i)^2$$

3. 90% confidence interval for p

$$\hat{p} \pm 1.644853626951 \sqrt{v(\hat{p})}$$

4. Estimated universe total having the attribute of interest

$$\hat{T} = M\hat{p}$$

5. Estimated variance of \hat{T}

$$v(\hat{T}) = M^2 v(\hat{p})$$

6. 90% confidence interval for T

$$\hat{T} \pm 1.644853626951 \sqrt{v(\hat{T})}$$

NOTE: For the Precision at the 95% Confidence Level, replace 1.644853626951 with 1.959963984540 and for the Precision at the 80% Confidence Level, replace 1.644853626951 with 1.281551565545.

VARIABLE APPRAISALS

A variable appraisal is carried out to estimate a particular universe total (T) and its corresponding sampling error. For example, the audit intent may be to determine the dollar value of an inventory or the amount of duplicate payments made by an organization.

A variety of procedures can be used to obtain and appraise a variable sample. There are ten sampling strategies utilized in the Variables Appraisals modules. They are listed below and described in the sections to follow.

- Unrestricted
- Stratified
- Two-Stage Unrestricted
- Three-Stage Unrestricted
- RHC Two Stage
- RHC Three Stage
- Stratified Cluster
- Stratified Multistage
- Post Stratification
- Unknown Universe Size

Unrestricted Variable Appraisal

An **unrestricted** sample is the same as a **simple random sample**. Consequently, every sample of size n has the same chance of being selected. For an unrestricted sample, a sample of size n is randomly obtained and the variable of interest is recorded for each sample item. Actually, the user may input a set of single values (examined amounts, audit amounts, or difference amounts) or a set of two values (examined and audit amounts, examined and difference amounts, or audit and difference amounts).

Example 1. An unrestricted sample of 50 items resulted in the 50 examined/audited values contained in data set DATASRS.TXT. For this sample, all of the resulting differences (examined value - audited value) were nonzero since all the examined (book) values were unequal to the corresponding audit (actual) values.

Data file DATASRS.TXT

1	300	267	◀ Each line contains a line counter, examined value, and audited value separated by one or more spaces, a tab delimiter, or a comma.
2	900	774	
3	300	255	
4	200	174	
5	900	810	
6	700	560	
7	1000	820	
8	100	80	
9	900	765	
10	700	630	
11	700	630	
12	400	332	
13	300	255	
14	100	84	
15	200	168	
16	100	88	
17	600	528	
18	400	340	

19	900	747
20	1000	800
21	1000	862
22	600	504
23	800	648
24	200	176
25	200	172
26	1000	890
27	900	792
28	600	540
29	500	525
30	200	172
31	200	178
32	500	425
33	200	164
34	500	420
35	500	400
36	400	324
37	200	160
38	600	540
39	500	425
40	300	264
41	900	765
42	100	84
43	100	85
44	900	810
45	300	240
46	500	415
47	500	425
48	300	237
49	500	435
50	100	86

The output on the next page was obtained from the **Unrestricted Variable Appraisal** program.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/5/2010 VARIABLE UNRESTRICTED APPRAISAL Time: 11:23
 AUDIT/REVIEW: Variable SRS
 DATA FILE USED: C:\Temp\DATASRS.TXT

SAMPLE SIZE	EXAMINED VALUE	NONZERO DIFFS	TOTAL OF DIFF VALUES	TOTAL OF AUD VALUES
50	24,800.00	50	3,530.00	21,270.00

----- E X A M I N E D -----

MEAN / UNIVERSE	496.00	10,000
STANDARD DEVIATION	296.90	
SKEWNESS	.32	
KURTOSIS	1.81	
STANDARD ERROR (MEAN)	41.88	
STANDARD ERROR (TOTAL)	418,823	
POINT ESTIMATE	4,960,000	

CONFIDENCE LIMITS

80% CONFIDENCE LEVEL

LOWER LIMIT	4,415,921
UPPER LIMIT	5,504,079
PRECISION AMOUNT	544,079
PRECISION PERCENT	10.97%
T-VALUE USED	1.299068784748

90% CONFIDENCE LEVEL

LOWER LIMIT	4,257,823
UPPER LIMIT	5,662,177
PRECISION AMOUNT	702,177
PRECISION PERCENT	14.16%
T-VALUE USED	1.676550892617

95% CONFIDENCE LEVEL

LOWER LIMIT	4,118,344
UPPER LIMIT	5,801,656
PRECISION AMOUNT	841,656
PRECISION PERCENT	16.97%
T-VALUE USED	2.009575237129

----- A U D I T E D -----

MEAN / UNIVERSE	425.40	10,000
STANDARD DEVIATION	256.20	
SKEWNESS	.30	
KURTOSIS	1.78	
STANDARD ERROR (MEAN)	36.14	
STANDARD ERROR (TOTAL)	361,412	
POINT ESTIMATE	4,254,000	

CONFIDENCE LIMITS

80% CONFIDENCE LEVEL

LOWER LIMIT	3,784,500
UPPER LIMIT	4,723,500
PRECISION AMOUNT	469,500
PRECISION PERCENT	11.04%
T-VALUE USED	1.299068784748

	90% CONFIDENCE LEVEL
LOWER LIMIT	3,648,074
UPPER LIMIT	4,859,926
PRECISION AMOUNT	605,926
PRECISION PERCENT	14.24%
T-VALUE USED	1.676550892617

	95% CONFIDENCE LEVEL
LOWER LIMIT	3,527,715
UPPER LIMIT	4,980,285
PRECISION AMOUNT	726,285
PRECISION PERCENT	17.07%
T-VALUE USED	2.009575237129

----- D I F F E R E N C E -----		
MEAN / UNIVERSE	70.60	10,000
STANDARD DEVIATION	48.25	
SKEWNESS	.64	
KURTOSIS	2.98	
STANDARD ERROR (MEAN)	6.81	
STANDARD ERROR (TOTAL)	68,068	
POINT ESTIMATE	706,000	

	CONFIDENCE LIMITS
	80% CONFIDENCE LEVEL
LOWER LIMIT	617,575
UPPER LIMIT	794,425
PRECISION AMOUNT	88,425
PRECISION PERCENT	12.52%
T-VALUE USED	1.299068784748

	90% CONFIDENCE LEVEL
LOWER LIMIT	591,881
UPPER LIMIT	820,119
PRECISION AMOUNT	114,119
PRECISION PERCENT	16.16%
T-VALUE USED	1.676550892617

	95% CONFIDENCE LEVEL
LOWER LIMIT	569,213
UPPER LIMIT	842,787
PRECISION AMOUNT	136,787
PRECISION PERCENT	19.37%
T-VALUE USED	2.009575237129

Explanation.

NOTE: The following discussion can be applied to the examined values, the audited values, or the difference values. The difference values will be used when discussing the computer output.

The estimated mean of the difference amounts in the universe is the sample mean, $\bar{x} = \$70.60$.

The estimated total difference for the universe (T) is the sample mean times the universe size;

that is, $\hat{T} = (70.60)(10,000) = \$706,000$. This is referred to as the POINT ESTIMATE in the computer output.

The sample standard deviation is $s = 48.2519$ and the corresponding (estimated) standard error for the mean is $48.2519 \sqrt{\frac{10000 - 50}{(50)(10000)}} = 6.80677$. The (estimated) standard error for the total is $10,000 \times 6.80677 = 68,067.7$ (68,068 rounded). The sample skewness is a measure of the symmetry of the sample data. This value is SKEWNESS = 0.64, indicating a very slight positive (right-tail) skew. The sample kurtosis is a measure of the sample “peakedness” and is equal to KURTOSIS = 2.98. Essentially, this value is small whenever the frequency of observations close to the mean is high and the frequency of observations far from the mean is low.

The 95% confidence interval for the universe total of the difference amounts is $706,000 \pm (2.009575237129)(68,067.7) = 706,000 \pm 136,787$; that is, 569,213 to 842,787. The

PRECISION AMOUNT is the amount added and subtracted to the POINT ESTIMATE; that is, \$136,787. This value is 19.37% of the point estimate and is referred to as the PRECISION PERCENT.¹

¹ When the POINT ESTIMATE is negative, the PRECISION PERCENT is set equal to zero.

FORMULAS

$$\text{STANDARD DEVIATION} = s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n-1}}$$

$$\text{STANDARD ERROR (MEAN)} = s\sqrt{\frac{N-n}{nN}} \text{ and } \text{STANDARD ERROR (TOTAL)} = Ns\sqrt{\frac{N-n}{nN}}$$

where n = sample size, N = universe size.

$$\text{SKEWNESS} = \frac{\frac{1}{n} \sum_{i=1}^n (x - \bar{x})^3}{\left[\frac{1}{n} \sum_{i=1}^n (x - \bar{x})^2 \right]^{3/2}}$$

$$\text{KURTOSIS} = \frac{\frac{1}{n} \sum_{i=1}^n (x - \bar{x})^4}{\left[\frac{1}{n} \sum_{i=1}^n (x - \bar{x})^2 \right]^2}$$

95% confidence interval for the universe total (T)

$$\hat{T} \pm t_{.025, n-1} \cdot s \cdot \sqrt{\frac{N(N-n)}{n}}$$

where (1) $\hat{T} = \bar{x} \cdot N$

(2) $t_{.025, n-1}$ is the t-value with n - 1 df having a right-tail area = .025 (RAT-STATS provides t-values accurate to 12 decimal places).

NOTE: For a 90% confidence interval, replace $t_{.025, n-1}$ with $t_{.05, n-1}$ and for an 80% confidence interval, replace $t_{.025, n-1}$ with $t_{.10, n-1}$.

Stratified Variable Appraisal

In a stratified variable sampling plan, the universe is divided into two or more nonoverlapping categories (strata). As with an unrestricted sample, the intent is to make a statistical estimate for a universe total (T) for a particular variable of interest. This plan involves obtaining a random sample from each of the strata. The program will request the number of universe items in each stratum and these values must be known. The program will develop estimates for each stratum as well as for the entire universe.

Using a Stratified Sample

Purpose: To divide (partition) the universe into separate strata so that variation within individual strata is less than variation within the entire universe.

Simple Illustration:

Universe consists of {5 7 8 10 55 60 66 70 120 133 145 150}

Mean of universe is $\mu = 69.08$

Variance of universe is $\sigma^2 = 2871.9$

Partition the universe into three strata:

{5 7 8 10}	{55 60 66 70}	{120 133 145 150}
#1	#2	#3

The strata variances are: Stratum Variance

1	3.25	◀ Compare these to $\sigma^2 = 2871.9$
2	32.69	
3	134.50	

Consequently, the individual strata are much more homogeneous.

So, when obtaining a stratified sample, the user can take a larger sample (perhaps 100%) from the stratum containing the large dollar items.

Reasons for Using Stratified Sampling:**A. Improved Sampling Precision**

Stratification tends to make the sampling more efficient; that is, the user will obtain narrower confidence intervals for the same sample size. When a sample is skewed or has a high degree of variability, the sample size required to provide a reasonable degree of precision using simple random sampling may be quite large. Precision is improved because each stratum should have a relatively small variance and the weighted sum of the strata variances is less than the variance for the entire universe.

B. Separate Information About Strata and the Universe

Strata may be formed because separate estimates are desired for subuniverses. For example, a nationwide audit of nursing homes can be planned in advance such that separate estimates are published for each state (stratum). When an auditor selects a simple random sample from the entire universe, he/she cannot control the sample size within each stratum. Stratified sampling permits the auditor to also impose different precision requirements on different strata, such as requiring more precise estimates for large accounts.

C. Accommodation of Different Techniques

It may be desirable to employ different sampling methods or audit techniques in various portions of the universe. For example, in a sample of health service employees, the headquarters employees (Stratum 1) may be sampled as individuals and the employees scattered throughout the state (Stratum 2) may be sampled as clusters to save travel time and cost.

Comments:

- (1) Defining effective strata is no accident! The user can incorporate all sorts of prior knowledge in defining the strata. Such a technique does not introduce any bias into the final estimate since strata are defined prior to obtaining the sample and each sampling item has a known (although not the same) chance of being selected. As a result, a well-designed stratified plan can provide audit protection and/or improved precision.
- (2) Strata can be defined after sample data are obtained provided the proportions of the universe in each stratum are known (with negligible error) and samples of at least 20 are obtained from each stratum.
- (3) Generally, it is not a good idea to stratify for convenience (unlike cluster or multistage sampling) since the resulting estimator may be less efficient than the estimator which uses a single simple random sample.
- (4) Even though random selection is performed within strata, this does not mean that the user cannot take a close look at the individual findings to determine nature, source, cause, trend and impact.
- (5) A careful balance must be maintained between the gains expected in sample precision and the additional time and resources involved in introducing a stratified scheme into the sample design.

Strata Formation

Strata are typically defined using the dollar value of the items being sampled. An alternative is to stratify using some other variable which is highly correlated with the principal variable, such as using the number of hospital beds to measure the “size” of a hospital.

Basic Rule: Select strata so that their means are as different as possible and their standard deviations are as small as possible.

Guidelines:

- A few strata yield most of the gains (say, 2 to 6).
- Experience, intuition, and the judgment of the auditor are extremely useful in improving the sampling precision through effective stratification.
- Quantitative rather than qualitative (sex, race, etc.) variables are preferable for defining strata.
- Coarser divisions of several stratifying variables are preferable to finer divisions of one variable.
- It is better to use unrelated stratifying variables.

Example 2. Random samples of size 25 were obtained from two strata:

Stratum 1: Examined amounts under \$200 ($N_1 = 5,200$) and

Stratum 2: Examined amounts \geq \$200 ($N_2 = 3,500$)

NOTE: These sample sizes are too small to meet OAS standards and are used for illustrative purposes only.

The sample difference amounts for the two strata are stored in data file DATASTRAT.TXT and the universe/sample sizes are stored in file UNIVSTRAT.TXT.

Data set DATASTRAT.TXT

1	80
2	43
3	133
4	125
5	116
6	84
7	111
8	148
9	104
10	114
11	83
12	132
13	96
14	86
15	66
16	89
17	72
18	114
19	135
20	71
21	127
22	105
23	102
24	69
25	76
26	354
27	328
28	313
29	250
30	261
31	294
32	380
33	296
34	248
35	277
36	331
37	305
38	360
39	348
40	318
41	290
42	249
43	362
44	348
45	355
46	295
47	277
48	355
49	314
50	277

Universe File UNIVSTRAT.TXT

1	5200	25
2	3500	25

The sample results are:

Stratum 1: $n_1 = 25$, mean = 99.24, std. dev. = 26.3317

Stratum 2: $n_2 = 25$, mean = 311.40, std. dev. = 39.6432

The following computer output was obtained from the **Stratified Variables Appraisal** program:

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 4/5/2010      STRATIFIED VARIABLE APPRAISAL      Time: 12:17
                    AUDIT/REVIEW: Variable - Stratified

DATA FILE USED: C:\Temp\DATASTRAT.TXT

STRATUM      SAMPLE
NUMBER      SIZE      VALUE OF SAMPLE      NONZERO ITEMS
1           25           2,481.00           25
2           25           7,785.00           25
TOTALS      50           10,266.00           50

----- D I F F E R E N C E -----
Stratum 1  MEAN / UNIVERSE           99.24           5,200
            STANDARD DEVIATION       26.33
            SKEWNESS                 -.07
            KURTOSIS                 2.24
            STANDARD ERROR (MEAN)     5.25
            STANDARD ERROR (TOTAL)    27,319
            POINT ESTIMATE            516,048

                                CONFIDENCE LIMITS
                                80% CONFIDENCE LEVEL
LOWER LIMIT           480,046
UPPER LIMIT           552,050
PRECISION AMOUNT      36,002
PRECISION PERCENT     6.98%
T-VALUE USED          1.317835933673

                                90% CONFIDENCE LEVEL
LOWER LIMIT           469,308
UPPER LIMIT           562,788
PRECISION AMOUNT      46,740
PRECISION PERCENT     9.06%
T-VALUE USED          1.710882079909

                                95% CONFIDENCE LEVEL
LOWER LIMIT           459,664
UPPER LIMIT           572,432
PRECISION AMOUNT      56,384
PRECISION PERCENT     10.93%
T-VALUE USED          2.063898561628

```

Stratum 2	MEAN / UNIVERSE	311.40	3,500
	STANDARD DEVIATION	39.64	
	SKEWNESS	-.06	
	KURTOSIS	1.85	
	STANDARD ERROR (MEAN)	7.90	
	STANDARD ERROR (TOTAL)	27,651	
	POINT ESTIMATE	1,089,900	
	CONFIDENCE LIMITS		
	80% CONFIDENCE LEVEL		
	LOWER LIMIT	1,053,461	
	UPPER LIMIT	1,126,339	
	PRECISION AMOUNT	36,439	
	PRECISION PERCENT	3.34%	
	T-VALUE USED	1.317835933673	
	90% CONFIDENCE LEVEL		
	LOWER LIMIT	1,042,592	
	UPPER LIMIT	1,137,208	
	PRECISION AMOUNT	47,308	
	PRECISION PERCENT	4.34%	
	T-VALUE USED	1.710882079909	
	95% CONFIDENCE LEVEL		
	LOWER LIMIT	1,032,831	
	UPPER LIMIT	1,146,969	
	PRECISION AMOUNT	57,069	
	PRECISION PERCENT	5.24%	
	T-VALUE USED	2.063898561628	
OVERALL	POINT ESTIMATE / UNIVERSE	1,605,948	8,700
	STANDARD ERROR	38,870	
	CONFIDENCE LIMITS		
	80% CONFIDENCE LEVEL		
	LOWER LIMIT	1,556,134	
	UPPER LIMIT	1,655,762	
	PRECISION AMOUNT	49,814	
	PRECISION PERCENT	3.10%	
	Z-VALUE USED	1.281551565545	
	90% CONFIDENCE LEVEL		
	LOWER LIMIT	1,542,012	
	UPPER LIMIT	1,669,884	
	PRECISION AMOUNT	63,936	
	PRECISION PERCENT	3.98%	
	Z-VALUE USED	1.644853626951	
	95% CONFIDENCE LEVEL		
	LOWER LIMIT	1,529,764	
	UPPER LIMIT	1,682,132	
	PRECISION AMOUNT	76,184	
	PRECISION PERCENT	4.74%	
	Z-VALUE USED	1.959963984540	

Discussion. The point estimates for the universe total difference amounts are \$516,048 (Stratum 1) and \$1,089,900 (Stratum 2). Referring to the formula section and the OVERALL section in the computer output, the estimate of the universe total difference is

$$\hat{T} = (5200)(99.24) + (3500)(311.40) = \$1,605,948$$

The estimated variance of \hat{T} is

$$5200^2 \left(\frac{5200 - 25}{5200} \right) \frac{26.3317^2}{25} + 3500^2 \left(\frac{3500 - 25}{3500} \right) \frac{39.6432^2}{25} = 1,510,906,287$$

The (estimated) standard error of \hat{T} is $\sqrt{1,510,906,287} = 38,870$.

The 95% confidence interval for universe total (T) is

$$1,605,948 \pm (1.959963984540)(38,870)$$

that is, $1,605,948 \pm 76,184$ (\$1,529,764 to \$1,682,132).

The PRECISION AMOUNT here is \$76,184 and is 4.74% of the point estimate, \hat{T} .

NOTE: When the POINT ESTIMATE is negative, the PRECISION PERCENT is set equal to zero.

FORMULAS

NOTE: For definitions and formulas of the statistics within each stratum (standard deviation, standard error, skewness, and kurtosis), refer to the previous section (**Unrestricted Variable Appraisal**).

1. Estimate of universe mean (μ):

$$\bar{y}_{st} = (N_1 / N)\bar{y}_1 + (N_2 / N)\bar{y}_2 + \cdots + (N_L / N)\bar{y}_L$$

where L = number of strata

N_i = number of items in i -th stratum (universe)

$$N = N_1 + N_2 + \cdots + N_L$$

\bar{y}_i = average of sample items in the i-th stratum

2. Estimate of universe total (T):

$$\hat{T} = N \cdot \bar{y}_{st} = N_1 \cdot \bar{y}_1 + N_2 \cdot \bar{y}_2 + \cdots + N_L \cdot \bar{y}_L$$

3. Estimated variance of \bar{y}_{st} :

$$v(\bar{y}_{st}) = \frac{1}{N^2} \sum_{i=1}^L N_i^2 \left(\frac{N_i - n_i}{N_i} \right) \frac{s_i^2}{n_i}$$

where n_i = number of sampled items in i-th stratum

s_i^2 = sample variance for i-th stratum

4. Estimated variance of \hat{T} :

$$v(\hat{T}) = N^2 v(\bar{y}_{st})$$

5. Approximate 95% confidence interval for universe mean (μ):

$$\bar{y}_{st} \pm Z_{.025} \sqrt{v(\bar{y}_{st})} \quad \text{where } Z_{.025} = 1.959963984540.$$

6. Approximate 95% confidence interval for universe total (T):

$$\hat{T} \pm Z_{.025} \sqrt{v(\hat{T})}$$

- NOTES:**
1. For a 90% confidence interval, replace $Z_{.025}$ with $Z_{.05} = 1.644853626951$ and for an 80% confidence interval, replace $Z_{.025}$ with $Z_{.10} = 1.281551565545$.
 2. The confidence intervals for each stratum total use t-values that are accurate to 12 decimal places.

Two-Stage Unrestricted Variable Appraisal

This is a special case of **multistage sampling**. Multistage sampling is a very cost-effective sampling procedure when (1) obtaining a frame that lists all elements in the universe is very costly or impossible or (2) the cost of obtaining observations increases as the distance separating the elements increases. Put another way, multistage sampling is cost effective when it is more costly to get to the sampling unit than it is to audit the sampling unit. The goal of multistage sampling is to get the most precise results per unit of examination cost.

General Comments

1. This is a very convenient sampling procedure for many situations. You don't have to visit all locations.
2. For a two-stage procedure, the universe can be broken down into “subgroups.”

Example: 1st Stage: Carriers (primary units, P.U.s)

 2nd Stage: Hospitals (secondary units, S.U.s)

So, the procedure is to first obtain a random sample of P.U.s. These are called clusters.

Then, obtain a random sample of S.U.s within each selected P.U. Notice that at the first stage, clusters are the sampling unit (sampling units are not always individual people, records, etc.). The program will accept a maximum of 20 clusters.

3. You can estimate cost recoveries for the entire universe with multistage sampling and it is very useful for large, widespread universes.

Example 3. In a particular region of the U.S. there are $N = 90$ universities with government research grants. Because these universities are so widespread, it was decided to use a cluster

sample of $n = 10$ universities. The 10 universities to be sampled may be obtained using the **Single-Stage Random Numbers** module discussed in the **Random Numbers** section.

Enter the values shown in the following input screen:

Single Stage Random Numbers

Do you want to enter a seed number? ☐ no ☒ yes

Enter the seed number below: 1357

Name of the audit/review: select

Enter the quantity of numbers to be generated in:

Sequential Order: 10 Spares in Random Order: 0

The sampling frame:

Low Number: 1 High Number: 90

HELP

Main Menu

EXIT

OUTPUT TO

☐ Printer

☒ Text File

☐ Access File

☐ Excel File

☐ Flat File

Click on File Name(s) when the desired output formats have been checked in the OUTPUT TO box. File Name(s)

CONTINUE

The resulting output is shown on the next page. Note that the selected universities are in sequential order:

Universities: 2, 5, 7, 23, 28, 46, 56, 67, 70, 76

Department of Health and Human Services
 OIG - Office of Audit Services
 Date: 5/26/2010 Random Number Generator Time: 11:03
 AUDIT: select

SEED NUMBER: 1357.00 FRAME SIZE: 90

FILE OF RANDOM NUMBERS: C:\TEMP\SELECT.TXT

TOTAL RANDOM NUMBERS GENERATED: 10

THE NUMBERS ARE IN THE FOLLOWING FORMAT IN YOUR FILE:

POSITIONS 1 THROUGH 6 - ORDER OF SELECTION

POSITIONS 7 THROUGH 17 - RANDOM NUMBER

EACH COLUMN OF NUMBERS IS RIGHT JUSTIFIED.

Selection Order	Value
4	2
7	5
9	7
10	23
5	28
2	46
6	56
1	67
8	70
3	76

SUMMATION OF RANDOM NUMBERS = 380

Example 3 -- continued. Rather than audit all grants at a selected university, it was decided (based on available resources) to audit roughly 20% of the grants at each selected university.

There are a total of $M = 4500$ grants in all 90 universities (see NOTE 5 under Formula 2). The following data were obtained where y_i is the dollars (in thousands) of improper charges for the i th

sampld university, M_i is the total number of grants at the i -th university, and m_i is the number of audited grants at the i -th university. Also, \bar{y}_i and s_i^2 are the mean and variance of the sample values from the i -th university.

For ease of illustration, university 1 refers to university 2, university 2 refers to university 5, and so on.

Univ.	M_i	m_i	Dollars (y_i , in thousands)	\bar{y}_i	s_i^2
1	50	10	5, 7, 9, 0, 11, 2, 8, 4, 3, 5	5.40	11.38
2	65	13	4, 3, 7, 2, 11, 0, 1, 9, 4, 3, 2, 1, 5	4.00	10.67
3	45	9	5, 6, 4, 11, 12, 0, 1, 8, 4	5.67	16.75
4	48	10	6, 4, 0, 1, 0, 9, 8, 4, 6, 10	4.80	13.29
5	52	10	11, 4, 3, 1, 0, 2, 8, 6, 5, 3	4.30	11.12
6	58	12	12, 11, 3, 4, 2, 0, 0, 1, 4, 3, 2, 4	3.83	14.88
7	42	8	3, 7, 6, 7, 8, 4, 3, 2	5.00	5.14
8	66	13	3, 6, 4, 3, 2, 2, 8, 4, 0, 4, 5, 6, 3	3.85	4.31
9	40	8	6, 4, 7, 3, 9, 1, 4, 5	4.88	6.13
10	56	11	6, 7, 5, 10, 11, 2, 1, 4, 0, 5, 4	5.00	11.80

NOTE: This example violates OAS minimum sample sizes of at least 30 grants at each university. It is used for illustration only. These data are in data set DATA2STG.TXT. The universe sizes (the M_i 's) are in data set UNIV2STG.TXT. The program output immediately follows.

<u>Dataset DATA2STG.TXT</u>		<u>Dataset UNIV2STG.TXT</u>		
1	5	1	50	10
2	7	2	65	13
3	9	3	45	9
4	0	4	48	10
5	11	5	52	10
6	2	6	58	12
7	8	7	42	8
8	4	8	66	13
9	3	9	40	8
10	5	10	56	11
11	4			
12	3			
13	7			
14	2			
15	11			
16	0			
17	1			
18	9			
19	4			
20	3			
21	2			
22	1			

23	5
.	
.	
.	
94	6
95	7
96	5
97	10
98	11
99	2
100	1
101	4
102	0
103	5
104	4

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/26/2010

TWO-STAGE UNRESTRICTED VARIABLE APPRAISAL

Time: 10:07

AUDIT/REVIEW: Variable 2-Stage

DATA FILE USED: C:\Temp\DATA2STG.TXT

----- D I F F E R E N C E -----						
UNIT NBR	SAMPLE SIZE/ NONZERO ITEMS	SAMPLE MEAN	VARIANCE	UNIVERSE SIZE	POINT ESTIMATE	
1	10/9	5.40	11.38	50	270	
2	13/12	4.00	10.67	65	260	
3	9/8	5.67	16.75	45	255	
4	10/8	4.80	13.29	48	230	
5	10/9	4.30	11.12	52	224	
6	12/10	3.83	14.88	58	222	
7	8/8	5.00	5.14	42	210	
8	13/12	3.85	4.31	66	254	
9	8/8	4.88	6.13	40	195	
10	11/10	5.00	11.80	56	280	
	104/94	4.80		522	2,400	
NOT SAMPLED						
	80			3,978		
OVERALL TOTALS						
	90			4,500	21,602	
STANDARD ERROR						867
CONFIDENCE LIMITS						
80% CONFIDENCE LEVEL						
	LOWER LIMIT			20,491		
	UPPER LIMIT			22,712		
	PRECISION AMOUNT			1,111		
	PRECISION PERCENT			5.14%		
	Z-VALUE USED			1.281551565545		

90% CONFIDENCE LEVEL	
LOWER LIMIT	20,176
UPPER LIMIT	23,027
PRECISION AMOUNT	1,425
PRECISION PERCENT	6.60%
Z-VALUE USED	1.644853626951
95% CONFIDENCE LEVEL	
LOWER LIMIT	19,903
UPPER LIMIT	23,300
PRECISION AMOUNT	1,699
PRECISION PERCENT	7.86%
Z-VALUE USED	1.959963984540

Discussion. The point estimate (highlighted) for the universe total difference amount is 21,602 (that is, \$21,602,000). The 95% confidence interval for this amount is from 19,903 to 23,300 (\$19,903,000 to \$23,300,000). The PRECISION AMOUNT at the 95% confidence level is (Z-value)(standard error of \hat{T}) = (1.959963984540)(867) = 1,699 (that is, \$1,699,000). This value is 7.86% of the point estimate.

Notice that the output also contains the estimated totals for each primary unit (university). For example, the estimated total difference for university 1 is 270 (\$270,000). The sample average of these estimates is $2,400/10 = 240$ (\$240,000). Since there are 90 universities in the universe, the point estimate for the universe total is $(90)(240) = 21,600$; that is, \$21,600,000 (more precisely, 21,602,000).

FORMULAS

1. The point estimate for the universe total (T) is

$$\hat{T} = N \frac{\sum_{i=1}^n M_i \bar{y}_i}{n}$$

2. The estimated variance of \hat{T} is

$$v(\hat{T}) = \left(\frac{N(N-n)}{n} \right) \left(\frac{\sum_{i=1}^n \left(M_i \bar{y}_i - \frac{\hat{T}}{N} \right)^2}{n-1} \right) + \frac{N}{n} \sum_{i=1}^n M_i^2 \left(\frac{M_i - m_i}{M_i} \right) \frac{s_i^2}{m_i}$$

NOTES: 1. n = number of primary units in the sample and N = number of primary units in the universe.

2. The STANDARD ERROR of \hat{T} is the square root of $v(\hat{T})$.

3. The PRECISION AMOUNT at the 95% confidence level for the universe total is (1.959963984540)(standard error of \hat{T}).

4. For the PRECISION AMOUNT at the 90% confidence level, replace 1.959963984540 with 1.644853626951. For the PRECISION AMOUNT at the 80% confidence level, replace 1.959963984540 with 1.281551565545.

5. The total number of secondary units in the universe (M) may be known or unknown and is not used in any of the calculations.

3. The approximate 95% confidence interval for T is

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

Three-Stage Unrestricted Variable Appraisal

Example 4. The situation discussed in Example 3 was extended the following year to a three-stage procedure by defining:

Stage 1: REGION (select 4 out of 12 regions)

Stage 2: UNIVERSITY (select 10 from each selected region)

Stage 3: GRANT (select approximately 20% of all grants at each university)

NOTE: This example violates OAS minimum sample sizes and is used for illustration only.

Using the **Single-Stage Random Numbers** module, regions 5, 7, 8, and 10 were selected as the sampled primary units. Next, 10 universities (secondary units) were randomly selected from the available universities in each of the four selected regions. The data on the following four pages were obtained, where M_i is the number of grants in the universe for each university, m_i is the number of sampled grants at each university (chosen to be roughly 20% of M_i),

\bar{y}_{ij} is the sample average of the items from the j -th university within the i -th region, and s_{ij} is the sample standard deviation of the items from the j -th university within the i -th region. The resulting data are stored in file DATA3STG.TXT.

REGION 5 -- Universe contains 90 universities.

Univ.	M_i	m_i
1	47	9
2	51	10
3	45	9
4	46	9
5	46	9
6	50	10
7	50	10
8	57	11
9	54	11
10	64	13

Data (total thousands of dollars of improper charges)

Obs.	University									
	1	2	3	4	5	6	7	8	9	10
1	8	13	10	11	14	5	0	0	2	12
2	0	13	0	9	14	11	14	3	9	13
3	6	4	12	12	5	0	4	0	15	14
4	6	6	14	1	1	4	10	6	4	11
5	0	0	13	10	2	13	13	1	13	6
6	13	15	13	11	11	8	10	0	12	10
7	1	12	13	15	14	15	8	3	14	11
8	7	9	2	7	15	2	0	13	6	0
9	2	0	9	8	11	0	7	7	11	7
10		13				14	3	5	0	12
11								4	1	9
12										11
13										7
\bar{y}_{1j}	4.78	8.50	9.56	9.33	9.67	7.20	6.90	3.82	7.91	9.46
s_{1j}	4.38	5.64	5.13	3.91	5.52	5.79	5.02	3.92	5.52	3.73

REGION 7 -- Universe contains 110 universities.

Univ.	M_i	m_i
1	53	11
2	59	12
3	52	10
4	67	13
5	59	12
6	73	15
7	51	10
8	75	15
9	66	13
10	58	12

Data (total thousands of dollars of improper charges)

Obs.	University									
	1	2	3	4	5	6	7	8	9	10
1	0	12	4	5	0	2	4	19	17	16
2	4	0	19	10	4	18	16	13	8	17
3	0	15	16	2	6	0	0	0	15	6
4	10	11	12	10	9	0	8	13	12	10
5	11	0	4	12	13	12	3	4	0	2
6	18	18	2	7	19	17	8	0	20	6
7	18	0	1	3	9	0	13	0	17	13
8	16	17	5	0	18	0	0	0	6	0
9	2	8	1	20	16	16	0	0	9	12

10	18	12	8	10	15	3	9	13	16	20
11	18	13		0	0	0		4	7	19
12		5		0	13	4		16	2	0
13				15		3		13	7	
14						6		0		
15						5		4		
\bar{y}_{2j}	10.45	9.25	7.20	7.23	10.17	5.73	6.10	6.60	10.46	10.08
s_{2j}	7.69	6.59	6.41	6.30	6.59	6.65	5.65	7.01	6.21	7.23

REGION 8 -- Universe contains 85 universities.

Univ.	M_i	m_i
1	45	9
2	39	8
3	43	9
4	34	7
5	54	11
6	54	11
7	34	7
8	59	12
9	49	10
10	43	9

Data (total thousands of dollars of improper charges)

Obs.	University									
	1	2	3	4	5	6	7	8	9	10
1	6	0	10	4	13	0	13	0	6	14
2	5	8	15	1	13	0	10	13	8	0
3	1	0	11	2	15	14	10	9	3	0
4	3	1	6	5	10	11	6	3	12	10
5	12	10	12	2	10	14	9	14	4	12
6	7	15	2	9	7	5	8	14	8	0
7	0	1	14	4	0	1	1	12	1	7
8	3	14	7		0	13		11	2	6
9	12		0		3	0		11	4	0
10					13	5		6	3	
11					1	10		15		
12								11		
\bar{y}_{3j}	5.44	6.12	8.56	3.86	7.73	6.64	8.14	9.92	5.10	5.44
s_{3j}	4.33	6.40	5.20	2.67	5.78	5.90	3.80	4.66	3.38	5.68

REGION 10 -- Universe contains 120 universities.

Univ.	M_i	m_i
1	59	12
2	68	14
3	57	11
4	72	14
5	70	14
6	73	15
7	83	17
8	89	18
9	73	15
10	77	15

Data (total thousands of dollars of improper charges)

Obs.	University									
	1	2	3	4	5	6	7	8	9	10
1	0	3	8	9	7	15	18	16	20	18
2	10	13	10	10	6	15	10	2	14	0
3	14	0	1	15	3	7	17	14	5	8
4	0	12	8	16	15	12	18	0	0	4
5	18	12	19	6	12	15	0	0	19	5
6	0	7	18	17	10	4	0	0	0	17
7	8	1	15	5	17	0	8	6	12	0
8	20	13	0	2	0	14	0	19	5	0
9	19	2	0	8	0	18	3	17	15	13
10	0	0	17	9	0	17	18	13	15	8
11	0	16	18	16	0	13	12	12	2	10
12	3	14		7	0	6	0	13	15	7
13		17		0	7	0	11	12	2	10
14		5		0	16	3	19	11	2	0
15						4	0	14	1	0
16							9	13		
17							6	0		
18								1		
\bar{y}_{4j}	7.67	8.21	10.36	8.57	6.64	9.53	8.76	9.06	8.47	6.67
s_{4j}	8.25	6.27	7.55	5.80	6.44	6.32	7.38	6.77	7.39	6.17

Using the file construction suggested in the User's Guide for this module, the primary unit file and secondary unit file could be constructed as shown below. In the secondary unit file, each line begins with a counter (1, 2, 3, ...), a value identifying the primary unit number in the second

column, and a value identifying the secondary unit number within each primary unit in the third column.

Data set PRIMARY3STG.TXT

1	REGION	5	90	10
2	REGION	7	110	10
3	REGION	8	85	10
4	REGION	10	120	10

Data set SECONDARY3STG.TXT

1	1	1	UNIV1	47	9
2	1	2	UNIV2	51	10
3	1	3	UNIV3	45	9
4	1	4	UNIV4	46	9
5	1	5	UNIV5	46	9
6	1	6	UNIV6	50	10
7	1	7	UNIV7	50	10
8	1	8	UNIV8	57	11
9	1	9	UNIV9	54	11
10	1	10	UNIV10	64	13
11	2	1	UNIV1	53	11
12	2	2	UNIV2	59	12
13	2	3	UNIV3	52	10
14	2	4	UNIV4	67	13
15	2	5	UNIV5	59	12
16	2	6	UNIV6	73	15
17	2	7	UNIV7	51	10
18	2	8	UNIV8	75	15
19	2	9	UNIV9	66	13
20	2	10	UNIV10	58	12
21	3	1	UNIV1	45	9
22	3	2	UNIV2	39	8
23	3	3	UNIV3	43	9
24	3	4	UNIV4	34	7
25	3	5	UNIV5	54	11
26	3	6	UNIV6	54	11
27	3	7	UNIV7	34	7
28	3	8	UNIV8	59	12
29	3	9	UNIV9	49	10
30	3	10	UNIV10	43	9
31	4	1	UNIV1	59	12
32	4	2	UNIV2	68	14
33	4	3	UNIV3	57	11
34	4	4	UNIV4	72	14
35	4	5	UNIV5	70	14
36	4	6	UNIV6	73	15
37	4	7	UNIV7	83	17
38	4	8	UNIV8	89	18
39	4	9	UNIV9	73	15
40	4	10	UNIV10	77	15

Once again using the file construction suggested in the User's Guide for this module, the data file could be constructed as shown below. The data lines for the first two universities in Region 5 and the last two universities in Region 10 are shown. Each line begins with a counter (1, 2, 3, ...), a value identifying the primary unit number in the second column, a value identifying the secondary unit number within each primary unit in the third column, and a value identifying the third-stage unit number within each sampled primary/secondary unit in the fourth column. The sample value appears in the fifth column.

Data set DATA3STG.TXT

1	1	1	1	8
2	1	1	2	0
3	1	1	3	6
4	1	1	4	6
5	1	1	5	0
6	1	1	6	13
7	1	1	7	1
8	1	1	8	7
9	1	1	9	2
10	1	2	1	13
11	1	2	2	13
12	1	2	3	4
13	1	2	4	6
14	1	2	5	0
15	1	2	6	15
16	1	2	7	12
17	1	2	8	9
18	1	2	9	0
19	1	2	10	13
.				
.				
.				
433	4	9	1	20
434	4	9	2	14
435	4	9	3	5
436	4	9	4	0
437	4	9	5	19
438	4	9	6	0
439	4	9	7	12
440	4	9	8	5
441	4	9	9	15

442	4	9	10	15
443	4	9	11	2
444	4	9	12	15
445	4	9	13	2
446	4	9	14	2
447	4	9	15	1
448	4	10	1	18
449	4	10	2	0
450	4	10	3	8
451	4	10	4	4
452	4	10	5	5
453	4	10	6	17
454	4	10	7	0
455	4	10	8	0
456	4	10	9	13
457	4	10	10	8
458	4	10	11	10
459	4	10	12	7
460	4	10	13	10
461	4	10	14	0
462	4	10	15	0

The program output using these three files is shown on the following pages.

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 3/4/2010  THREE-STAGE UNRESTRICTED VARIABLE APPRAISAL  Time: 17:34
AUDIT/REVIEW: Variable 3-Stage

DATA FILE USED: C:\Temp\DATA3STG.TXT

----- D I F F E R E N C E -----
FIRST STAGE      SAMPLE  NON-
SECOND STAGE     SIZE    ZEROES  SAMPLE MEAN  VARIANCE  UNIVERSE  POINT ESTIMATE
=====
REGION 5
UNIV1             9        7        4.78    19.19        47        225
UNIV2            10        8        8.50    31.83        51        434
UNIV3             9        8        9.56    26.28        45        430
UNIV4             9        9        9.33    15.25        46        429
UNIV5             9        9        9.67    30.50        46        445
UNIV6            10        8        7.20    33.51        50        360
UNIV7            10        8        6.90    25.21        50        345
UNIV8            11        8        3.82    15.36        57        218
UNIV9            11       10        7.91    30.49        54        427
UNIV10           13       12        9.46    13.94        64        606

COMBINED          101                392                90        35,256

```

REGION 7						
UNIV1	11	9	10.45	59.07	53	554
UNIV2	12	9	9.25	43.48	59	546
UNIV3	10	10	7.20	41.07	52	374
UNIV4	13	10	7.23	39.69	67	484
UNIV5	12	10	10.17	43.42	59	600
UNIV6	15	10	5.73	44.21	73	419
UNIV7	10	7	6.10	31.88	51	311
UNIV8	15	9	6.60	49.11	75	495
UNIV9	13	12	10.46	38.60	66	690
UNIV10	12	10	10.08	52.27	58	585
COMBINED	123		506		110	55,643

REGION 8						
UNIV1	9	8	5.44	18.78	45	245
UNIV2	8	6	6.12	40.98	39	239
UNIV3	9	8	8.56	27.03	43	368
UNIV4	7	7	3.86	7.14	34	131
UNIV5	11	9	7.73	33.42	54	417
UNIV6	11	8	6.64	34.85	54	358
UNIV7	7	7	8.14	14.48	34	277
UNIV8	12	11	9.92	21.72	59	585
UNIV9	10	10	5.10	11.43	49	250
UNIV10	9	5	5.44	32.28	43	234
COMBINED	93		310		85	26,388

----- D I F F E R E N C E -----						
FIRST STAGE SECOND STAGE	SAMPLE SIZE	NON- ZEROES	SAMPLE MEAN	VARIANCE	UNIVERSE	POINT ESTIMATE
=====						
REGION 10						
UNIV1	12	7	7.67	68.06	59	452
UNIV2	14	12	8.21	39.26	68	559
UNIV3	11	9	10.36	57.05	57	591
UNIV4	14	12	8.57	33.65	72	617
UNIV5	14	9	6.64	41.48	70	465
UNIV6	15	13	9.53	39.98	73	696
UNIV7	17	12	8.76	54.44	83	727
UNIV8	18	14	9.06	45.82	89	806
UNIV9	15	13	8.47	54.55	73	618
UNIV10	15	10	6.67	38.10	77	513
COMBINED	145		604		120	72,534

STAGES	UNIVERSE	SAMPLED
FIRST	12	4
SECOND	405	40
THIRD	2,298	462

OVERALL POINT ESTIMATE 569,464
OVERALL STANDARD ERROR 102,337

CONFIDENCE LIMITS
80% CONFIDENCE LEVEL
LOWER LIMIT 438,314
UPPER LIMIT 700,615
PRECISION AMOUNT 131,151
PRECISION PERCENT 23.03%
Z-VALUE USED 1.281551565545

	90% CONFIDENCE LEVEL
LOWER LIMIT	401,134
UPPER LIMIT	737,795
PRECISION AMOUNT	168,330
PRECISION PERCENT	29.56%
Z-VALUE USED	1.644853626951

	95% CONFIDENCE LEVEL
LOWER LIMIT	368,887
UPPER LIMIT	770,042
PRECISION AMOUNT	200,578
PRECISION PERCENT	35.22%
Z-VALUE USED	1.959963984540

Some highlighted values: (1) 405 is $90 + 110 + 85 + 120$, (2) 2,298 is the total number of third-stage units (universe) for the four sampled primary units, (3) 462 is $101 + 123 + 93 + 145$.

The point estimate and confidence intervals: The point estimate (highlighted) for the universe total difference amount is 569,464 (that is, \$569,464,000). The 95% confidence interval for this amount is from 368,887 to 770,042 (\$368,887,000 to \$770,042,000). The PRECISION

AMOUNT at the 95% confidence level is $(Z\text{-value})(\text{standard error of } \hat{T}) =$

$(1.959963984540)(102,337) = 200,578$ (that is, \$200,578,000). This value is 35.22% of the point estimate.

Notice that the output also contains the estimated totals for each primary unit (region) and each secondary unit (university). For example, the estimated total difference for UNIV1 within region 5 is 225 (\$225,000). The sample average of the 10 university estimates in region 5 is 392 (\$392,000). Since there are 90 universities in this region, the point estimate for the region 5 total is $(90)(392) = 35,280$ (more precisely, 35,256 or \$35,256,000).

The average of the four regional estimates is $(35,256 + 55,643 + 26,388 + 72,534)/4 = 47,455.25$ (\$47,455,250). Since there are 12 regions in the universe, the (unbiased) point estimate for the

universe total is $(12)(47,455,250) =$ approximately \$569,463,000. The actual amount (highlighted) is \$569,464,000.

FORMULAS

1. Point estimate of the universe total (T):

$$\hat{T} = \frac{N}{n} \sum_{i=1}^n \hat{T}_i$$

where $\hat{T}_i = \frac{M_i}{m_i} \sum_{j=1}^{m_i} \hat{T}_{ij}$ is the estimate of the total for the i-th sampled P.U. and where

$$\hat{T}_{ij} = \frac{B_{ij}}{b_{ij}} \sum_{k=1}^{b_{ij}} y_{ijk}$$

is the estimate of the total for the j-th S.U. within the i-th sampled P.U.

Notation: $n =$ number of primary units (P.U.s) in the sample

$N =$ number of P.U.s in the universe

$m_i =$ number of sampled secondary units (S.U.s) in the i-th P.U. ($i = 1, \dots, n$)

$M_i =$ number of S.U.s in the universe in the i-th P.U. ($i = 1, \dots, n$)

$b_{ij} =$ number of 3rd-stage items in the sample for the i-th P.U. and j-th S.U.
($i = 1, \dots, n$ and $j = 1, \dots, m_i$)

$B_{ij} =$ number of 3rd-stage items in the universe for the i-th P.U. and j-th S.U.
($i = 1, \dots, n$ and $j = 1, \dots, m_i$)

$y_{ijk} =$ sample value of the k-th item from the i-th P.U. and j-th S.U.
($i = 1, \dots, n$ and $j = 1, \dots, m_i$, $k = 1, \dots, b_{ij}$)

NOTE: The value of n , N , along with m_i , M_i , b_{ij} , B_{ij} for each **sampled** primary and secondary unit must be known.

2. Estimated variance of \hat{T} :

$$v(\hat{T}) = \frac{N(N-n)}{n} s^2 + \frac{N}{n} \sum_{i=1}^n \frac{M_i(M_i - m_i)}{m_i} s_i^2 + \frac{N}{n} \sum_{i=1}^n \frac{M_i}{m_i} \sum_{j=1}^{m_i} \frac{B_{ij}(B_{ij} - b_{ij})}{b_{ij}} s_{ij}^2$$

$$\text{where } s^2 = \left[\sum_{i=1}^n \hat{T}_i^2 - \left(\sum_{i=1}^n \hat{T}_i \right)^2 / n \right] / (n-1)$$

$$s_i^2 = \left[\sum_{j=1}^{m_i} \hat{T}_{ij}^2 - \left(\sum_{j=1}^{m_i} \hat{T}_{ij} \right)^2 / m_i \right] / (m_i - 1)$$

$$s_{ij}^2 = \left[\sum_{k=1}^{b_{ij}} y_{ijk}^2 - \left(\sum_{k=1}^{b_{ij}} y_{ijk} \right)^2 / b_{ij} \right] / (b_{ij} - 1)$$

3. The approximate 95% confidence interval for T:

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval, replace 1.959963984540 with 1.281551565545.

RHC Two-Stage Variable Sampling

For a discussion on the motivation behind the RHC (developed by Rao, Hartley, and Cochran) sampling procedure, refer to the **RHC Sample Selection** section, contained in the **Random Numbers** section of this manual. It provides a method of sample selection that allows sampling without replacement while maintaining the flavor of sampling using probability proportional to size. When the primary units (P.U.s) are selected, the size of each P.U. is considered rather than obtaining a simple random sample of P.U.s.

The size of each P.U. is rather arbitrary and can be the number of people, dollars, beds (for hospitals), and so forth. In general, you can expect improved precision using the RHC procedure if there is a high correlation between the size of each P.U. and the number of secondary units (S.U.s) within each P.U.. In other words, larger P.U.s should contain a larger number of S.U.s.

The P.U.s are selected using the **RHC Sample Selection** program. A random sample is then obtained for each selected P.U. and the variable(s) of interest (e.g., dollars in error) is/are recorded.

Example 5. (Note: This is the same example used in Example 8 in the **RHC Sample Selection** discussion contained in the **Random Numbers** section). In a particular region of the United States there are $N = 90$ universities (primary units) with Government research grants. Because these universities are so widespread, it was decided to use a cluster sample of $n = 10$ universities. Rather than audit all grants at a selected university, it was decided (based on available resources) to audit roughly 20% of the grants at each selected university. As a measure of the size for each university, the total grant dollars were used.

DATA: University ID, number of grants, total grant dollars (90 rows)

The data are contained in UNIVRHC.TXT.

OUTPUT: The 10 universities to use in the sample (see last page of computer output):

UNIV78, UNIV42, UNIV49, UNIV5, UNIV19,

UNIV38, UNIV62, UNIV28, UNIV60, UNIV75

Here there are 10 groups with 9 universities per group. The output file created by this program is OUTRHC.TXT. This program is required by the RHC appraisal program.

Data set UNIVRHC.TXT and the program output are contained in the pages to follow.

Dataset UNIVRHC.TXT

< - - - continued - - - >
 (1) (2)
 (3)

< - - - continued - - - >

UNIV1	42	8	UNIV31	52	11	UNIV61	66	13
UNIV2	21	4	UNIV32	66	14	UNIV62	77	18
UNIV3	63	13	UNIV33	25	5	UNIV63	31	7
UNIV4	74	16	UNIV34	60	12	UNIV64	46	9
UNIV5	51	11	UNIV35	19	4	UNIV65	32	7
UNIV6	43	9	UNIV36	24	5	UNIV66	68	14
UNIV7	57	11	UNIV37	44	9	UNIV67	41	9
UNIV8	49	10	UNIV38	76	17	UNIV68	28	6
UNIV9	63	13	UNIV39	41	9	UNIV69	66	14
UNIV10	18	4	UNIV40	77	18	UNIV70	31	7
UNIV11	64	13	UNIV41	37	8	UNIV71	27	6
UNIV12	56	11	UNIV42	63	12	UNIV72	33	7
UNIV13	19	4	UNIV43	52	11	UNIV73	23	4
UNIV14	44	9	UNIV44	76	17	UNIV74	71	15
UNIV15	20	4	UNIV45	51	10	UNIV75	75	16
UNIV16	34	7	UNIV46	23	4	UNIV76	47	10
UNIV17	25	6	UNIV47	24	5	UNIV77	50	10
UNIV18	38	9	UNIV48	68	15	UNIV78	37	7
UNIV19	72	16	UNIV49	34	7	UNIV79	77	18
UNIV20	46	10	UNIV50	49	10	UNIV80	49	10
UNIV21	44	9	UNIV51	55	11	UNIV81	76	17
UNIV22	64	13	UNIV52	38	9	UNIV82	66	14
UNIV23	45	9	UNIV53	72	16	UNIV83	28	6
UNIV24	55	11	UNIV54	51	10	UNIV84	77	17
UNIV25	29	7	UNIV55	71	15	UNIV85	27	6
UNIV26	36	7	UNIV56	59	12	UNIV86	75	17
UNIV27	40	9	UNIV57	23	4	UNIV87	71	15
UNIV28	78	18	UNIV58	57	11	UNIV88	59	12
UNIV29	49	10	UNIV59	53	11	UNIV89	71	15
UNIV30	60	12	UNIV60	64	13	UNIV90	72	16

Columns: (1) unit ID
 (2) number of grants
 (3) grant dollar amount (x \$100,000) ◀ This is the size of the university.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 3/15/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 12:52
 NAME OF INPUT FILE: C:\TEMP\UNIVRHC.TXT

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV51                          11                      55
UNIV44                          17                      76
UNIV32                          14                      66
UNIV78 <-- selected              7                      37
UNIV79                          18                      77
UNIV2                           4                      21
UNIV52                          9                      38
UNIV33                          5                      25
UNIV47                          5                      24

GROUP TOTALS:    9                      90                      419
  
```

```

***** GROUP 2 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV6                           9                      43
UNIV42 <-- selected              12                     63
UNIV65                          7                      32
UNIV40                          18                     77
UNIV45                          10                     51
UNIV1                           8                      42
UNIV80                          10                     49
UNIV36                          5                      24
UNIV70                          7                      31

GROUP TOTALS:    9                      86                      412
  
```

< Groups 3 Through 9 Are Omitted Here >

```

***** GROUP 10 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
UNIV22                          13                      64
UNIV39                          9                      41
UNIV88                          12                      59
UNIV55                          15                      71
UNIV29                          10                      49
UNIV75 <-- selected              16                      75
UNIV87                          15                      71
UNIV13                          4                      19
UNIV53                          16                      72

GROUP TOTALS:    9                      110                     521
  
```

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 3/15/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 12:52
 NAME OF OUTPUT FILE: C:\TEMP\OutRHCsummary.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF PRIMARY UNITS IN THE POPULATION: 90
 NUMBER OF PRIMARY UNITS SAMPLED: 10

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV78	37	7	90	9
UNIV42	63	12	86	9
UNIV49	34	7	96	9
UNIV5	51	11	84	9
UNIV19	72	16	89	9
UNIV38	76	17	89	9
UNIV62	77	18	92	9
UNIV28	78	18	115	9
UNIV60	64	13	99	9
UNIV75	75	16	110	9

Example--continued. The sample size for each selected university (P.U.) was chosen to be approximately 20% of the universe size. This leads to the following table where M_i is the total number of grants at the i -th university, and m_i is the number of audited grants at the i -th university.

University	M_i	m_i
UNIV78	37	7
UNIV42	63	13
UNIV49	34	7
UNIV5	51	10
UNIV19	72	14
UNIV38	76	15
UNIV62	77	15
UNIV28	78	16
UNIV60	64	13
UNIV75	75	<u>15</u>
		125

Data from these 125 secondary units (grants) were obtained by recording the total amount that was charged to each grant after the scheduled completion of this grant (dollars in error). The error amounts (in thousands of dollars) for each grant are contained in data set DATARHC2.TXT.

The output file created using the sample selection program is also used as input to the appraisal program in this two-step procedure. This file (PRIMRHC2.TXT), along with DATARHC2.TXT, are used as the input files for the **RHC Two-Stage** appraisal program. Both files are listed on the next page and the computer output from the appraisal program immediately follows. The illustrated data file (DATARHC2.TXT) contains the data for the first two universities (primary units) and the last two universities.

Dataset DATARHC2.TXT

1	9
2	2
3	9
4	6
5	0
6	5
7	7
8	2
9	7
10	6
11	0
12	6
13	0
14	3
15	4
16	1
17	13
18	8
19	0
20	6
21	11
22	8
23	8
24	0
.	
.	
.	
100	7
101	10
102	2
103	6
104	0
105	8
106	4
107	0
108	10
109	3
110	2
111	5
112	10
113	0
114	0
115	0
116	0

117 0
 118 8
 119 9
 120 0
 121 2
 122 8
 123 4
 124 6
 125 2

Output/Input file PRIMRHC2.TXT

UNIV78	37	7	7	90	9
UNIV42	63	13	12	86	9
UNIV49	34	7	7	96	9
UNIV5	51	10	11	84	9
UNIV19	72	14	16	89	9
UNIV38	76	15	17	89	9
UNIV62	77	15	18	92	9
UNIV28	78	16	18	115	9
UNIV60	64	13	13	99	9
UNIV75	75	15	15	110	9

NOTE: File PRIMRHC2.TXT was created by adding the third column containing sample sizes to the output file created by the **RHC Sample Selection** program.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 3/15/2010 RHC TWO-STAGE VARIABLE APPRAISAL Time: 13:26
 AUDIT/REVIEW: RHC 2-Stage

DATA FILE USED: C:\TEMP\DATARHC2.TXT

PRIMARY UNIT	SAMPLE SIZE	==DIFFERENCE== SAMPLE TOTAL	NUMBER OF NONZERO ITEMS
1	7	38.00	6
2	13	56.00	10
3	7	33.00	4
4	10	55.00	8
5	14	67.00	13
6	15	76.00	14
7	15	79.00	14
8	16	84.00	13
9	13	61.00	11
10	15	54.00	9
TOTALS	125	603.00	102

PRIMARY UNIT FILE USED: C:\TEMP\PRIMRHC2.txt

P.U. NBR	PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
1	UNIV78	37	7	90	9
2	UNIV42	63	12	86	9
3	UNIV49	34	7	96	9
4	UNIV5	51	11	84	9
5	UNIV19	72	16	89	9
6	UNIV38	76	17	89	9
7	UNIV62	77	18	92	9
8	UNIV28	78	18	115	9
9	UNIV60	64	13	99	9
10	UNIV75	75	15	110	9
TOTALS		627	134	950	90

--- POINT ESTIMATES ---

* **Note:** 2,582 is the product of 5.43, 37, and 12.857. This is the point estimate for P.U. 1.

P.U. NBR	SAMPLE SIZE	==DIFFERENCE== SAMPLE MEAN	SECONDARY UNIVERSE	SIZES RATIO	POINT ESTIMATE
1	7	5.43	37	12.857	2,582 *
2	13	4.31	63	7.167	1,945
3	7	4.71	34	13.714	2,198
4	10	5.50	51	7.636	2,142
5	14	4.79	72	5.563	1,917
6	15	5.07	76	5.235	2,016
7	15	5.27	77	5.111	2,073

8	16	5.25	78	6.389	2,616
9	13	4.69	64	7.615	2,287
10	15	3.60	75	7.333	1,980

TOTALS: 125 627 21,756

Note: 21,756 is the point estimate
of the universe total (\hat{T}).

--- VARIANCE COMPONENTS ---

P.U. NBR	WITHIN VARIANCE	BETWEEN VARIANCE	TOTAL VARIANCE
1	23,689	283,356	307,044
2	25,870	659	26,529
3	38,797	0	38,797
4	22,443	53,230	75,674
5	19,701	15,570	35,271
6	13,977	523	14,500
7	33,107	1,192	34,299
8	29,738	247	29,985
9	23,704	369	24,074
10	31,366	247,927	279,293
TOTALS:	262,392	603,072	865,464

Note: 865,464 is equal to $v(\hat{T})$.

PRIMARY UNITS SAMPLED: 10
PRIMARY UNITS NOT SAMPLED: 80
PRIMARY UNITS IN POPULATION: 90

POINT ESTIMATE OF POPULATION TOTAL: 21,756

STANDARD ERROR 930

CONFIDENCE LIMITS

80% CONFIDENCE LEVEL

LOWER LIMIT 20,564
UPPER LIMIT 22,948
PRECISION AMOUNT 1,192
PRECISION PERCENT 5.48%
Z-VALUE USED 1.281551565545

90% CONFIDENCE LEVEL

LOWER LIMIT 20,226
UPPER LIMIT 23,286
PRECISION AMOUNT 1,530
PRECISION PERCENT 7.03%
Z-VALUE USED 1.644853626951

95% CONFIDENCE LEVEL

LOWER LIMIT 19,933
UPPER LIMIT 23,579
PRECISION AMOUNT 1,823
PRECISION PERCENT 8.38%
Z-VALUE USED 1.959963984540

Discussion: The (highlighted) estimate of the universe total on the previous page obtained using formula 1 is:

$$\begin{aligned}\hat{T} &= (\text{Estimate of group 1 total}) + \cdots + (\text{Estimate of group 10 total}) \\ &= (90/7)(5.43)(37) + (86/12)(4.31)(63) + \cdots + (110/15)(3.60)(75) \\ &= 2,582 + 1,945 + \cdots + 1,980 \\ &= 21,756 \text{ } (\$21,756,000)\end{aligned}$$

Using formula 2, the estimated variance of \hat{T} is:

$$v(\hat{T}) = 865,464$$

and the estimated standard error of \hat{T} is the square root of 865,464; that is, 930 (highlighted on the previous page).

The approximate 95% confidence interval is:

$$\begin{aligned}21,756 \pm 1.959963984540(930) \\ 21,756 \pm 1,823\end{aligned}$$

that is, 19,933 to 23,579 (\$19,933,000 to \$23,579,000).

FORMULAS

Definitions

1. P.U. stands for “primary unit” and S.U. is “secondary unit”
2. A_i = size of i-th P.U.
3. S_i = (size of i-th P.U.)/(size of entire population) = A_i /(size of entire population)
4. B_i = total size for i-th group
5. π_i = (total size for i-th group)/(size of entire population) = B_i /(size of entire population)
6. N = number of P.U.s in the population
7. N_i = number of P.U.s in the i-th group
8. n = number of P.U.s in the sample
9. M_i = number of S.U.s in the i-th sampled P.U. (population)
10. m_i = number of S.U.s in the i-th sampled P.U. (sample)

Estimate of population total (T)

$$\hat{T} = \sum_{i=1}^n \left(\frac{B_i}{A_i} \right) M_i \bar{y}_i$$

where \bar{y}_i = average of m_i sampled S.U.s and B_i/A_i is labeled SIZES RATIO in the computer output.

Estimated variance of \hat{T}

$$v(\hat{T}) = V_1 + V_2 \quad \text{where}$$

$$V_1 = \frac{\sum_{i=1}^n N_i^2 - N}{N^2 - \sum_{i=1}^n N_i^2} \sum_{i=1}^n \pi_i \left(\frac{M_i \bar{y}_i}{S_i} - \hat{T} \right)^2$$

and

$$V_2 = \sum_{i=1}^n \pi_i \frac{M_i}{S_i} \left(\frac{M_i - m_i}{m_i} \right) s_i^2$$

where s_i^2 = variance of the m_i sampled S.U.s.

NOTE: The estimated standard error of \hat{T} is $\sqrt{v(\hat{T})}$

Approximate 95% confidence interval for the population total (T)

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

RHC Three-Stage Variable Sampling

The RHC sampling procedure can be used for a three-stage design.

The steps for such a procedure are the following:

1. A sample of primary units (clusters) is obtained as in the one- and two-stage procedures. The size of the primary units is considered for this sample, where pps sampling is used for each group of primary units.
2. A sample of secondary units is obtained within each chosen primary unit by partitioning the primary unit into random groups. The group sizes are chosen to be as nearly equal as possible. Using pps sampling and the size of each secondary unit, one secondary unit is chosen from each of the secondary groups.
3. A random sample of third-stage units is obtained for each of the chosen secondary units. No attention is paid to “size” here. This is a random sample.

Prior to running the appraisal program, the user must run the **RHC Sample Selection** program in the OAS software.

Example 6. The situation discussed in Example 4 in the **Three-Stage Unrestricted** section will be appraised using the RHC methodology. For this example, the stages are:

Stage 1: REGION (select 4 out of 12 regions)

Stage 2: UNIVERSITY (select 10 from each selected region)

Stage 3: GRANT (select approximately 20% of all grants at each university)

Selection of Primary Units

A file must be constructed containing (for each region) (1) the number of secondary units (universities) in this region and (2) the size of this region (total dollars of grants). This file is GRANTSPU.TXT. The selected regions are 4, 6, 8, and 10.

NOTE: Seed values of 100 and 200 were used to select the primary units. In practice, it is recommended that the user not set these seed values.

--- Data set GRANTSPU.TXT ---

	(A)	(B)
REGION1	117	1250
REGION2	63	610
REGION3	91	720
REGION4	123	1320
REGION5	107	1160
REGION6	116	1240
REGION7	102	960
REGION8	118	1300
REGION9	122	1320
REGION10	85	640
REGION11	94	930
REGION12	62	550

NOTE: It is okay to set the number of S.U.s [column (A)] equal to one in this file. The actual number of S.U.s must be known for the selected P.U.s. The correct number of S.U.s must then be inserted into file GRANTSPUOUT.TXT (the highlighted values).

Columns: (A) number of universities (S.U.s)
 (B) size of each P.U. (total grant amount x \$100,000)

--- Data set GRANTSPUOUT.TXT ---

REGION6	116	1240	3100	3
REGION4	123	1320	3410	3
REGION8	118	1300	3170	3
REGION10	85	640	2320	3

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF PRIMARY UNIT SAMPLE Time: 15:09
 NAME OF INPUT FILE: C:\TEMP\GRANTSPU.TXT

GROUPS OF PRIMARY UNITS

```

***** GROUP 1 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
REGION2                          610                      63
REGION6 <-- Selected             1,240                    116
REGION1                          1,250                    117

GROUP TOTALS: 3                  3,100                    296
  
```

```

***** GROUP 2 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
REGION4 <-- Selected             1,320                    123
REGION5                          1,160                    107
REGION11                         930                      94

GROUP TOTALS: 3                  3,410                    324
  
```

```

***** GROUP 3 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
REGION12                         550                      62
REGION8 <-- Selected             1,300                    118
REGION9                          1,320                    122

GROUP TOTALS: 3                  3,170                    302
  
```

```

***** GROUP 4 *****
PRIMARY UNIT IDENTIFICATION      PRIMARY UNIT      SECONDARY
=====                     SIZE                     UNIVERSE
=====                     =====
REGION3                          720                      91
REGION7                          960                    102
REGION10 <-- Selected            640                      85

GROUP TOTALS: 3                  2,320                    278
  
```

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

In practice, do not set these seed values.

NUMBER OF PRIMARY UNITS IN THE POPULATION: 12
 NUMBER OF PRIMARY UNITS SAMPLED: 4

< Program output - continued >

PRIMARY UNIT ID	SECONDARY UNIVERSE	PRIMARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
REGION6	116	1,240	3,100	3
REGION4	123	1,320	3,410	3
REGION8	118	1,300	3,170	3
REGION10	85	640	2,320	3

NOTE: The above four lines make up file GRANTSPUOUT.TXT

Selection of Secondary Units

The input for three-stage RHC can be greatly simplified if you only obtain information for each **selected** primary unit (that is, Regions 4, 6, 8, and 10 here). The information consists of the size of each secondary unit (university, here) and the number of third-stage units in the universe for each secondary unit. This input is shown in files REGION4.TXT, REGION6.TXT, REGION8.TXT, and REGION10.TXT. Each line in the files contains the number of third-stage units (grants) in the universe and the size of that secondary unit (total grant dollars x 100,000), in that order. After each of these four files is the computer output using the **RHC Sample Selection** program. A sample of 10 universities is selected for each region. The results are:

REGION	UNIVERSITIES
4	85, 46, 7, 82, 30, 34, 27, 66, 65, 80
6	113, 43, 78, 104, 89, 112, 30, 65, 3, 99
8	112, 6, 7, 93, 75, 111, 62, 115, 70, 99
10	78, 43, 7, 73, 55, 33, 10, 59, 64, 39

The previous five program runs (one at the primary level and four at the secondary level) created five output files. Using a word processor or spreadsheet, these files can be joined to form one of the input files (the one containing primary/secondary unit information) for the three-stage RHC program which calculates the confidence interval. The file for this example is PUSURHC3.TXT.

Data set REGION4.TXT

< --- continued --- >

< --- continued --- >

(1)	(2)	(3)
UNIV1	52	11
UNIV2	37	9
UNIV3	38	9
UNIV4	20	5
UNIV5	69	15
UNIV6	69	15
UNIV7	77	17
UNIV8	32	7
UNIV9	49	10
UNIV10	73	15
UNIV11	21	5
UNIV12	62	13
UNIV13	55	11
UNIV14	59	12
UNIV15	55	11
UNIV16	36	8
UNIV17	51	11
UNIV18	26	7
UNIV19	25	6
UNIV20	73	15
UNIV21	71	15
UNIV22	47	10
UNIV23	34	8
UNIV24	25	6
UNIV25	39	9
UNIV26	49	10
UNIV27	76	16
UNIV28	21	5
UNIV29	33	8
UNIV30	54	11
UNIV31	45	10
UNIV32	74	16
UNIV33	69	14
UNIV34	50	10
UNIV35	29	7
UNIV36	56	12
UNIV37	64	14
UNIV38	66	14
UNIV39	63	14
UNIV40	57	12
UNIV41	71	15
UNIV42	45	10
UNIV43	21	5
UNIV44	46	10
UNIV45	48	10
UNIV46	44	9
UNIV47	71	15
UNIV48	67	14
UNIV49	23	6
UNIV50	54	11

UNIV51	62	13
UNIV52	52	11
UNIV53	56	11
UNIV54	70	15
UNIV55	41	9
UNIV56	65	14
UNIV57	76	16
UNIV58	30	7
UNIV59	75	16
UNIV60	27	7
UNIV61	36	8
UNIV62	61	13
UNIV63	58	12
UNIV64	61	13
UNIV65	62	14
UNIV66	76	16
UNIV67	71	15
UNIV68	34	8
UNIV69	62	13
UNIV70	23	6
UNIV71	28	7
UNIV72	46	10
UNIV73	62	14
UNIV74	67	14
UNIV75	25	6
UNIV76	24	6
UNIV77	57	12
UNIV78	44	10
UNIV79	73	16
UNIV80	70	15
UNIV81	45	10
UNIV82	52	11
UNIV83	34	8
UNIV84	59	12
UNIV85	54	11
UNIV86	31	7
UNIV87	69	14
UNIV88	22	6
UNIV89	47	10
UNIV90	57	12
UNIV91	31	7
UNIV92	73	15
UNIV93	52	11
UNIV94	22	6
UNIV95	22	6
UNIV96	29	7
UNIV97	56	12
UNIV98	74	16
UNIV99	43	9
UNIV100	57	12

UNIV101	34	8
UNIV102	28	7
UNIV103	73	15
UNIV104	65	14
UNIV105	68	14
UNIV106	28	7
UNIV107	55	11
UNIV108	37	9
UNIV109	54	11
UNIV110	47	10
UNIV111	44	9
UNIV112	24	6
UNIV113	50	10
UNIV114	52	11
UNIV115	66	14
UNIV116	50	10
UNIV117	66	14
UNIV118	34	8
UNIV119	73	16
UNIV120	37	8
UNIV121	42	9
UNIV122	59	12
UNIV123	45	11

**NOTE: This file has
123 lines.**

Columns: (1) unit ID
(2) number of grants

(3) size of university (grant amount x \$100,000)

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 14:21

NAME OF INPUT FILE: C:\TEMP\REGION4.TXT

GROUPS OF SECONDARY UNITS

***** GROUP 1 *****

SECONDARY UNIT IDENTIFICATION	SECONDARY UNIT SIZE	3RD STAGE UNIVERSE
UNIV57	16	76
UNIV48	14	67
UNIV35	7	29
UNIV107	11	55
UNIV85 <-- Selected	11	54
UNIV103	15	73
UNIV86	7	31
UNIV2	9	37
UNIV81	10	45
UNIV58	7	30
UNIV36	12	56
UNIV49	6	23
GROUP TOTALS: 12	125	576

***** GROUP 2 *****

SECONDARY UNIT IDENTIFICATION	SECONDARY UNIT SIZE	3RD STAGE UNIVERSE
UNIV52	11	52
UNIV6	15	69
UNIV46 <-- Selected	9	44
UNIV69	13	62
UNIV108	9	37
UNIV44	10	46
UNIV50	11	54
UNIV121	9	42
UNIV1	11	52
UNIV43	5	21
UNIV87	14	69
UNIV39	14	63
GROUP TOTALS: 12	131	611

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

***** GROUP 10 *****

SECONDARY UNIT IDENTIFICATION	SECONDARY UNIT SIZE	3RD STAGE UNIVERSE
UNIV53	11	56
UNIV24	6	25
UNIV42	10	45
UNIV120	8	37
UNIV105	14	68
UNIV97	12	56

UNIV119	16	73
UNIV32	16	74
UNIV80 <-- Selected	15	70
UNIV96	7	29
UNIV13	11	55
UNIV62	13	61
UNIV59	16	75

GROUP TOTALS: 13	155	724
------------------	-----	-----

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 14:21

NAME OF OUTPUT FILE: C:\TEMP\OutRegion4.txt

FIRST SEED NUMBER:	100.00	SECOND SEED NUMBER:	200.00
--------------------	--------	---------------------	--------

NUMBER OF SECONDARY UNITS IN THE POPULATION: 123

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
UNIV85	54	11	125	12
UNIV46	44	9	131	12
UNIV7	77	17	119	12
UNIV82	52	11	129	12
UNIV30	54	11	141	12
UNIV34	50	10	140	12
UNIV27	76	16	138	12
UNIV66	76	16	128	13
UNIV65	62	14	125	13
UNIV80	70	15	155	13

<-- continued -->

UNIV1	56	10
UNIV2	27	5
UNIV3	56	11
UNIV4	23	5
UNIV5	72	13
UNIV6	24	5
UNIV7	61	11
UNIV8	65	12
UNIV9	68	13
UNIV10	40	8
UNIV11	64	12
UNIV12	66	13
UNIV13	80	14
UNIV14	53	9
UNIV15	36	7
UNIV16	53	10
UNIV17	47	9
UNIV18	73	14
UNIV19	41	8
UNIV20	58	11
UNIV21	45	9
UNIV22	43	8
UNIV23	56	10
UNIV24	35	7
UNIV25	34	7
UNIV26	65	13
UNIV27	78	14
UNIV28	35	7
UNIV29	31	6
UNIV30	58	11
UNIV31	29	6
UNIV32	76	14
UNIV33	57	10
UNIV34	42	8
UNIV35	69	13
UNIV36	58	11
UNIV37	31	6
UNIV38	33	6
UNIV39	40	8
UNIV40	51	9
UNIV41	60	11
UNIV42	78	14
UNIV43	39	7
UNIV44	46	9
UNIV45	58	11
UNIV46	59	11
UNIV47	53	10
UNIV48	57	10
UNIV49	28	6
UNIV50	63	12
UNIV51	31	6
UNIV52	60	11
UNIV53	30	6
UNIV54	30	6
UNIV55	40	8
UNIV56	26	5
UNIV57	24	5
UNIV58	44	8
UNIV59	67	13
UNIV60	56	10
UNIV61	33	7
UNIV62	40	8
UNIV63	68	13
UNIV64	70	13
UNIV65	57	10
UNIV66	40	7
UNIV67	54	10
UNIV68	65	12
UNIV69	62	12
UNIV70	28	5
UNIV71	56	10
UNIV72	41	8
UNIV73	31	6
UNIV74	31	6
UNIV75	46	9
UNIV76	38	7
UNIV77	62	12
UNIV78	63	12
UNIV79	50	9
UNIV80	53	9
UNIV81	39	7
UNIV82	39	7
UNIV83	39	7
UNIV84	25	5
UNIV85	67	13
UNIV86	47	9
UNIV87	54	10
UNIV88	50	9
UNIV89	35	7
UNIV90	66	13
UNIV91	65	12
UNIV92	71	13
UNIV93	29	6
UNIV94	74	14
UNIV95	66	13
UNIV96	71	13
UNIV97	43	8
UNIV98	62	11
UNIV99	80	14
UNIV100	57	11
UNIV101	22	5
UNIV102	33	6
UNIV103	78	5
UNIV104	25	9
UNIV105	76	8
UNIV106	39	5
UNIV107	48	5
UNIV108	54	7
UNIV109	63	12
UNIV110	28	8
UNIV111	69	8
UNIV112	27	10
UNIV113	33	8
UNIV114	52	7
UNIV115	33	15
UNIV116	23	10

Data set REGION6.TXT

**NOTE: This file has
116 lines.**

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 13:57
 NAME OF INPUT FILE: C:\TEMP\REGION6.txt

GROUPS OF SECONDARY UNITS

```

***** GROUP 1 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
UNIV52                             11                  60
UNIV45                             11                  58
UNIV32                             14                  76
UNIV86                              9                  47
UNIV113 <-- Selected                8                  33
UNIV85                             13                  67
UNIV109                            12                  63
UNIV87                             10                  54
UNIV2                               5                   27
UNIV80                              9                   53
UNIV53                              6                   30

GROUP TOTALS: 11                      108                 568
  
```

```

***** GROUP 2 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
UNIV33                             10                  57
UNIV48                             10                  57
UNIV6                               5                   24
UNIV43 <-- Selected                7                   39
UNIV68                             12                  65
UNIV41                             11                  60
UNIV46                             11                  59
UNIV1                              10                  56
UNIV40                              9                   51
UNIV88                              9                   50
UNIV36                             11                  58

GROUP TOTALS: 11                      105                 576
  
```

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

```

***** GROUP 10 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
UNIV20                             11                  58
UNIV22                              8                   43
UNIV39                              8                   40
UNIV111                             8                   69
UNIV100                             11                  57
UNIV29                              6                   31
UNIV105                             8                   76
UNIV79                              9                   50
UNIV99 <-- Selected                14                  80
  
```

UNIV13	14	80
UNIV60	10	56
UNIV54	6	30
GROUP TOTALS: 12	113	670

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 13:57

NAME OF OUTPUT FILE: C:\TEMP\OutRegion6.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 116

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
UNIV113	33	8	108	11
UNIV43	39	7	105	11
UNIV78	63	12	104	11
UNIV104	25	9	96	11
UNIV89	35	7	124	12
UNIV112	27	10	108	12
UNIV30	58	11	95	12
UNIV65	57	10	109	12
UNIV3	56	11	115	12
UNIV99	80	14	113	12

Data set REGION8.TXT

< - - - continued - - - >

< - - - continued - - - >

UNIV1 72 15
 UNIV2 44 10
 UNIV3 43 10
 UNIV4 55 12
 UNIV5 27 7
 UNIV6 34 8
 UNIV7 51 11
 UNIV8 42 10
 UNIV9 54 12
 UNIV10 25 6
 UNIV11 82 17
 UNIV12 65 14
 UNIV13 33 8
 UNIV14 48 10
 UNIV15 32 8
 UNIV16 82 17
 UNIV17 35 8
 UNIV18 54 12
 UNIV19 34 8
 UNIV20 62 14
 UNIV21 26 6
 UNIV22 31 7
 UNIV23 58 13
 UNIV24 61 13
 UNIV25 61 14
 UNIV26 54 12
 UNIV27 53 11
 UNIV28 56 12
 UNIV29 57 12
 UNIV30 26 6
 UNIV31 25 5
 UNIV32 37 9
 UNIV33 79 16
 UNIV34 60 13
 UNIV35 57 12
 UNIV36 27 7
 UNIV37 31 7
 UNIV38 75 15
 UNIV39 26 6
 UNIV40 36 9
 UNIV41 36 9
 UNIV42 49 10
 UNIV43 83 17
 UNIV44 71 15
 UNIV45 31 7
 UNIV46 42 10
 UNIV47 62 14
 UNIV48 54 11
 UNIV49 31 7
 UNIV50 80 16

UNIV51 77 16
 UNIV52 36 9
 UNIV53 75 16
 UNIV54 68 15
 UNIV55 34 8
 UNIV56 55 12
 UNIV57 42 10
 UNIV58 36 9
 UNIV59 36 9
 UNIV60 66 15
 UNIV61 61 13
 UNIV62 64 14
 UNIV63 72 15
 UNIV64 65 14
 UNIV65 58 13
 UNIV66 49 11
 UNIV67 30 7
 UNIV68 75 16
 UNIV69 33 8
 UNIV70 65 14
 UNIV71 55 12
 UNIV72 38 9
 UNIV73 36 9
 UNIV74 60 13
 UNIV75 52 11
 UNIV76 65 14
 UNIV77 49 10
 UNIV78 27 7
 UNIV79 48 10
 UNIV80 36 9
 UNIV81 66 15
 UNIV82 62 14
 UNIV83 70 15
 UNIV84 68 15
 UNIV85 53 11
 UNIV86 38 9
 UNIV87 35 8
 UNIV88 36 9
 UNIV89 26 6
 UNIV90 26 6
 UNIV91 51 11
 UNIV92 25 5
 UNIV93 54 11
 UNIV94 56 12
 UNIV95 81 17
 UNIV96 73 15
 UNIV97 44 10
 UNIV98 50 11
 UNIV99 60 13
 UNIV100 31 7

UNIV101 24 5
 UNIV102 26 6
 UNIV103 40 10
 UNIV104 77 16
 UNIV105 27 6
 UNIV106 65 15
 UNIV107 61 13
 UNIV108 36 9
 UNIV109 26 6
 UNIV110 38 9
 UNIV111 84 17
 UNIV112 75 16
 UNIV113 26 6
 UNIV114 45 10
 UNIV115 59 13
 UNIV116 59 13
 UNIV117 57 12
 UNIV118 58 12

NOTE: This file has 118 lines.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 14:03
 NAME OF INPUT FILE: C:\TEMP\REGION8.TXT

GROUPS OF SECONDARY UNITS

***** GROUP 1 *****

SECONDARY UNIT IDENTIFICATION	SECONDARY UNIT SIZE	3RD STAGE UNIVERSE
UNIV54	15	68
UNIV46	10	42
UNIV33	16	79
UNIV86	9	38
UNIV112 <-- Selected	16	75
UNIV85	11	53
UNIV108	9	36
UNIV87	8	35
UNIV2	10	44
UNIV55	8	34
UNIV34	13	60
GROUP TOTALS: 11	125	564

***** GROUP 2 *****

SECONDARY UNIT IDENTIFICATION	SECONDARY UNIT SIZE	3RD STAGE UNIVERSE
UNIV47	14	62
UNIV50	16	80
UNIV6 <-- Selected	8	34
UNIV44	15	71
UNIV68	16	75
UNIV42	10	49
UNIV48	11	54
UNIV1	15	72
UNIV41	9	36
UNIV89	6	26
UNIV37	7	31
GROUP TOTALS: 11	127	590

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

***** GROUP 10 *****

SECONDARY UNIT IDENTIFICATION	SECONDARY UNIT SIZE	3RD STAGE UNIVERSE
UNIV21	6	26
UNIV23	13	58
UNIV40	9	36
UNIV110	9	38
UNIV100	7	31
UNIV30	6	26
UNIV104	16	77
UNIV81	15	66
UNIV99 <-- Selected	13	60
UNIV13	8	33

UNIV60	15	66
UNIV56	12	55
GROUP TOTALS: 12	129	572

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

GENERATION OF SECONDARY UNIT SAMPLE

Time: 14:03

NAME OF OUTPUT FILE: C:\TEMP\OutRegion8.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 118

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV112	75	16	125	11
UNIV6	34	8	127	11
UNIV7	51	11	120	12
UNIV93	54	11	136	12
UNIV75	52	11	126	12
UNIV111	84	17	134	12
UNIV62	64	14	123	12
UNIV115	59	13	137	12
UNIV70	65	14	143	12
UNIV99	60	13	129	12

Data set
REGION10.TXT

<--continued -->

UNIV1 34 6
 UNIV2 32 5
 UNIV3 69 10
 UNIV4 23 4
 UNIV5 60 9
 UNIV6 72 11
 UNIV7 56 9
 UNIV8 28 5
 UNIV9 38 6
 UNIV10 60 9
 UNIV11 58 9
 UNIV12 37 6
 UNIV13 70 10
 UNIV14 37 6
 UNIV15 81 12
 UNIV16 53 9
 UNIV17 63 10
 UNIV18 32 5
 UNIV19 33 5
 UNIV20 37 6
 UNIV21 77 11
 UNIV22 52 8
 UNIV23 63 10
 UNIV24 41 7
 UNIV25 45 8
 UNIV26 34 6
 UNIV27 61 10
 UNIV28 70 10
 UNIV29 34 5
 UNIV30 22 4
 UNIV31 66 10
 UNIV32 69 10
 UNIV33 65 10
 UNIV34 26 4
 UNIV35 43 7
 UNIV36 65 10
 UNIV37 80 12
 UNIV38 74 11
 UNIV39 38 6
 UNIV40 43 7
 UNIV41 47 8
 UNIV42 59 9
 UNIV43 42 7
 UNIV44 54 9
 UNIV45 73 11

UNIV46 78 12
 UNIV47 72 11
 UNIV48 30 5
 UNIV49 47 8
 UNIV50 52 8
 UNIV51 24 4
 UNIV52 26 4
 UNIV53 22 4
 UNIV54 57 9
 UNIV55 78 12
 UNIV56 62 10
 UNIV57 57 9
 UNIV58 68 10
 UNIV59 52 8
 UNIV60 54 9
 UNIV61 41 7
 UNIV62 61 10
 UNIV63 79 12
 UNIV64 50 8
 UNIV65 54 9
 UNIV66 53 9
 UNIV67 40 7
 UNIV68 44 7
 UNIV69 39 7
 UNIV70 72 11
 UNIV71 76 11
 UNIV72 34 5
 UNIV73 27 5
 UNIV74 40 7
 UNIV75 41 7
 UNIV76 25 4
 UNIV77 41 7
 UNIV78 39 7
 UNIV79 58 9
 UNIV80 71 11
 UNIV81 37 6
 UNIV82 30 5
 UNIV83 78 12
 UNIV84 59 9
 UNIV85 29 5

Note: This file has 85 lines.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 13:49
 NAME OF INPUT FILE: C:\TEMP\REGION10.TXT

GROUPS OF SECONDARY UNITS

```

***** GROUP 1 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV44                             9                  54
UNIV32                             10                 69
UNIV77                             7                  41
UNIV78 <-- Selected                 7                  39
UNIV2                              5                  32
UNIV50                             8                  52
UNIV34                             4                  26
UNIV46                             12                 78

GROUP TOTALS:   8                      62                 391

```

```

***** GROUP 2 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV6                              11                 72
UNIV43 <-- Selected                 7                  42
UNIV62                             10                 61
UNIV41                             8                  47
UNIV1                              6                  34
UNIV40                             7                  43
UNIV79                             9                  58
UNIV36                             10                 65

GROUP TOTALS:   8                      68                 422

```

< GROUPS 3 THROUGH 9 ARE OMITTED HERE >

```

***** GROUP 10 *****
SECONDARY UNIT IDENTIFICATION      SECONDARY UNIT      3RD STAGE
=====                          SIZE              UNIVERSE
=====                          =====
UNIV71                             11                 76
UNIV9                              6                  38
UNIV21                             11                 77
UNIV23                             10                 63
UNIV39 <-- Selected                 6                  38
UNIV29                             5                  34
UNIV72                             5                  34
UNIV13                             10                 70
UNIV51                             4                  24

GROUP TOTALS:   9                      68                 454

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/22/2010 GENERATION OF SECONDARY UNIT SAMPLE Time: 13:49
 NAME OF OUTPUT FILE: C:\TEMP\OutRegion10.txt

FIRST SEED NUMBER: 100.00 SECOND SEED NUMBER: 200.00

NUMBER OF SECONDARY UNITS IN THE POPULATION: 85

NUMBER OF SECONDARY UNITS SAMPLED: 10

SECONDARY UNIT ID	3RD STAGE UNIVERSE	SECONDARY UNIT SIZE	GROUP SIZE	UNITS IN GROUP
=====	=====	=====	=====	=====
UNIV78	39	7	62	8
UNIV43	42	7	68	8
UNIV7	56	9	54	8
UNIV73	27	5	63	8
UNIV55	78	12	70	8
UNIV33	65	10	77	9
UNIV10	60	9	76	9
UNIV59	52	8	71	9
UNIV64	50	8	73	9
UNIV39	38	6	68	9

Constructing the data file

The data file for this example (PUSURHC3.TXT) is shown on the next page. This file was constructed using the **RHC Sample Selection** program to select the primary units (regions) and within each selected primary unit, the 10 secondary units (universities). The four lines beginning with REGIONx are from the output file created during the primary unit selection (GRANTSPUOUT.TXT). The 10 lines after each REGIONx line consist of the output file created when selecting the universities from each region (OUTREGION4.TXT, . . . , OUTREGION10.TXT). Using a word processor or spreadsheet, a column containing the sample sizes (highlighted) must be added to the files created by the five RHC Sample Selection programs.

--- Data set PUSURHC3.TXT ---

REGION4	123	10	1320	3410	3
UNIV85	54	11	11	125	12
UNIV46	44	9	9	131	12
UNIV7	77	15	17	119	12
UNIV82	52	10	11	129	12
UNIV30	54	11	11	141	12
UNIV34	50	10	10	140	12
UNIV27	76	15	16	138	12
UNIV66	76	15	16	128	13
UNIV65	62	12	14	125	13
UNIV80	70	14	15	155	13
REGION6	116	10	1240	3100	3
UNIV113	33	7	8	108	11
UNIV43	39	8	7	105	11
UNIV78	63	13	12	104	11
UNIV104	25	5	9	96	11
UNIV89	35	7	7	124	12
UNIV112	27	5	10	108	12
UNIV30	58	12	11	95	12
UNIV65	57	11	10	109	12
UNIV3	56	11	11	115	12
UNIV99	80	16	14	113	12
REGION8	118	10	1300	3170	3
UNIV112	75	15	16	125	11
UNIV6	34	7	8	127	11
UNIV7	51	10	11	120	12
UNIV93	54	11	11	136	12
UNIV75	52	10	11	126	12
UNIV111	84	17	17	134	12
UNIV62	64	13	14	123	12
UNIV115	59	12	13	137	12
UNIV70	65	13	14	143	12
UNIV99	60	12	13	129	12
REGION10	85	10	640	2320	3
UNIV78	39	8	7	62	8
UNIV43	42	8	7	68	8
UNIV7	56	11	9	54	8
UNIV73	27	5	5	63	8
UNIV55	78	16	12	70	8
UNIV33	65	13	10	77	9
UNIV10	60	12	9	76	9
UNIV59	52	10	8	71	9
UNIV64	50	10	8	73	9
UNIV39	38	8	6	68	9

Selection of Third-Stage Units

Since approximately 20% of the grants at each selected university are to be audited, the following sample sizes are determined:

Region 4:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV85	54	11
	UNIV46	44	9
	UNIV7	77	15
	UNIV82	52	10
	UNIV30	54	11
	UNIV34	50	10
	UNIV27	76	15
	UNIV66	76	15
	UNIV65	62	12
	UNIV80	70	<u>14</u>
			122

Region 6:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV113	33	7
	UNIV43	39	8
	UNIV78	63	13
	UNIV104	25	5
	UNIV89	35	7
	UNIV112	27	5
	UNIV30	58	12
	UNIV65	57	11
	UNIV3	56	11
	UNIV99	80	<u>16</u>
			95

Region 8:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV112	75	15
	UNIV6	34	7
	UNIV7	51	10
	UNIV93	54	11
	UNIV75	52	10
	UNIV111	84	17
	UNIV62	64	13
	UNIV115	59	12
	UNIV70	65	13
	UNIV99	60	<u>12</u>
			120

Region 10:	<u>University</u>	<u>Grants in universe</u>	<u>Number to be audited</u>
	UNIV78	39	8
	UNIV43	42	8
	UNIV7	56	11
	UNIV73	27	5
	UNIV55	78	16
	UNIV33	65	13
	UNIV10	60	12
	UNIV59	52	10
	UNIV64	50	10
	UNIV39	38	<u>8</u>
			101

The data file containing the errors for these 438 audited grants is DATARHC3.TXT, shown on the next page. The values for the first two universities in Region 4 and the last two universities in Region 10 are illustrated. Each sample value is equal to the total charges after the scheduled completion of the grant (in thousands of dollars). Notice that each line begins with a counter.

Finally, the **RHC Three-Stage** program is run, which generates a confidence interval for the universe total using input files PUSURHC3.TXT and DATARHC3.TXT. The output from this program is shown at the end of this section.

--- Data set DATARHC3.TXT ---

1	8
2	0
3	6
4	6
5	0
6	13
7	1
8	7
9	2
10	13
11	13
12	4
13	6
14	0
15	15
16	12
17	9
18	0
19	13
20	10

These are the sample values for UNIV85 and UNIV46 in Region 4.

.	
.	
.	
421	0
422	6
423	19
424	17
425	13
426	12
427	13
428	12
429	11
430	14
431	13
432	0
433	1
434	5
435	16
436	0
437	0
438	8

These are the sample values for UNIV64 and UNIV39 in Region 10.

Summary of results. Referring to the last page in the computer output, the estimate of the universe total (all 12 regions) is the OVERALL POINT ESTIMATE of 463,526 (\$463,526,000) with a corresponding estimated OVERALL STANDARD ERROR of 53,521 (\$53,521,000).

NOTE: This estimate does not require knowing the number of grants in the universe.

The program also provides estimates for the total error amount for each sampled P.U. (region) and for each of the groups of S.U.s (universities) within each sampled region. For example, the estimated error amount for Region 4 is 50,529 (\$50,529,000) and the estimated error amount for the group of 12 universities containing UNIV85 is 3,849 (\$3,849,000). The SIZES RATIO refers to the ratio of the size of the group containing this university to the size of this university. To illustrate, UNIV85 in Region 4 has a size of 11 and is in a group of size 125 (look at file REGION4.TXT). The SIZES RATIO here is $125/11 = 11.3636$.

The 80% confidence interval for the total error amount is 394,936 to 532,116 (\$394,936,000 to \$532,116,000). The PRECISION AMOUNT is the amount added and subtracted to the point estimate (463,526) to obtain the corresponding confidence interval. In the 80% confidence interval, the lower limit of 394,936 is obtained by subtracting the precision amount of 68,590 from 463,526. The PRECISION PERCENT is the precision amount divided by the point estimate, expressed as a percentage.

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

RHC THREE-STAGE VARIABLE APPRAISAL

Time: 16:27

AUDIT/REVIEW: RHC 3-Stage

DATA FILE USED: C:\TEMP\DATARHC3.TXT

PRIMARY/SECONDARY UNIVERSE FILE USED: C:\TEMP\PUSURHC3.txt

**** SAMPLED UNITS ****	THIRD STAGE	*****D I F F E R E N C E*****		
PRIMARY / SECONDARY IDENTIFICATION	UNIVERSE	SAMPLE SIZE	SAMPLE VALUE	NONZERO COUNT
=====	=====	=====	=====	=====
REGION4				
UNIV85	54	11	69.00	9
UNIV46	44	9	69.00	7
UNIV7	77	15	145.00	14
UNIV82	52	10	91.00	10
UNIV30	54	11	83.00	9
UNIV34	50	10	69.00	8
UNIV27	76	15	72.00	12
UNIV66	76	15	134.00	13
UNIV65	62	12	97.00	10
UNIV80	70	14	153.00	11
Total	615	122		103
REGION6				
UNIV113	33	7	81.00	7
UNIV43	39	8	36.00	8
UNIV78	63	13	83.00	9
UNIV104	25	5	56.00	5
UNIV89	35	7	82.00	6
UNIV112	27	5	29.00	2
UNIV30	58	12	65.00	9
UNIV65	57	11	82.00	8
UNIV3	56	11	67.00	6
UNIV99	80	16	164.00	15
Total	473	95		75
REGION8				
UNIV112	75	15	78.00	12
UNIV6	34	7	60.00	7
UNIV7	51	10	48.00	9
UNIV93	54	11	85.00	9
UNIV75	52	10	63.00	7
UNIV111	84	17	154.00	16
UNIV62	64	13	83.00	13
UNIV115	59	12	61.00	8
UNIV70	65	13	87.00	9
UNIV99	60	12	90.00	9
Total	598	120		99
REGION10				
UNIV78	39	8	75.00	7
UNIV43	42	8	71.00	6
UNIV7	56	11	123.00	11
UNIV73	27	5	32.00	3
UNIV55	78	16	123.00	11
UNIV33	65	13	113.00	11
UNIV10	60	12	104.00	8
UNIV59	52	10	77.00	7
UNIV64	50	10	117.00	9
UNIV39	38	8	43.00	5
Total	507	101		78

**** SAMPLED UNITS ****		THIRD STAGE	*****D I F F E R E N C E*****		
PRIMARY /	SECONDARY IDENTIFICATION	UNIVERSE	SAMPLE SIZE	SAMPLE VALUE	NONZERO COUNT
=====		=====	=====	=====	=====
TOTALS		2,193	438	3,414.00	355

--- POINT ESTIMATES---

**** SAMPLED UNITS ****		*****D I F F E R E N C E*****	
PRIMARY /	SECONDARY IDENTIFICATION	SAMPLE MEAN	POINT ESTIMATE
=====		=====	=====
REGION4			
UNIV85	Note: 3,849 is the estimate	6.27	11.3636
UNIV46	for the group containing	7.67	14.5556
UNIV7	UNIV85 (not just UNIV85).	9.67	7.0000
UNIV82	This group contained 12	9.10	11.7273
UNIV30	universities show earlier	7.55	12.8182
UNIV34	in the output using data	6.90	14.0000
UNIV27	set REGION4.TXT.	4.80	8.6250
UNIV66		8.93	8.0000
UNIV65		8.08	8.9286
UNIV80		10.93	10.3333

TOTAL	Estimate for Region 4 →	50,529
-------	-------------------------	--------

REGION6			
UNIV113		11.57	13.5000
UNIV43		4.50	15.0000
UNIV78		6.38	8.6667
UNIV104		11.20	10.6667
UNIV89		11.71	17.7143
UNIV112		5.80	10.8000
UNIV30		5.42	8.6364
UNIV65		7.45	10.9000
UNIV3		6.09	10.4545
UNIV99		10.25	8.0714

TOTAL	40,744
-------	--------

REGION8			
UNIV112		5.20	7.8125
UNIV6		8.57	15.8750
UNIV7		4.80	10.9091
UNIV93		7.73	12.3636
UNIV75		6.30	11.4545
UNIV111		9.06	7.8824
UNIV62		6.38	8.7857
UNIV115		5.08	10.5385
UNIV70		6.69	10.2143
UNIV99		7.50	9.9231

TOTAL	40,913
-------	--------

--- POINT ESTIMATES---

**** SAMPLED UNITS ****		*****D I F F E R E N C E*****	
PRIMARY /	SECONDARY IDENTIFICATION	SAMPLE MEAN	POINT ESTIMATE
=====		=====	=====
REGION10			
UNIV78		9.38	8.8571
UNIV43		8.88	9.7143
UNIV7		11.18	6.0000
UNIV73		6.40	12.6000
UNIV55		7.69	5.8333

UNIV33	8.69	7.7000	4,351
UNIV10	8.67	8.4444	4,391
UNIV59	7.70	8.8750	3,554
UNIV64	11.70	9.1250	5,338
UNIV39	5.38	11.3333	2,315
TOTAL			36,240

--- VARIANCE COMPONENTS FOR PRIMARY UNITS ---

**** SAMPLED UNITS **** PRIMARY UNIT IDENTIFICATION	WITHIN VARIANCE	BETWEEN VARIANCE	TOTAL VARIANCE
=====	=====	=====	=====
REGION4	757,373	10,383,517	11,140,890
REGION6	798,923	29,209,394	30,008,317
REGION8	710,631	8,065,650	8,776,280
REGION10	781,064	6,460,707	7,241,771

(Values of V_4) (Values of V_3)

--- OVERALL VARIANCE COMPONENTS ---

STAGE 1	STAGES 2 AND 3	TOTAL VARIANCE
=====	=====	=====
2,713,022,822	151,453,466	2,864,476,288
(Value of V_1)	(Value of V_2)	

*****D I F F E R E N C E*****

--- SUMMARY OF APPRAISAL RESULTS ---

PRIMARY UNITS SAMPLED	4
PRIMARY UNITS NOT SAMPLED	8
TOTAL PRIMARY UNITS	12
OVERALL POINT ESTIMATE	463,526
OVERALL STANDARD ERROR	53,521

CONFIDENCE LIMITS

80% CONFIDENCE LEVEL

LOWER LIMIT	394,936
UPPER LIMIT	532,116
PRECISION AMOUNT	68,590
PRECISION PERCENT	14.80%
Z-VALUE USED	1.281551565545

90% CONFIDENCE LEVEL

LOWER LIMIT	375,492
UPPER LIMIT	551,560
PRECISION AMOUNT	88,034
PRECISION PERCENT	18.99%
Z-VALUE USED	1.644853626951

95% CONFIDENCE LEVEL

LOWER LIMIT	358,627
UPPER LIMIT	568,425
PRECISION AMOUNT	104,899
PRECISION PERCENT	22.63%
Z-VALUE USED	1.959963984540

Discussion. In general, you can expect greater precision with the RHC procedure, provided there is a significant correlation between NUMBER OF UNITS and SIZE OF UNIT. To illustrate, consider the file containing the primary unit information used in the three-stage RHC illustration:

	(A)	(B)
REGION1	117	1250
REGION2	63	610
REGION3	91	720
REGION4	123	1320
REGION5	107	1160
REGION6	116	1240
REGION7	102	960
REGION8	118	1300
REGION9	122	1320
REGION10	85	640
REGION11	94	930
REGION12	62	550

Columns: (A) number of universities (S.U.s)
 (B) size of each P.U. (grant amount x \$100,000)

For this example, the correlation between columns (A) and (B) is .958, and we would expect a single- and two-stage RHC procedure to work quite well. For a three-stage procedure, this correlation rule must also apply within the sampled primary units, at the secondary unit level.

As mentioned earlier, the benefits of RHC sampling include the following:

- Precision is increased if the above correlation rule is satisfied.
- The flavor of pps sampling is maintained, since pps sampling is used to select a unit from each random group.
- Computations are relatively simple and straightforward.
- The point estimate (\hat{T}) is stable. This implies that when sampling indefinitely, the point estimate will exhibit relatively small variation.

- The point estimate of the variance of \hat{T} is stable, producing more reliable confidence intervals.

This implies that when sampling indefinitely, the lower confidence limits will exhibit relatively small variation.

FORMULAS

Definitions

1. $S_i = (\text{size of } i\text{-th P.U.})/(\text{size of entire population})$
2. $\pi_i = \Sigma S_i \text{ over the } i\text{-th group of P.U.s}$
3. $S_{ij} = (\text{size of } j\text{-th S.U. in the } i\text{-th sampled P.U.})/(\text{size of } i\text{-th sampled P.U.})$
(Note: denominator of S_{ij} = numerator of S_i)
4. $\pi_{ij} = \Sigma S_{ij} \text{ over the } j\text{-th group in } i\text{-th sampled P.U.}$
5. $N = \text{number of P.U.s (population)}$
6. $n = \text{number of P.U.s (sample)}$
7. $M_i = \text{number of S.U.s in } i\text{-th sampled P.U. (population)}$
8. $m_i = \text{number of S.U.s in } i\text{-th sampled P.U. (sample)}$
9. $K_{ij} = \text{number of third-stage units for } j\text{-th sampled S.U. in } i\text{-th sampled P.U. (population)}$
10. $k_{ij} = \text{number of third-stage units for } j\text{-th sampled S.U. in } i\text{-th sampled P.U. (sample)}$

Estimator of population total (T)

$$\hat{T} = \sum_{i=1}^n \pi_i \left(\frac{\hat{T}_i}{S_i} \right)$$

where \hat{T}_i = estimator of total for i -th sampled P.U.

$$= \sum_{j=1}^{m_i} \pi_{ij} \left(\frac{\hat{T}_{ij}}{S_{ij}} \right) \quad (\text{equation 1})$$

and \hat{T}_{ij} = estimator of population total for j -th sampled S.U. in i -th sampled P.U.

$$= K_{ij} \bar{y}_{ij} \text{ where } \bar{y}_{ij} = \text{average of } k_{ij} \text{ units at the third stage}$$

NOTE: It can be shown that \hat{T} is an unbiased estimator of T.

Estimated variance of \hat{T}

$$v(\hat{T}) = V_1 + V_2 \quad \text{where}$$

$$V_1 = \frac{\sum_{i=1}^n N_i^2 - N}{N^2 - \sum_{i=1}^n N_i^2} \sum_{i=1}^n \pi_i \left(\frac{\hat{T}_i}{S_i} - \hat{T} \right)^2 \quad (\text{equation 2})$$

and

$$V_2 = \sum_{i=1}^n \left(\frac{\pi_i}{S_i} \right) v(\hat{T}_i) \quad (\text{equation 3})$$

and N_i = number of P.U.s in the i -th group after the random split into n groups.

$v(\hat{T}_i)$ is obtained by applying the two-stage RHC procedure within the i -th sampled P.U.;

i.e., the i -th P.U. is viewed as the entire population. Consequently,

$$v(\hat{T}_i) = V_{3,i} + V_{4,i}$$

where

$$V_{3,i} = \frac{\sum_{j=1}^{m_i} M_{ij}^2 - M_i}{M_i^2 - \sum_{j=1}^{m_i} M_{ij}^2} \sum_{j=1}^{m_i} \pi_{ij} \left(\frac{K_{ij} \bar{y}_{ij}}{S_{ij}} - \hat{T}_i \right)^2$$

and

$$V_{4,i} = \sum_{j=1}^{m_i} \pi_{ij} \frac{K_{ij}}{S_{ij}} \left(\frac{K_{ij} - k_{ij}}{k_{ij}} \right) s_{ij}^2$$

and where

- (1) M_{ij} = the number of S.U.s in the j-th random group within the i-th sampled P.U.
- (2) \bar{y}_{ij} = average of the k_{ij} items for the j-th sampled S.U. within the i-th sampled P.U.

- Comments**
1. V_1 is essentially the same expression obtained for the single-stage RHC procedure and will be referred to as the “between unit” variation.
 2. V_2 is the contribution of the 2nd- and 3rd-stage variation and is obtained by treating each sampled P.U. as the population to be sampled using two (additional) stages.
 3. The estimated standard error of \hat{T} is $\sqrt{v(\hat{T})}$.

Approximate 95% confidence interval for the population total (T)

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

Stratified Cluster Variable Appraisal

With this procedure, you first stratify, then obtain a cluster (single-stage) sample within each stratum. This is motivated by the discussion in the RAT-STATS User's Guide.

The estimate of a universe total is the sum of the estimates for each stratum. The estimated variance of this estimator is the sum of the estimated variances for each stratum.

Example 7. In a large section of the U.S., an audit was conducted for 583 universities with health related research grants. It was decided to define two strata:

Stratum 1: state-supported universities ($N_1 = 415$)

Stratum 2: private universities ($N_2 = 168$)

Within each stratum, a **single-stage cluster** sample was obtained with $n_1 = 25$ universities selected from Stratum 1 and $n_2 = 10$ universities from Stratum 2. For each of the sampled universities, all health-related grants would be audited (since there weren't that many at each university) to determine the amount of charges improperly charged to these grants. The following data were obtained, where y_j is the total of the improper charges (in thousands of dollars) for the j -th university (cluster) and M_j is the number of grants (universe) for this university, all of which are audited.

NOTE: The number of grants audited at each university (the M_j values) are not used in the program calculations. They are supplied for informational purposes only. For example, if all the M_j values are set equal to 1, the resulting confidence intervals will be unchanged.

Stratum 1

Univ.	M _j	y _j	Univ.	M _j	y _j	
1	8	96	14	10	49	
2	12	121	15	9	53	
3	4	42	16	3	50	
4	5	65	17	6	32	
5	6	52	18	5	22	
6	6	40	19	5	45	$\Sigma M_j = 151$
7	7	75	20	4	37	
8	5	65	21	6	51	$\Sigma y_j = 1,329$
9	8	45	22	8	30	
10	3	50	23	7	39	
11	2	85	24	3	47	
12	6	43	25	8	41	
13	5	54				

Stratum 2

Univ.	M _j	y _j	Univ.	M _j	y _j	
1	2	18	6	8	96	
2	5	52	7	6	64	$\Sigma M_j = 49$
3	7	68	8	10	115	
4	4	36	9	3	41	$\Sigma y_j = 547$
5	3	45	10	1	12	

These values were stored in data file DATASTRCLUS.TXT. Immediately following the listing of this data file is the resulting computer output using the VARIABLE STRATIFIED CLUSTER program.

Data file DATASTRCLUS.TXT

```
STATE UNIVERSITIES  415  25
UNIV1      8      96
UNIV2     12     121
UNIV3      4      42
UNIV4      5      65
UNIV5      6      52
UNIV6      6      40
UNIV7      7      75
UNIV8      5      65
UNIV9      8      45
UNIV10     3      50
UNIV11     2      85
UNIV12     6      43
UNIV13     5      54
UNIV14    10      49
```

UNIV15	9	53		
UNIV16	3	50		
UNIV17	6	32		
UNIV18	5	22		
UNIV19	5	45		
UNIV20	4	37		
UNIV21	6	51		
UNIV22	8	30		
UNIV23	7	39		
UNIV24	3	47		
UNIV25	8	41		
PRIVATE UNIVERSITIES	168	10		
UNIV1	2	18		
UNIV2	5	52		
UNIV3	7	68		
UNIV4	4	36		
UNIV5	3	45		
UNIV6	8	96		
UNIV7	6	64		
UNIV8	10	115		
UNIV9	3	41		
UNIV10	1	12		

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 3/23/2010 STRATIFIED CLUSTER VARIABLE APPRAISAL Time: 14:36
 AUDIT/REVIEW: Variable - Stratified Cluster

DATA FILE USED: C:\Temp\DATASTRCLUS.TXT

STRATUM IDENTIFICATION CLUSTER IDENTIFICATION	SAMPLE UNIVERSE	SAMPLE SIZE	SAMPLED VALUE	POINT ESTIMATE
=====	=====	=====	=====	=====
STATE UNIVERSITIES	415	25		
UNIV1	8	8	96.00	
UNIV2	12	12	121.00	
UNIV3	4	4	42.00	
UNIV4	5	5	65.00	
UNIV5	6	6	52.00	
UNIV6	6	6	40.00	
UNIV7	7	7	75.00	
UNIV8	5	5	65.00	
UNIV9	8	8	45.00	
UNIV10	3	3	50.00	
UNIV11	2	2	85.00	
UNIV12	6	6	43.00	
UNIV13	5	5	54.00	
UNIV14	10	10	49.00	
UNIV15	9	9	53.00	
UNIV16	3	3	50.00	
UNIV17	6	6	32.00	
UNIV18	5	5	22.00	
UNIV19	5	5	45.00	
UNIV20	4	4	37.00	
UNIV21	6	6	51.00	
UNIV22	8	8	30.00	

UNIV23	7	7	39.00	
UNIV24	3	3	47.00	
UNIV25	8	8	41.00	
STRATUM TOTALS	151	151	1,329.00	22,061
PRIVATE UNIVERSITIES	168	10		
UNIV1	2	2	18.00	
UNIV2	5	5	52.00	
UNIV3	7	7	68.00	
UNIV4	4	4	36.00	
UNIV5	3	3	45.00	
UNIV6	8	8	96.00	
UNIV7	6	6	64.00	
UNIV8	10	10	115.00	
UNIV9	3	3	41.00	
UNIV10	1	1	12.00	
STRATUM TOTALS	49	49	547.00	9,190
STRATUM IDENTIFICATION	SAMPLE	SAMPLE		
CLUSTER IDENTIFICATION	UNIVERSE	SIZE	SAMPLED VALUE	POINT ESTIMATE
=====	=====	=====	=====	=====
STRATA TOTALS	583	35		
CLUSTER UNIT TOTALS	200	200	1,876.00	
OVERALL POINT ESTIMATE				31,251
OVERALL STANDARD ERROR				2,418
CONFIDENCE LEVEL	---80 PERCENT--	---90 PERCENT--	---95 PERCENT--	
LOWER LIMIT	28,152	27,273	26,511	
UPPER LIMIT	34,350	35,229	35,991	
PRECISION AMOUNT	3,099	3,978	4,740	
PRECISION PERCENT	9.92%	12.73%	15.17%	
Z-VALUE USED	1.281551565545	1.644853626951	1.959963984540	

Discussion. For stratum 1, the unbiased estimate of the universe total is

$$\hat{T}_1 = (415/25)(1,329) = 22,061 \text{ } (\$22,061,000)$$

The unbiased estimate of the universe total for stratum 2 is

$$\hat{T}_2 = (168/10)(547) = 9,190 \text{ } (\$9,190,000)$$

Consequently, an unbiased estimate of the universe total (highlighted) is

$$\hat{T} = \hat{T}_1 + \hat{T}_2 = 31,251 \text{ } (\$31,251,000)$$

Using formula 2, the estimated variance of \hat{T} is $v(\hat{T}) = v(\hat{T}_1) + v(\hat{T}_2) = 5,848,565$ and the corresponding standard error (highlighted) is 2,418.

The approximate 95% confidence interval for the universe total is

$$31,251 \pm 1.959963984540(2418)$$

that is, 26,511 to 35,991 (\$26,511,000 to \$35,991,000).

FORMULAS

1. Estimated total in the universe (T)

$$\hat{T} = \sum_{h=1}^L \frac{N_h}{n_h} \left(\sum_{j=1}^{n_h} y_{j,h} \right) = \sum_{h=1}^L N_h \bar{y}_h$$

where L = number of strata

N_h = number of clusters (universe) for stratum h

n_h = number of clusters (sample) for stratum h

$y_{j,h}$ = total of the variable of interest (e.g., errors) for the j-th P.U. within stratum h

\bar{y}_h = sample average for stratum h

NOTE: Let \hat{T}_h = estimated total for stratum h. Then $\hat{T}_h = N_h \bar{y}_h$ and $\hat{T} = \sum \hat{T}_h$

2. Estimated variance of \hat{T}

$$v(\hat{T}) = \sum_{h=1}^L \frac{N_h(N_h - n_h)}{n_h(n_h - 1)} \sum_{j=1}^{n_h} (y_{j,h} - \bar{y}_h)^2 = \sum_{h=1}^L v(\hat{T}_h)$$

3. Approximate 95% confidence interval for T

$$\hat{T} \pm 1.959963984540 \sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

Stratified Multistage Variable Appraisal

As with the stratified cluster procedure, you must first stratify the universe. Rather than take a cluster (single-stage) sample within each stratum, you will obtain a multistage (two-stage or three-stage) sample within each stratum. These multistage samples may be random (using the **Two-Stage Unrestricted** or **Three-Stage Unrestricted** programs) or may be obtained using the RHC procedure and the **RHC Two-Stage** or **RHC Three-Stage** programs.

Unlike the **Stratified Cluster** program, this program requires that you first run the appropriate multistage program on each stratum and record the results. The output results are then used as input to the **Stratified Multistage** program. You may store the results from each stratum (point estimate, standard error) in a file or simply input these values interactively.

NOTE: The “universe size” refers to the number of units at the most detailed level of the multistage sample. For example, if you are obtaining a three-stage sample within each stratum, then the “universe size” refers to the total number of third-stage units within this stratum.

Example 8. This example is similar to Example 7 in the Stratified Cluster section. In a particular region, the universe consisting of university grants is stratified by defining

Stratum 1: state-supported universities ($N_1 = 120$ univ.)

and Stratum 2: private universities ($N_2 = 85$ univ.)

Because these universities are so widespread, it was decided to employ a two-stage sample using 15 state supported universities and 10 private universities. Rather than audit all grants at a selected university, it was decided (based on available resources) to audit roughly 20% of the grants at each selected university to estimate the amount of charges improperly charged to these

grants. We know that there are a total of 5,800 grants within the universe of the 120 state supported universities and 4,500 grants within the 85 private universities.

The following data were obtained where y_{ij} is the dollars (in thousands) of improper charges for the j -th grant within the i -th sampled university, M_i is the total number of grants at the i -th university, and m_i is the number of audited grants at the i -th university. Also, \bar{y}_i and s_i^2 are the mean and variance of the sample values from the i -th university.

NOTE: The 15 state-supported universities and 10 private universities were obtained using the **Single-Stage Random Numbers** program. For ease of illustration, they will be referred to as University 1, 2, 3, . . . within each stratum.

The corresponding data files are the input files for the **Two-Stage Unrestricted** program. These are files STRMULT1.TXT and STRMULT2.TXT. The files containing the universe sizes are UNIV1.TXT and UNIV2.TXT.

State-supported universities

Univ.	M_i	m_i	Dollars ($y_{i,j}$, in thousands)												\bar{y}_i	s_i^2
1	60	12	4	0	0	6	7	11	0	5	4	0	8	2	3.92	13.36
2	50	10	4	7	0	0	6	10	3	3	2	0			3.50	11.17
3	45	9	3	1	0	5	5	8	10	0	4				4.00	12.00
4	40	8	2	7	10	0	6	5	4	0					4.25	12.21
5	55	11	7	1	6	0	0	0	5	12	8	2	4		4.09	15.49
6	58	12	0	0	6	5	12	8	2	0	7	7	2	5	4.50	14.27
7	62	12	4	1	0	3	10	7	6	6	0	5	4	4	4.17	8.70
8	52	10	3	8	0	6	2	10	0	0	5	2			3.60	12.49
9	50	10	7	1	4	0	2	6	0	0	0	5			2.50	7.61
10	45	9	2	1	0	0	0	5	8	6	10				3.56	14.53
11	40	8	4	7	3	8	0	1	6	10					4.88	12.13
12	48	10	0	0	6	5	2	3	8	0	12	0			3.60	16.93
13	57	11	3	0	1	10	5	0	6	6	7	3	8		4.45	11.07
14	60	12	4	1	0	0	0	6	8	10	4	3	2	5	3.58	10.63
15	54	11	3	0	1	1	4	8	6	9	5	0	3		3.64	9.65

Private universities

Univ.	M_i	m_i	Dollars ($y_{i,j}$, in thousands)												\bar{y}_i	s_i^2	
1	66	13	4	4	0	0	0	3	6	7	5	2	0	0	4	2.69	6.40
2	52	10	10	1	6	4	0	0	5	8	12	7				5.30	17.12
3	47	9	2	8	1	0	6	6	11	5	0					4.33	14.75
4	55	11	3	8	0	0	6	5	2	1	12	8	5			4.55	14.47
5	48	10	0	5	8	1	1	6	10	7	9	3				5.00	12.89
6	60	12	7	3	0	5	6	6	0	0	8	2	3	5		3.75	8.02
7	57	11	2	4	0	6	10	7	5	0	0	3	8			4.09	11.89
8	50	10	3	5	1	1	0	0	3	8	10	7				3.80	12.62
9	62	12	8	4	0	2	2	6	5	1	12	5	0	0		3.75	13.66
10	56	11	5	0	1	2	8	7	10	6	0	4	2			4.09	11.49

The data files are shown on the next page. The corresponding computer outputs immediately follow.

Data file STRMULT1.TXT

1	4
2	0
3	0
4	6
5	7
6	11
7	0
8	5
9	4
10	0
11	8
12	2
13	4
14	7
15	0
16	0
17	6
18	10
19	3
20	3
21	2
22	0
.	
.	
.	
133	4
134	1
135	0
136	0
137	0
138	6
139	8
140	10
141	4
142	3
143	2
144	5
145	3
146	0
147	1
148	1
149	4
150	8
151	6
152	9
153	5
154	0
155	3

Data file STRMULT2.TXT

1	4
2	4
3	0
4	0
5	0
6	3
7	6
8	7
9	5
10	2
11	0
12	0
13	4
14	10
15	1
16	6
17	4
18	0
19	0
20	5
21	8
22	12
23	7
.	
.	
.	
87	8
88	4
89	0
90	2
91	2
92	6
93	5
94	1
95	12
96	5
97	0
98	0
99	5
100	0
101	1
102	2
103	8
104	7
105	10
106	6
107	0
108	4
109	2

Data file UNIV1.TXT

1	60	12
2	50	10
3	45	9
4	40	8
5	55	11
6	58	12
7	62	12
8	52	10
9	50	10
10	45	9
11	40	8
12	48	10
13	57	11
14	60	12
15	54	11

Data file UNIV2.TXT

1	66	13
2	52	10
3	47	9
4	55	11
5	48	10
6	60	12
7	57	11
8	50	10
9	62	12
10	56	11

Output using the state-supported universities

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/20/2010 TWO-STAGE UNRESTRICTED VARIABLE APPRAISAL Time: 14:14
 AUDIT/REVIEW: Stratum 1

DATA FILE USED: C:\TEMP\STRMULT1.TXT

----- D I F F E R E N C E -----						
UNIT NBR	SAMPLE SIZE/ NONZERO ITEMS	SAMPLE MEAN	VARIANCE	UNIVERSE SIZE	POINT ESTIMATE	
1	12/8	3.92	13.36	60	235	
2	10/7	3.50	11.17	50	175	
3	9/7	4.00	12.00	45	180	
4	8/6	4.25	12.21	40	170	
5	11/8	4.09	15.49	55	225	
6	12/9	4.50	14.27	58	261	
7	12/10	4.17	8.70	62	258	
8	10/7	3.60	12.49	52	187	
9	10/6	2.50	7.61	50	125	
10	9/6	3.56	14.53	45	160	
11	8/7	4.88	12.12	40	195	
12	10/6	3.60	16.93	48	173	
13	11/9	4.45	11.07	57	254	
14	12/9	3.58	10.63	60	215	
15	11/9	3.64	9.65	54	196	
	155/114	4.15		776	3,010	
NOT SAMPLED						
	105			5,024		
OVERALL TOTALS						
	120			5,800	24,077	
STANDARD ERROR						
					1,277	

```

CONFIDENCE LIMITS
80% CONFIDENCE LEVEL
LOWER LIMIT                22,440
UPPER LIMIT                25,714
PRECISION AMOUNT           1,637
PRECISION PERCENT          6.80%
Z-VALUE USED               1.281551565545

90% CONFIDENCE LEVEL
LOWER LIMIT                21,976
UPPER LIMIT                26,178
PRECISION AMOUNT           2,101
PRECISION PERCENT          8.73%
Z-VALUE USED               1.644853626951

95% CONFIDENCE LEVEL
LOWER LIMIT                21,573
UPPER LIMIT                26,580
PRECISION AMOUNT           2,503
PRECISION PERCENT          10.40%
Z-VALUE USED               1.959963984540

```

Output using the private universities

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 4/20/2010    TWO-STAGE UNRESTRICTED VARIABLE APPRAISAL    Time: 14:13
AUDIT/REVIEW: Stratum 2

```

DATA FILE USED: C:\TEMP\STRMULT2.TXT

```

----- D I F F E R E N C E -----
UNIT   SAMPLE SIZE/
NBR     NONZERO ITEMS   SAMPLE MEAN   VARIANCE   UNIVERSE SIZE   POINT ESTIMATE

1       13/8           2.69         6.40       66              178
2       10/8           5.30        17.12      52              276
3        9/7           4.33        14.75      47              204
4       11/9           4.55        14.47      55              250
5       10/9           5.00        12.89      48              240
6       12/9           3.75         8.02      60              225
7       11/8           4.09        11.89      57              233
8       10/8           3.80        12.62      50              190
9       12/9           3.75        13.66      62              232
10      11/9           4.09        11.49      56              229

109/84           4.26                    553          2,257

NOT SAMPLED
75                                3,947
OVERALL TOTALS
85                                4,500          19,182

STANDARD ERROR                                873

```

CONFIDENCE LIMITS	
80% CONFIDENCE LEVEL	
LOWER LIMIT	18,064
UPPER LIMIT	20,300
PRECISION AMOUNT	1,118
PRECISION PERCENT	5.83%
Z-VALUE USED	1.281551565545
90% CONFIDENCE LEVEL	
LOWER LIMIT	17,747
UPPER LIMIT	20,617
PRECISION AMOUNT	1,435
PRECISION PERCENT	7.48%
Z-VALUE USED	1.644853626951
95% CONFIDENCE LEVEL	
LOWER LIMIT	17,472
UPPER LIMIT	20,892
PRECISION AMOUNT	1,710
PRECISION PERCENT	8.92%
Z-VALUE USED	1.959963984540

When running the STRATIFIED MULTISTAGE program, you should see the following input window containing values for the first stratum. The program output immediately follows.

Variable - Stratified Multistage

Name of Audit/Review Combining the Strata

Number of strata for this appraisal 2

Have you created a text data file for this appraisal? ☐ Yes ☒ No

Stratum 1

Point Estimate: 24077

Standard Error: 1277

OUTPUT TO

- ☒ Text File and Screen
- ☐ Printer and Screen
- ☐ Text File, Printer, and Screen
- ☐ Screen Only

Click here to save the data set

Next Stratum

Previous Stratum

CONTINUE

HELP EXIT Main Menu

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 4/20/2010 STRATIFIED MULTISTAGE VARIABLE APPRAISAL Time: 14:41
 AUDIT/REVIEW: Combining the Strata

THE ESTIMATORS ARE BASED ON THE FOLLOWING ENTRIES:

STRATUM	POINT ESTIMATE	STANDARD ERROR
1	24,077	1,277
2	19,182	873

= = = = = RESULTS = = = = =

POINT ESTIMATE	STANDARD ERROR
43,259	1,547

CONFIDENCE LEVEL	---80 PERCENT--	---90 PERCENT--	---95 PERCENT--
LOWER LIMIT	41,277	40,715	40,227
UPPER LIMIT	45,241	45,803	46,291
PRECISION AMOUNT	1,982	2,544	3,032
PRECISION PERCENT	4.58%	5.88%	7.01%
Z-VALUE USED	1.281551565545	1.644853626951	1.959963984540

Discussion. The point estimate for the universe total is the sum of the point estimates for the

two strata; that is, $\hat{T} = 24,077 + 19,182 = 43,259$ (\$43,259,000). The estimated variance of \hat{T}

is $v(\hat{T}) = (1277)^2 + (873)^2 = 2,392,858$ and the corresponding standard error is $\sqrt{2,392,858}$

= 1,547. The approximate 95% confidence interval for the universe total is

$$43,259 \pm 1.959963984540(1,547)$$

that is, 40,227 to 46,291 (\$40,227,000 to \$46,291,000).

FORMULAS

1. Estimated total in the universe (T)

$$\hat{T} = \sum_{h=1}^L \hat{T}_h$$

where \hat{T}_h is the point estimate for the universe total in stratum h and L is the number of strata.

2. Estimated variance of \hat{T}

$$v(\hat{T}) = \sum_{h=1}^L v(\hat{T}_h)$$

where $v(\hat{T}_h)$ is the estimated variance of \hat{T}_h and is equal to the square of the standard error of \hat{T}_h .

3. Approximate 95% confidence interval for T

$$\hat{T} \pm 1.959963984540\sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

Poststratification

Oftentimes sampling problems arise in which the user would like to stratify on a key variable but cannot place the sampling units into their correct strata until after the sample is selected.

Another situation arises when an auditor does not recognize a need to stratify prior to obtaining a simple random sample and the sample items are evaluated. The poststratification program is designed for such situations and provides reliable results if the overall sample size is large and the poststratified sample sizes are large (say, at least 20). It is however, less efficient than using a prestratified sample; that is, produces a slightly wider confidence interval for the same sample size.

A key thing to keep in mind here is that **the universe strata sizes must be known**.

Consequently, before you define a set of strata, make sure that you know the number of universe items in each of the strata. The program does not allow you to estimate these universe sizes. Poststratification is often appropriate when a simple random sample is not properly balanced according to major groupings of the population.

Example 9. In a recent hospital audit, the amount of unallowable bad debts was determined for a particular year. For the universe, it was known that there were $N_1 = 373$ inpatient bad debts (Stratum 1) and $N_2 = 1,146$ outpatient bad debts (Stratum 2). The total universe size is $N = N_1 + N_2 = 1,519$. So there are roughly 20% inpatient and 80% outpatient bad debts in the universe. Suppose that a simple random sample of 100 bad debts revealed:

Inpatient bad debts

$$n_1 = 45$$

$$\bar{y}_1 = \$240.00$$

$$s_1^2 = 22.04$$

Outpatient bad debts

$$n_2 = 55$$

$$\bar{y}_2 = \$30.00$$

$$s_2^2 = 198.56$$

The data files (POSTDATA.TXT and UNIVPOST.TXT) are shown on the next page and the resulting computer output immediately follows.

Data file POSTDATA.TXT

< - continued - >

Universe file UNIVPOST.TXT

1	242.27	51	25.99
2	240.43	52	31.75
3	243.43	53	42.37
4	241.00	54	28.82
5	235.71	55	14.80
6	239.74	56	22.44
7	249.53	57	12.93
8	232.61	58	37.22
9	243.51	59	12.29
10	243.70	60	21.02
11	241.36	61	24.76
12	238.86	62	49.69
13	241.20	63	50.43
14	246.14	64	9.03
15	236.91	65	32.47
16	244.91	66	28.43
17	231.91	67	18.54
18	249.58	68	31.48
19	227.84	69	4.45
20	239.07	70	54.76
21	232.89	71	9.45
22	242.15	72	19.73
23	238.93	73	33.71
24	243.41	74	59.84
25	241.61	75	22.73
26	240.04	76	15.54
27	229.43	77	22.14
28	238.12	78	21.81
29	240.82	79	53.10
30	241.84	80	38.72
31	236.08	81	36.58
32	248.11	82	39.26
33	239.39	83	53.71
34	236.00	84	40.95
35	238.67	85	24.82
36	242.88	86	45.05
37	241.90	87	25.30
38	238.96	88	58.31
39	234.03	89	25.30
40	237.86	90	9.48
41	239.01	91	15.93
42	239.21	92	25.00
43	245.45	93	29.27
44	242.08	94	26.29
45	241.39	95	38.26
46	46.32	96	31.30
47	37.59	97	37.97
48	31.23	98	38.52
49	22.92	99	11.53
50	44.96	100	3.64

1	373	45
2	1146	55

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 3/24/2010 POSTSTRATIFIED VARIABLE APPRAISAL Time: 9:51
 AUDIT/REVIEW: Poststratification

DATA FILE USED: C:\Temp\POSTDATA.TXT

```

-----D I F F E R E N C E-----
Stratum 1  SAMPLE SIZE / UNIVERSE SIZE          45          373
           MEAN                                240.00
           STANDARD DEVIATION                  4.69
           STANDARD ERROR (TOTAL)             347.13
           POINT ESTIMATE                      89,520

                                   CONFIDENCE LIMITS
                                   80% CONFIDENCE LEVEL
           LOWER LIMIT                        89,075
           UPPER LIMIT                        89,965
           PRECISION AMOUNT                   445
           PRECISION PERCENT                  .50%
           Z-VALUE USED                       1.281551565545

                                   90% CONFIDENCE LEVEL
           LOWER LIMIT                        88,949
           UPPER LIMIT                        90,091
           PRECISION AMOUNT                   571
           PRECISION PERCENT                  .64%
           Z-VALUE USED                       1.644853626951

                                   95% CONFIDENCE LEVEL
           LOWER LIMIT                        88,839
           UPPER LIMIT                        90,200
           PRECISION AMOUNT                   680
           PRECISION PERCENT                  .76%
           Z-VALUE USED                       1.959963984540

Stratum 2  SAMPLE SIZE / UNIVERSE SIZE          55          1,146
           MEAN                                30.00
           STANDARD DEVIATION                  14.09
           STANDARD ERROR (TOTAL)             1,800.03
           POINT ESTIMATE                      34,379

                                   CONFIDENCE LIMITS
                                   80% CONFIDENCE LEVEL
           LOWER LIMIT                        32,072
           UPPER LIMIT                        36,685
           PRECISION AMOUNT                   2,307
           PRECISION PERCENT                  6.71%
           Z-VALUE USED                       1.281551565545

                                   90% CONFIDENCE LEVEL
           LOWER LIMIT                        31,418
           UPPER LIMIT                        37,339
           PRECISION AMOUNT                   2,961
           PRECISION PERCENT                  8.61%
           Z-VALUE USED                       1.644853626951

```

		95% CONFIDENCE LEVEL	
		LOWER LIMIT	30,851
		UPPER LIMIT	37,907
		PRECISION AMOUNT	3,528
		PRECISION PERCENT	10.26%
		Z-VALUE USED	1.959963984540
OVERALL	SAMPLE SIZE / UNIVERSE SIZE	100	1,519
	POINT ESTIMATE	123,898	
	STANDARD ERROR	1,833	
		CONFIDENCE LIMITS	
		80% CONFIDENCE LEVEL	
		LOWER LIMIT	121,549
		UPPER LIMIT	126,248
		PRECISION AMOUNT	2,349
		PRECISION PERCENT	1.90%
		Z-VALUE USED	1.281551565545
		90% CONFIDENCE LEVEL	
		LOWER LIMIT	120,883
		UPPER LIMIT	126,914
		PRECISION AMOUNT	3,015
		PRECISION PERCENT	2.43%
		Z-VALUE USED	1.644853626951
		95% CONFIDENCE LEVEL	
		LOWER LIMIT	120,305
		UPPER LIMIT	127,491
		PRECISION AMOUNT	3,593
		PRECISION PERCENT	2.90%
		Z-VALUE USED	1.959963984540

Discussion. Using the usual estimator for a simple random sample, the estimate of the universe mean is

$$\bar{y} = [(45)(240) + (55)(30)]/100 = \$124.50$$

and the estimate of the universe total is

$$\hat{T} = (1519)(124.50) = \$189,115.50$$

Since there is an unusually high number of inpatient bad debts (and low outpatient), a better procedure would be to use the poststratified estimate of the universe total; namely

$$\hat{T}_{pst} = (373)(240.00) + (1146)(30.00) = \$123,900$$

The (more precise) computer-generated point estimate is \$123,898 (highlighted).

Also, the point estimate for the inpatient stratum is $\hat{T}_1 = (373)(240.00) = \$89,520$ (highlighted);

for the outpatient stratum, this estimate is $\hat{T}_2 = (1146)(30.00) = \$34,380$ (more precisely, the

highlighted value of \$34,379). Notice that $\hat{T}_{pst} = 89,520 + 34,379 = \$123,899$ (more precisely, \$123,898).

The estimated variance of \hat{T}_{pst} is $v(\hat{T}_{pst}) = v(\hat{T}_1) + v(\hat{T}_2)$

$$= \left[\frac{1419}{100} (373)(4.69484)^2 + \frac{1519}{100^2} (1146)(4.69484)^2 \right] + \left[\frac{1419}{100} (1146)(14.09098)^2 + \frac{1519}{100^2} (373)(14.09098)^2 \right]$$

$$= 120,499.88 + 3,240,111.44 = 3,360,611 \text{ and the estimated standard error is } \sqrt{3,360,611} =$$

\$1,833 (highlighted)

NOTE: The estimated standard error for stratum 1 is $\sqrt{120,499.88} = 347.13$ and the estimated standard error for stratum 2 is $\sqrt{3,240,111.44} = 1,800.03$.

The corresponding 95% confidence interval for the universe total is

$$123,898 \pm 1.959963984540(1,833)$$

that is, \$120,305 to \$127,491.

Comments

1. Poststratification allows you to obtain a single simple random sample (easier than obtaining a simple random sample from each stratum) and then stratify provided the strata sizes in the universe (N_i) are known. The value of N_i is multiplied by the sample mean \bar{y}_i to estimate

the total for the i-th stratum. These estimates are then summed over all the strata to estimate the universe total.

2. A minimum of 20 sampling units per stratum is required as well as 6 nonzero items per stratum (OA Policy and Procedures). The total sample size should be at least 100.
3. With poststratification, the sample sizes (n_i) are unknown in advance (random variables).
With stratified sampling, the sample sizes are fixed (nonrandom).

FORMULAS

1. Estimate of the universe total for the i-th stratum (T_i)

$$\hat{T}_i = N_i \bar{y}_i$$

where N_i = number of items (universe) in stratum i

\bar{y}_i = average of sample items in i-th stratum

L = number of strata

2. Estimate of universe total (T)

$$\hat{T}_{pst} = \sum_{i=1}^L N_i \bar{y}_i = \sum_{i=1}^L \hat{T}_i$$

3. Estimated variance of $\hat{T}_{pst} = v(\hat{T}_{pst}) = \sum_{i=1}^L v(\hat{T}_i)$ where

$$v(\hat{T}_i) = \left(\frac{N - n}{n} \right) N_i s_i^2 + \frac{N}{n^2} (N - N_i) s_i^2$$

where N = universe total = $\sum N_i$ and n is the total sample size.

4. Estimated standard error of $\hat{T}_i = \sqrt{v(\hat{T}_i)}$

5. Approximate 95% confidence interval for stratum total (T_i):

$$\hat{T}_i \pm 1.959963984540\sqrt{v(\hat{T}_i)}$$

6. Estimated standard error of \hat{T}_{pst} is $\sqrt{v(\hat{T}_{pst})}$

7. Approximate 95% confidence interval for universe total (T):

$$\hat{T}_{pst} \pm 1.959963984540\sqrt{v(\hat{T}_{pst})}$$

NOTE: For a 90% confidence interval in equations 5 and 7, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

Unknown Universe Size

This program calculates a confidence interval for a universe total when using variable sampling, as does the Unrestricted Variable Appraisal program. When using the Unrestricted Variable Appraisal program, one of the user queries is for the universe size (N) and this value must be known. For situations where N is unknown, the Unknown Universe Size program can be used.

Use of this program requires that two random samples be used -- one to estimate the universe size and the other to estimate one or more variable characteristics. Both samples must be appraised using the Unrestricted Variable Appraisal program prior to running this module since this program will ask for the mean and standard deviation of each sample.

The population of interest is a subset of some other universe. For example, the larger sampling frame might consist of 575 file drawers containing a mixture of dental claims and the population of interest consists of all claims related to a particular dental procedure (say, procedure ABC). Suppose we sample 70 of the file drawers and count the number of claims related to procedure ABC. The first step is to estimate the number of claims related to procedure ABC in all 575 drawers. The results on the next page were obtained and are stored in data file DATAUNIV.TXT.

<u>Sampled Drawer</u>	<u>Number of Claims Related to Procedure ABC</u>	
1	9	
2	12	
3	9	<u>Sample summary</u>
.	.	$\bar{x} = 10.33$
.	.	$s = 2.75$
.	.	
70	<u>10</u>	
Total	723	

Data file DATAUNIV.TXT

<continued>

1	9	36	9
2	12	37	12
3	9	38	8
4	6	39	6
5	12	40	14
6	13	41	9
7	10	42	14
8	9	43	14
9	10	44	6
10	6	45	14
11	13	46	12
12	7	47	9
13	12	48	10
14	7	49	9
15	12	50	9
16	12	51	8
17	10	52	14
18	9	53	9
19	10	54	14
20	10	55	8
21	13	56	12
22	10	57	14
23	10	58	10
24	10	59	12
25	6	60	9
26	8	61	12
27	14	62	12
28	22	63	14
29	10	64	10
30	8	65	10
31	8	66	6
32	8	67	9
33	12	68	8
34	10	69	12
35	8	70	10

**NOTE: This file contains 70
lines.**

The following computer output was obtained using the **Unrestricted Variable Appraisal** program. In the **Universe Size** box, the size of the larger universe (575 file drawers here) was used. With this procedure, you are able to see the estimated size of the universe of interest ($\hat{N} = 5,939$) in this output.

```

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 2/24/2010      VARIABLE UNRESTRICTED APPRAISAL      Time: 12:12
                     AUDIT/REVIEW: First Sample

DATA FILE USED: C:\Temp\DATAUNIV.TXT

      SAMPLE SIZE      VALUE OF SAMPLE      NONZERO ITEMS
           70              723.00              70

----- E X A M I N E D -----
MEAN / UNIVERSE      10.33      575
STANDARD DEVIATION      2.75
STANDARD ERROR          .31
SKEWNESS              1.02
KURTOSIS              5.95
POINT ESTIMATE      5,939  <--estimated universe
                               size

                                CONFIDENCE LIMITS
                                80% CONFIDENCE LEVEL
LOWER LIMIT              5,710
UPPER LIMIT              6,168
PRECISION AMOUNT          229
PRECISION PERCENT         3.86%
T-VALUE USED            1.293941609194

                                90% CONFIDENCE LEVEL
LOWER LIMIT              5,644
UPPER LIMIT              6,234
PRECISION AMOUNT          295
PRECISION PERCENT         4.97%
T-VALUE USED            1.667238548669

                                95% CONFIDENCE LEVEL
LOWER LIMIT              5,586
UPPER LIMIT              6,292
PRECISION AMOUNT          353
PRECISION PERCENT         5.95%
T-VALUE USED            1.994945415107

```

The next step is to independently obtain a random sample from the population of interest to appraise the variable(s) of interest. It was decided to sample 55 claims related to procedure ABC and record the amount in error for each sampled claim. The results are stored in data file DATAVAR.TXT.

Data file DATAVAR.TXT

< - continued - >

1	15.05		
2	14.05	29	16.07
3	18.55	30	20.10
4	13.62	31	17.29
5	18.78	32	21.38
6	6.97	33	16.60
7	17.17	34	6.98
8	23.75	35	16.85
9	19.32	36	19.25
10	21.32	37	20.30
11	25.17	38	29.51
12	14.12	39	6.95
13	15.85	40	19.00
14	13.50	41	13.93
15	14.62	42	12.45
16	6.97	43	15.23
17	22.62	44	15.09
18	6.97	45	6.98
19	15.61	46	20.48
20	12.69	47	6.98
21	21.45	48	16.47
22	21.62	49	19.75
23	17.95	50	6.98
24	14.65	51	17.65
25	6.97	52	15.26
26	18.89	53	6.98
27	15.34	54	14.15
28	6.98	55	22.05

NOTE: This file contains 55 lines.

This file is used as input to the **Unrestricted Variable Appraisal** program. The user can enter any value in the **Universe Size** box since this value has no effect on the results produced by the **Unknown Universe Size** program. The following computer output was produced.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 2/24/2010 VARIABLE UNRESTRICTED APPRAISAL Time: 13:12
 AUDIT/REVIEW: Second Sample

DATA FILE USED: C:\Temp\DATAVAR.TXT

SAMPLE SIZE	VALUE OF SAMPLE	NONZERO ITEMS
55	860.76	55

Any value can be
used for universe

----- D I F F E R E N C E -----

MEAN / UNIVERSE	15.65	1,000
STANDARD DEVIATION	5.45	
STANDARD ERROR	.71	NOTE: Input to Unknown
SKEWNESS	-.14	Universe Size program.
KURTOSIS	2.63	
POINT ESTIMATE	15,650	

CONFIDENCE LIMITS

80% CONFIDENCE LEVEL

LOWER LIMIT	14,723
UPPER LIMIT	16,577
PRECISION AMOUNT	927
PRECISION PERCENT	5.92%
T-VALUE USED	1.297426488209

90% CONFIDENCE LEVEL

LOWER LIMIT	14,455
UPPER LIMIT	16,846
PRECISION AMOUNT	1,196
PRECISION PERCENT	7.64%
T-VALUE USED	1.673564906352

95% CONFIDENCE LEVEL

LOWER LIMIT	14,218
UPPER LIMIT	17,082
PRECISION AMOUNT	1,432
PRECISION PERCENT	9.15%
T-VALUE USED	2.004879288188

Comment: This is the same example used in the RAT-STATS User's Guide illustration. The following computer output is produced by the Unknown Universe Size program:

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES
Date: 2/24/2010 VARIABLE APPRAISAL WITH UNKNOWN UNIVERSE SIZE Time: 13:48
AUDIT/REVIEW: Variable Unknown Universe Size

```

= = = = = I N P U T = = = = =
                SAMPLE TO          SAMPLE FOR
                ESTIMATE POPULATION  VARIABLE ATTRIBUTE
UNIVERSE                        575
SAMPLE                          70          55
MEAN                          10.33       15.65
STANDARD DEVIATION            2.75        5.45

= = = = = E S T I M A T I O N = = = = =
                80% CONFIDENCE      90% CONFIDENCE      95% CONFIDENCE
POINT ESTIMATE       $\hat{T}$  --> 92,957          92,957          92,957
STANDARD ERROR           5,152          5,152          5,152
LOWER LIMIT          86,355          84,483          82,859
UPPER LIMIT          99,560          101,431          103,055
PRECISION AMOUNT       6,603          8,474          10,098
PRECISION PERCENT      7.10%          9.12%          10.86%
Z-VALUE USED          1.281551565545    1.644853626951    1.959963984540

```

Discussion. The estimated total dollars in the universe is equal to

(estimated universe size)(mean of the variable sample)

$$= [(10.33)(575)](15.65) = \$92,957. \text{ This is } \hat{T}.$$

Using the formula section, the value of se_1 is

$$\frac{(2.75)(575)}{\sqrt{70}} \sqrt{1 - \frac{70}{575}} = 177.118$$

and the value of se_2 is

$$\frac{5.45}{\sqrt{55}} \sqrt{1 - \frac{55}{5939}} = .7315$$

The estimated variance of \hat{T} is

$$[(15.65)(177.118)]^2 + [(5939)(.7315)]^2 - [(177.118)(.7315)]^2 = 26,540,250.$$

The estimated standard error of \hat{T} is $\sqrt{26,540,250} = \$5,152$.

The PRECISION AMOUNT at the 90% confidence level is $(5,152)(1.644853626951) = \$8,474$

(highlighted). This amount is 9.12% of the point estimate, since $8,474/92,957 = .0912$.

The corresponding confidence interval is $92,957 \pm 8,474$; that is, \$84,483 to \$101,431.

FORMULAS

Given: Larger universe size: N_1 . Sample to estimate universe size: sample size is n_1 , mean is

\bar{x}_1 , and standard deviation is s_1 .

Sample to estimate variable of interest: sample size is n_2 , mean is \bar{x}_2 , and standard

deviation is s_2 .

1. Estimate of universe (of interest) size

$$\hat{N} = N_1 \cdot \bar{x}_1$$

2. Overall estimate for variable total (e.g., total error amount)

$$\hat{T} = \hat{N} \cdot \bar{x}_2$$

3. Estimated variance of \hat{T}

$$v(\hat{T}) = [\bar{x}_2 \cdot se_1]^2 + [\hat{N} \cdot se_2]^2 - [se_1 \cdot se_2]^2$$

where

$$se_1 = \frac{s_1 N_1}{\sqrt{n_1}} \sqrt{1 - \frac{n_1}{N_1}} \quad \text{and} \quad se_2 = \frac{s_2}{\sqrt{n_2}} \sqrt{1 - \frac{n_2}{\hat{N}}}$$

4. Approximate 95% confidence interval for universe total (T):

$$\hat{T} \pm 1.959963984540\sqrt{v(\hat{T})}$$

NOTE: For a 90% confidence interval, replace 1.959963984540 with 1.644853626951 and for an 80% confidence interval replace 1.959963984540 with 1.281551565545.

Reference:

Estimated variance: Kendall's Advanced Theory of Statistics, Volume 1, 5th ed., Alan Stuart and J. Keith Ord, New York: Oxford University Press, 1987, page 343, exercise 10.23.

SAMPLE SIZE DETERMINATION

A commonly encountered question in auditing is “How large a sample is necessary?” When using an unrestricted (simple random) sample, this depends on the desired precision of the point estimate. The programs in this section are listed below and are concerned with determining sample sizes for various data types and sample strategies.

- Variable
 - Unrestricted Using a Probe Sample
 - Unrestricted Using Estimated Error Rate
 - Stratified (Total Sample Size Known)
 - Stratified (Total Sample Size Unknown)

Variable Sample Size Determination

This RAT-STATS module can be used for two situations.

Situation 1: The program will help select the necessary sample size for an unrestricted or stratified variable appraisal. The program output includes sample sizes for each stratum that will provide precision percentages of 1%, 2%, 5%, 10%, 20% and “Other.” When selecting “Other,” the user will be prompted to enter the desired precision percentage. The user may also select any combination of the following confidence levels: 80%, 90%, 95%, and 99%.

Situation 2: The program also allows the user to determine the optimum distribution of a sample among strata when the overall sample size has already been determined. It will allocate the larger samples to those strata that are larger in size and/or contain a larger amount of variation (are nonhomogeneous). Any combination of the confidence levels 80%, 90%, 95%, and 99% can be selected.


Variable Sample Size Determination - Unrestricted Using a Probe Sample

This program allows the user to estimate sample sizes for specified precision percentages and specified confidence levels. The user has the option of having the program read a probe sample file to obtain an estimate of the universe mean and standard deviation or input these two estimates directly without reading a probe sample file. The probe sample may be stored in a text file, an Excel spreadsheet, or an Access table.

Example 1. This example illustrates Situation 1. A probe sample of 25 examined values was obtained. The audit objective was to determine the necessary sample sizes when estimating the total examined amount for the universe of 100,000 items. The probe sample (SAMPDATA.TXT) is shown below. The sample mean is \$400 and the sample standard deviation is \$50.

321
382
453
459
343
388
313
420
407
395
441
448
447
333
357
395
477
391
356
368
376
350
461
472
447

The input screen and resulting text file output are shown on the following page.



The dialog box is titled "Variable Sample Size Determination". It contains several sections for user input:

- Probe Sample Format:** A group box containing four radio buttons: "Text File", "Excel Spreadsheet", "Access Table", and "No Probe Sample File".
- Confidence Level:** A group box containing five checkboxes: "80%", "95%", "90%", "99%", and "All".
- Precision:** A group box containing five checkboxes: "1%", "10%", "2%", "15%", "5%", "25%", and "All".
- Universe Size:** A text input field with the value "100,000".
- OUTPUT TO:** A group box containing four radio buttons: "Text File and Screen", "Printer and Screen", "Text File, Printer, and Screen", and "Screen Only".
- Buttons:** On the left, there are three buttons: "HELP" (orange), "Main Menu" (cyan), and "EXIT" (pink). On the right, there is an "OK" button highlighted with a pink border.

The following text file output is obtained using the previous screen. A sample size under 30 will be flagged using “(*)” and the note immediately following the sample sizes will appear.

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/11/2010

Sample Size Determination

Time: 21:52

		Confidence Level			
		80%	90%	95%	99%
Precision Level	1%	256	421	597	1026
	2%	64	106	150	259
	5%	10 (*)	17 (*)	24 (*)	41
	10%	3 (*)	4 (*)	6 (*)	10 (*)
	15%	1 (*)	2 (*)	3 (*)	5 (*)
	25%	---	1 (*)	1 (*)	2 (*)

Estimated Mean: 400.00

Estimated Std. Deviation: 50.00

Universe Size: 100,000

NOTE (*): One or more sample sizes were under 30. The generated sample sizes were the result of mathematical formulas and did not incorporate management decisions concerning the purpose of the sample or current organizational sampling policies. You may need to increase the sample sizes in order to be in compliance with organizational objectives.

Explanation of Output

The output for each cell in the output table will consist of (1) the necessary sample size or (2) the text “- - -”. The necessary sample size is the number of sample items necessary to obtain the specified sample precision at the specified confidence level. For example, in this illustration, a sample size of 106 is necessary to obtain a point estimate having a precision percentage of plus or minus 2% using a 90% confidence level. If the calculated sample size is 0, a text value of “- - -” will appear in this cell. This occurred in the lower left cell for the sample illustration.

The output also contains the estimated mean and standard deviation, along with the specified universe size.

FORMULAS

Let PREC = the precision percentage (e.g., 1 for 1%, 10 for 10%)

ZVAL = the value from the standard normal (Z) distribution having a right-tail area equal to $(100 - \text{Confidence Level})/2$, where the right-tail area is expressed as a proportion between 0 and 1.

ZVAL is 1.281551565545(80%), 1.644853626951 (90%), 1.959963984540 (95%), and 2.575829303549 (99%).

N = the universe size

Mean = estimated universe mean obtained from the probe sample or specified by the user

StdDev = estimated universe standard deviation obtained from the probe sample or specified by the user

E = maximum error = $(\text{PREC}/100) \cdot \text{Mean} \cdot N$

For each selected value of PREC and ZVAL, the sample size is

$$n = \frac{(\text{StdDev} \cdot N)^2}{(E / ZVAL)^2 + N \cdot (\text{StdDev})^2}$$

The value of n is rounded up or down to the nearest integer.

Variable Sample Size Determination - Unrestricted Using Expected Error Rate

This procedure estimates the mean and standard deviation of the difference (error) amounts by assuming (1) any item found to be in error is 100% in error and (2) the mean and standard deviation of the *nonzero* error amounts is the same as the mean and standard deviation of the reported (examined) amounts. The mean and standard deviation of the error amounts are estimated by assuming the percentage of nonzero errors in the error population is equal to the expected error rate (one of the input values) and the nonzero errors resemble the reported amounts; that is, the mean and standard deviation of the nonzero errors are equal to the mean and standard deviation of the reported amounts.

Comment. Even though these assumptions may not be entirely true, this procedure will often give more reliable sample size estimates than those obtained using the Variable Unrestricted (Using Reported Amounts) module since the expected number of zero values in the error population is factored into the sample size calculation.

Example 2. This example illustrates another method of dealing with Situation 1. The estimated error rate is 15% for a universe of 10,000 transactions. The total reported amount is \$3,000,000 and the standard deviation of the reported amounts is \$125. Consequently, the mean reported amount is \$300. Of interest is the required sample size necessary in order to obtain plus or minus 15% using a 90% confidence level. The corresponding input screen follows where 25% was specified for the “Other” precision level.

Variable Sample Size Using Estimated Error Rate

Universe Size 10,000

Anticipated Error Rate
NOTE: Enter 5 for 5%, 10 for 10%, etc. 15 %

Reported Amounts

Total Amount 3,000,000.00

Standard Deviation 125.00

Confidence Level

☒ 80% ☒ 95%

☒ 90% ☒ 99%

☒ All

Precision

☒ 1% ☒ 10%

☒ 2% ☒ 15%

☒ 5% ☒ 25%

☒ All

OUTPUT TO

☐ Text File and Screen

☐ Printer and Screen

☒ Text File, Printer, and Screen

☐ Screen Only

HELP

Main Menu

EXIT

OK

The text file output shown on the next page is obtained using this input screen. A sample size under 30 will be flagged using “(*)” in the program output and a note informing the user of this fact will also appear in the program output immediately following the sample sizes.

DEPARTMENT OF HEALTH & HUMAN SERVICES

OIG - OFFICE OF AUDIT SERVICES

Date: 5/22/2010

Sample Size Determination

Time: 10:14

		Confidence Level			
		80%	90%	95%	99%
Precision Level	1%	9181	9486	9633	9784
	2%	7370	8219	8676	9188
	5%	3095	4248	5119	6443
	10%	1008	1559	2077	3117
	15%	474	758	1044	1675
	25%	176	287	403	675

Universe Size: 10,000

Anticipated Error Rate: 15%

Reported Amounts - - Total Amount: 3,000,000.00

Standard Deviation: 125.00

Difference Values - - Estimated Mean: 45.00

Estimated Standard Deviation: 117.55

Explanation of Output

The output for each cell in the output table will consist of (1) the necessary sample size or (2) the text “- - -”. The necessary sample size is the number of sample items necessary to obtain the specified sample precision at the specified confidence level. For example, in this illustration, a sample size of 287 is necessary to obtain a point estimate having a precision percentage of plus or minus 15% using a 90% confidence level. If the calculated sample size is 0, a text value of “- - -” will appear in this cell.

The output also contains the estimated mean and standard deviation of the difference (error) values. For this illustration, the estimated mean and standard deviation are \$45.00 and \$117.55, respectively.

FORMULAS

Let PREC = the precision percentage (e.g., 1 for 1%, 10 for 10%)

ZVAL = the value from the standard normal (Z) distribution having a right-tail area equal to $(100 - \text{Confidence Level})/2$, where the right-tail area is expressed as a proportion between 0 and 1.

ZVAL is 1.281551565545(80%), 1.644853626951 (90%), 1.959963984540 (95%), and 2.575829303549 (99%).

N = the universe size (input)

T_R = the total reported amount (input)

μ_R = mean reported amount = T_R / N

σ_R = standard deviation of reported amounts (input)

\hat{p} = the estimated error rate (input)

$\hat{\mu}_D$ = the estimated mean of the difference (error) values = $\hat{p}\mu_R$

$\hat{\sigma}_D$ = estimated standard deviation of the difference (error) values

$$= \sqrt{\hat{p}[\sigma_R^2 + (1 - \hat{p})\mu_R^2]}$$

E = maximum error = $(\text{PREC}/100) \cdot \hat{\mu}_D \cdot N$

For each selected value of PREC and ZVAL, the sample size is

$$n = \frac{(\hat{\sigma}_D \cdot N)^2}{(E / ZVAL)^2 + N \cdot (\hat{\sigma}_D)^2}$$

The value of n is rounded up or down to the nearest integer.

Variable Sample Size Determination - Stratified

Stratified Sample Sizes - Total Sample Size is Unknown

Example 3. This example illustrates Situation 1. Two strata have been defined: The high-income stratum ($N_1 = 100,000$ items) and the low-income stratum ($N_2 = 500,000$ items). Of interest is the total audit (claimed) amount for the universe. For the high-income stratum, the estimated mean of the audited amounts is \$10,000 and the estimated standard deviation is \$5,000. These values for the low-income stratum are \$5,000 (mean) and \$4,000 (standard deviation). At a confidence level of 95%, what sample size is required to obtain a precision percentage of $\pm 10\%$?

Solution. The following input screen was used for this example.

The screenshot shows a software window titled "Stratified Variable Sample Size Determination". The window contains several input fields and checkboxes. The "Number of strata" field is set to 2, with a note "(maximum = 12)". Under "Total sample size is known: Determine the optimum allocation", the radio button is unselected. Under "Total sample size is unknown", the radio button is selected. The "Confidence Level" section has checkboxes for 80%, 95%, 90%, 99%, and All, all of which are checked. The "Precision" section has checkboxes for 1%, 10%, 2%, 15%, 5%, 25%, and All, all of which are checked. On the left side, there are three buttons: "HELP" (orange), "Main Menu" (cyan), and "EXIT" (pink). In the center, under "OUTPUT TO", there are four radio buttons: "Text File and Screen" (selected), "Printer and Screen", "Text File, Printer, and Screen", and "Screen Only". On the right side, there is a large "OK" button with a pink border.

The following output is obtained using the previous screen. If one or more of the sample sizes are under 30, the note immediately following the total sample sizes will appear.

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES

Date: 5/19/2010 Sample Size Determination Time: 12:02

THE ESTIMATES ARE BASED ON THE FOLLOWING ENTRIES:

NBR	DESCRIPTION	-- MEAN --	-- STD.DEV. --	-- UNIVERSE --	-- RATIO --
1	High Income	10,000.00	5,000.00	100,000	20.00%
2	Low Income	5,000.00	4,000.00	500,000	80.00%
- TOTALS -		5,833.33	4,579.54	600,000	

= = = = =

Sample Sizes for Stratum 1: High Income

		Confidence Level			
		80%	90%	95%	99%
Precision Level	1%	1653	2699	3795	6406
	2%	418	687	972	1669
	5%	67	111	157	271
	10%	17 (*)	28 (*)	40	68
	15%	8 (*)	13 (*)	18 (*)	31
	25%	3 (*)	5 (*)	7 (*)	11 (*)

Sample Sizes for Stratum 2: Low Income

		Confidence Level			
		80%	90%	95%	99%
Precision Level	1%	6611	10793	15180	25624
	2%	1671	2745	3888	6676
	5%	268	442	627	1081
	10%	68	111	157	271
	15%	30	50	70	121
	25%	11 (*)	18 (*)	26 (*)	44

Total Sample Sizes

		Confidence Level			
		80%	90%	95%	99%
Precision Level	1%	8264	13492	18975	32030
	2%	2089	3432	4860	8345
	5%	335	553	784	1352
	10%	85	139	197	339
	15%	38	63	88	152
	25%	14 (*)	23 (*)	33	55

NOTE (*): One or more sample sizes were under 30. The generated sample sizes were the result of mathematical formulas and did not incorporate management decisions concerning the purpose of the sample or current organizational sampling policies. You may need to increase the sample sizes in order to be in compliance with organizational objectives.

If any of the calculated samples sizes exceeds the corresponding universe size, the program will conclude with the following reminder:

NOTE (#): The formulas calculated a sample size greater than the universe size. The program reduced the calculated sample size to the universe size. The additional sampling units were then distributed among the remaining strata based on optimal allocation formulas.

Discussion. For 10% precision and 95% confidence, the total sample size required is $n = 197$ with $n_1 = 40$ items to be obtained from the high-income stratum and $n_2 = 157$ from the low-income stratum. Consequently, a 95% confidence interval based on these sample sizes should result in a precision percentage of $\pm 10\%$. This assumes that the resulting sample means and standard deviations are the same as the values used as input to this program.

To demonstrate this, a data set was constructed that contained 40 items from stratum 1 with a sample mean and standard deviation of \$10,000 and \$5,000, respectively, and 157 items from stratum 2 with a sample mean and standard deviation of \$5,000 and \$4,000, respectively. When this data set (named STRATA.TXT) was used as input to the STRATIFIED VARIABLE APPRAISAL module, the computer output on the next page was obtained. In the final portion of the output, notice that the resulting point estimate for the universe total is 3,500,000,000. At the 95% confidence level, the precision amount is 349,043,863 and is in fact (approximately) 10% of the point estimate.

DEPARTMENT OF HEALTH & HUMAN SERVICES
 OIG - OFFICE OF AUDIT SERVICES
 Date: 5/23/2010 STRATIFIED VARIABLE APPRAISAL Time: 10:36
 AUDIT/REVIEW: Two Strata Example

DATA FILE USED: C:\Temp\STRATA.TXT

STRATUM NUMBER	SAMPLE SIZE	VALUE OF SAMPLE	NONZERO ITEMS
1	40	400,000.00	40
2	157	785,000.00	157
TOTALS	197	1,185,000.00	197

----- D I F F E R E N C E -----

Stratum 1 MEAN / UNIVERSE 10,000.00 100,000
 STANDARD DEVIATION 5,000.04 ← Approx. 5,000
 STANDARD ERROR 790.42
 SKEWNESS -.22
 KURTOSIS 2.30
 STANDARD ERROR (MEAN) 790.42
 STANDARD ERROR (TOTAL) 79,041,756
 POINT ESTIMATE 1,000,000,000

CONFIDENCE LIMITS

80% CONFIDENCE LEVEL

LOWER LIMIT 896,958,117
 UPPER LIMIT 1,103,041,883
 PRECISION AMOUNT 103,041,883
 PRECISION PERCENT 10.30%
 T-VALUE USED 1.303638588621

90% CONFIDENCE LEVEL

LOWER LIMIT 866,824,512
 UPPER LIMIT 1,133,175,488
 PRECISION AMOUNT 133,175,488
 PRECISION PERCENT 13.32%
 T-VALUE USED 1.684875121711

95% CONFIDENCE LEVEL

LOWER LIMIT 840,122,958
 UPPER LIMIT 1,159,877,042
 PRECISION AMOUNT 159,877,042
 PRECISION PERCENT 15.99%
 T-VALUE USED 2.022690920037

Stratum 2 MEAN / UNIVERSE 5,000.00 500,000
 STANDARD DEVIATION 3,999.81 ← Approx. 4,000
 SKEWNESS .87
 KURTOSIS 3.30
 STANDARD ERROR (MEAN) 319.17
 STANDARD ERROR (TOTAL) 159,584,887
 POINT ESTIMATE 2,500,000,000

				CONFIDENCE LIMITS
				80% CONFIDENCE LEVEL
LOWER LIMIT				2,294,613,944
UPPER LIMIT				2,705,386,056
PRECISION AMOUNT				205,386,056
PRECISION PERCENT				8.22%
T-VALUE USED				1.287001917850
				90% CONFIDENCE LEVEL
LOWER LIMIT				2,235,938,079
UPPER LIMIT				2,764,061,921
PRECISION AMOUNT				264,061,921
PRECISION PERCENT				10.56%
T-VALUE USED				1.654679995672
				95% CONFIDENCE LEVEL
LOWER LIMIT				2,184,773,966
UPPER LIMIT				2,815,226,034
PRECISION AMOUNT				315,226,034
PRECISION PERCENT				12.61%
T-VALUE USED				1.975287507703
OVERALL	POINT ESTIMATE / UNIVERSE	3,500,000,000	600,000	
	STANDARD ERROR	178,086,876		
				CONFIDENCE LIMITS
				80% CONFIDENCE LEVEL
LOWER LIMIT				3,271,772,485
UPPER LIMIT				3,728,227,515
PRECISION AMOUNT				228,227,515
PRECISION PERCENT				6.52%
Z-VALUE USED				1.281551565545
				90% CONFIDENCE LEVEL
LOWER LIMIT				3,207,073,156
UPPER LIMIT				3,792,926,844
PRECISION AMOUNT				292,926,844
PRECISION PERCENT				8.37%
Z-VALUE USED				1.644853626951
				95% CONFIDENCE LEVEL
LOWER LIMIT				3,150,956,137
UPPER LIMIT				3,849,043,863
PRECISION AMOUNT				349,043,863
PRECISION PERCENT				9.97% ←
Z-VALUE USED				1.959963984540

Comments

- (1) When the sample of size $n = 197$ is obtained, the values of the sample mean and standard deviation will likely not be exactly those specified in the input to this program.

Consequently, the best the user can hope for is that the resulting precision percentage will be approximately 10%.

- (2) For the preceding example, the specified precision was 10% of the point estimate. The point estimate for the universe total was 3,500,000,000. In the formula section, E is the desired precision amount expressed as a percentage of the point estimate for the universe total. Here this would be $E = 350,000,000$.
- (3) For situations in which you do not have an estimate of the universe standard deviation (σ) from previous audit results, a rough approximation for σ can be obtained for each stratum by estimating (1) the largest value (L) that you expect to see in the sample for this stratum and (2) the smallest value (S) that you expect to see in this stratum. Then, the approximate value of σ for this stratum is $\hat{\sigma} = \frac{L - S}{4}$. In the previous example, if the largest audit amount that you expect to see in the LOW INCOME stratum is $L = \$15,000$ and the smallest value is $S = \$1,000$, then the estimated standard deviation is $\hat{\sigma} = (15,000 - 1,000)/4 = \$3,500$.

Stratified Sample Sizes - Total Sample Size is Known

Example 4. This is an illustration of situation 2. The situation is the same as that described in Example 3, which used two strata -- the high-income stratum and the low-income stratum. The total sample size is set at 500. The input screen on the following page was used for this example. Notice that the user is unable to set the precision percentages for this situation.

Stratified Variable Sample Size Determination

Number of strata
(maximum = 12)

☒ Total sample size is known:
Determine the optimum allocation
☐ Total sample size is unknown

Total sample size

Confidence Level

☒ 80% ☒ 95%
☒ 90% ☒ 99%
☒ All

HELP **Main Menu** **EXIT**

OUTPUT TO

☒ Text File and Screen
☐ Printer and Screen
☐ Text File, Printer, and Screen
☐ Screen Only

OK

The following estimates were used as input to the program:

Stratum	Estimated Mean	Estimated Standard Deviation	Estimated Universe Size
High Income	10,000	5,000	100,000
Low Income	5,000	4,000	500,000

The program output on the next page is obtained. Notice that the resulting strata ratios (i.e., 20% and 80%) are identical to those obtained in Example 3.

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES

Date: 5/23/2010

Sample Size Determination

Time: 13:07

THE ESTIMATES ARE BASED ON THE FOLLOWING ENTRIES:

NBR	DESCRIPTION	-- MEAN --	-- STD.DEV. --	-- UNIVERSE --
1	High Income	10,000.00	5,000.00	100,000
2	Low Income	5,000.00	4,000.00	500,000
- TOTALS -		5,833.33	4,579.54	600,000
=====				

Precision Values:

NOTE: See the Discussion section.

Confidence Level	80%	90%	95%	99%
	4.09%	5.25%	6.26%	8.22%

The following sample sizes are based on a total sample size of 500.

Stratum 1: High Income

Sample	
Size	Ratio
100	20.00%

Stratum 2: Low Income

Sample	
Size	Ratio
400	80.00%

Discussion. The two sample sizes are $n_1 = 100$ and $n_2 = 400$, which total $n = 500$. For this example, $\sum N_i \hat{\sigma}_i^2$ is $(100,000)(5,000) + (500,000)(4,000) = 2,500,000,000$. Call this SUM.

The ratio value for stratum 1 is $(100,000)(5,000)$ divided by SUM; that is .2. So, 20% of the sample size is allocated to stratum 1; that is, n_1 is $(500)(.2) = 100$. Similarly, the ratio for stratum 2 is .8 and n_2 is $(500)(.8) = 400$. **NOTE:** This same discussion applies to Example 3.

What is the precision amount for this sampling design? This will be the value obtained by the **Stratified Variable Appraisal** program using these sample sizes and estimated standard deviations. This formula (borrowed from the **Stratified Variable Appraisal** formula section) is contained in the formula section to follow. For this example, the precision amount will be

$$1.95996 \sqrt{100,000^2 \left(\frac{100,000 - 100}{100,000} \right) \frac{5,000^2}{100} + 500,000^2 \left(\frac{500,000 - 400}{500,000} \right) \frac{4,000^2}{400}}$$

$$= 219,038,136.$$

The estimated universe total is

$$\hat{T} = \Sigma(\text{stratum mean})(\text{stratum size}) = (10,000)(100,000) + (5,000)(500,000) = 3,500,000,000.$$

The resulting precision percentage is

$$100 \cdot (219,038,136 / 3,500,000,000) = 6.26\%.$$

This value is called PERC in the formula section to follow and matches with the highlighted value in the computer output.

FORMULAS

Total Sample Size (n) is Known

Notation

L = Number of strata

N_i = the universe size for the i-th stratum

$(StdDev)_i$ = estimated universe standard deviation for the i-th stratum

$$SUM = \sum_{i=1}^L N_i \cdot (StdDev)_i$$

$$(Ratio)_i = [N_i \cdot (StdDev)_i] / SUM$$

The resulting sample size allocated to the i-th stratum is $n_i = n \cdot (Ratio)_i$.

Total Sample Size (n) is Unknown

Notation

L = Number of strata

N_i = the universe size for the i -th stratum

N = the total universe size = $\sum_{i=1}^L N_i$

$(Mean)_i$ = estimated universe mean for the i -th stratum

UnivTotal = estimated universe total = $\sum_{i=1}^L N_i \cdot (Mean)_i$

$(StdDev)_i$ = estimated universe standard deviation for the i -th stratum

$$SUM1 = \sum_{i=1}^L N_i \cdot (StdDev)_i$$

$$SUM2 = \sum_{i=1}^L N_i \cdot (StdDev)_i^2$$

$$(Ratio)_i = [N_i \cdot (StdDev)_i] / SUM1$$

PREC = the precision percentage (e.g., 1 for 1%, 10 for 10%)

ZVAL = the value from the standard normal (Z) distribution having a right-tail area equal to $(100 - \text{Confidence Level})/2$, where the right-tail area is expressed as a proportion between zero and one.

ZVAL is 1.281551565545 (80%), 1.644853626951 (90%), 1.959963984540 (95%), and 2.575829303549 (99%).

$$E = \text{the precision amount} = (PREC/100) \cdot (\text{UnivTotal})$$

For each selected value of PREC and ZVAL,

- (1) the total sample size is

$$n = \frac{(SUM1)^2}{(E / ZVAL)^2 + SUM2}$$

- (2) the sample size allocated to the i-th stratum is

$$n_i = n \cdot (\text{Ratio})_i$$

Comments

1. In the preceding calculation, the value of n is treated as a floating point number (e.g., n = 487.263) and the strata sample sizes (n_i) are calculated using this value. The n_i values are then rounded up to the nearest integer. After all strata sample sizes have been determined, n is reset to the sum of the n_i .
2. If the computed sample size for stratum i (n_i) is larger than the universe size N_i , then n_i is set equal to N_i . The remaining sample sizes are then obtained by applying the above formula and (1) omitting the i-th stratum in the denominator and (2) replacing n with n - N_i (the total sample size for the remaining L-1 strata).

The precision percentage at the 95% confidence level is \pm PERC, where

$$PERC = \frac{1.959963984540}{\hat{T}} \sqrt{\sum_{i=1}^L N_i^2 \left(\frac{N_i - n_i}{N_i} \right) \frac{(StdDev)_i^2}{n_i}}$$

and where \hat{T} is the estimated total for the universe. The value of \hat{T} is obtained by multiplying

N_i by the estimated mean for stratum i and summing over the L strata; that is, $\hat{T} = \sum_{i=1}^L N_i \hat{\mu}_i$

NOTE: Replace 1.959963984540 with 1.281551565545 for an 80% interval, 1.644853626951 for a 90% interval, and 2.575829303549 for a 99% interval.

Attribute Sample Size Determination

This program determines the sample size for an attribute simple random sample. The sample size is determined for specified degrees of precision (using the desired width of the confidence intervals) and for various levels of confidence. The resulting sample size is the smallest sample size capable of meeting the specified precision requirement at each of the specified confidence levels. The user may select any combination of the following confidence levels: 80%, 90%, 95%, and 99%.

Confidence intervals for attribute sampling are exact and are based on the hypergeometric distribution. As a result, such confidence intervals are usually not symmetric about the point estimate. For example, the point estimate might be 3% and the corresponding 95% confidence interval might be 2% to 6%. For this illustration, the width of the confidence interval is 4% and the confidence level is 95%. Consequently, attribute confidence intervals differ from the usual interval obtained by deriving the point estimate plus or minus the estimated precision, where the estimated precision is half the width of the resulting confidence interval. Because of this, the “desired precision” for the attribute sampling procedure must be specified as the desired width (rather than the half-width) of the confidence interval.

An approximate confidence interval for a universe proportion (discussed in many introductory statistics textbooks) is based on the normal approximation. This particular interval follows the “usual” procedure where the confidence interval is equal to (point estimate) \pm (estimated precision); that is, this interval is symmetric about the point estimate. However, this confidence interval is approximate and is unreliable whenever the estimated proportion is very small or very

large, unless the sample size is extremely large. The confidence interval using the RAT-STATS attribute sample size module discussed here is always exact.

The input screen includes (1) the size of the universe and (2) the anticipated rate of occurrence in the universe. This rate of occurrence is generally estimated from past experience, either from similar systems or a past review of this universe. If no information concerning the rate of occurrence is available, the most conservative procedure is to specify 50% for this value. If the actual rate of occurrence differs from the user-specified rate of occurrence, this in no way affects the sample's validity but the resulting precision (confidence interval width) may not meet the desired precision requirement.

Example 5. An audit is to be carried out using a universe of $N = 10,000$ documents to determine what proportion (p) of the documents do not have the proper approval signature. A confidence level of 95% will be used. It is estimated that 20% of the documents will not have the proper signature. Consequently, the estimate of p is $\hat{p} = .20$.

NOTE: This may be a rough guess if little information regarding this estimate is available from previous audit experience. If the user has no idea as to the value of p , $\hat{p} = .5$ should be used. This will produce the largest possible sample size (for fixed values of N and precision range) but the user will be guaranteed that the resulting confidence interval will meet the desired precision range.

Suppose that the desired precision range is 6%. This is equal to the desired value of (upper confidence limit - lower confidence limit). If the confidence limits were symmetric about the point estimate, the user would have specified the precision as $\pm 3\%$ for this situation, where 3% is half the width of the resulting confidence interval. Since the exact procedure used in this

program usually does not produce an interval symmetric about the point estimate, the user must specify the desired total width of the confidence interval. The following input screen is used for this example:

Attribute Sample Size Determination

Confidence Level

☒ 80% ☒ 95%

☒ 90% ☒ 99%

☒ All

The "anticipated rate of occurrence" should be entered as a percentage; that is, enter 10 for 10%, 20 for 20%, and so on. The most conservative value is 50. The minimum value is 0.5% and the maximum value is 98%.

Anticipated Rate of Occurrence

Universe Size

The "desired precision range" for the universe error rate is the desired width of the confidence interval. Enter 5 for 5%, 10 for 10%, and so on. For example, if the confidence interval (10% to 16%) satisfies your precision requirements, enter "6" in the box. The minimum value is 1% and the maximum value is 99%.

Desired Precision Range

OUTPUT TO

☐ Text File and Screen

☐ Printer and Screen

☐ Text File, Printer, and Screen

☒ Screen Only

HELP

Main Menu

EXIT

OK

The resulting computer output (saved to a text file) is shown on the next page.

DEPARTMENT OF HEALTH & HUMAN SERVICES
OIG - OFFICE OF AUDIT SERVICES

Date: 5/19/2010 Sample Size Determination Time: 8:46

	Confidence Level			
	80%	90%	95%	99%
Sample Size	314	488	666	1,079

Anticipated Rate of Occurrence: 20%

Desired Precision Range: 6%

Universe Size: 10,000

Explanation of Output

The output for each cell in the output table will consist of (1) the necessary sample size or (2) the text “- - -”. The necessary sample size is the number of sample items necessary to obtain the specified sample precision at each confidence level. For example, in this illustration a sample size of 488 is necessary to obtain a confidence interval having a width of 6% using a 90% confidence level. If the calculated sample size is zero, a text value of “- - -” will appear in this cell.

Discussion. The necessary sample size (highlighted) is $n = 666$. As a result, after the sample of 666 is obtained, if the resulting point estimate is close to $\hat{p} = .20$, then the resulting 95% confidence interval for p should have a width approximately equal to .06 (such as .1710 to .2310, with a width of $.2310 - .1710 = .06$). If the resulting sample produced 133 documents not containing the proper signature, then the rate of occurrence in this sample would be $133/666$; that is, 20%.

The resulting confidence interval will have a width equal to .06 (i.e., 6%). This can be seen in the computer output below, obtained using the **Unrestricted Attribute Appraisal** module. The

width of this 95% confidence interval is 23.10% -17.10% = 6% (the desired precision range).

Notice that $\hat{p} = .20$ (i.e., 20%) is inside this interval (it always is), but it is not in the center.

Department of Health and Human Services
 OIG - Office of Audit Services
 Date: 5/19/2010 Single Stage Attribute Appraisal Time: 12:24
 AUDIT/REVIEW: Example

UNIVERSE SIZE	10,000
SAMPLE SIZE	666
CHARACTERISTIC(S) OF INTEREST	
QUANTITY IDENTIFIED IN SAMPLE	133
PROJECTED QUANTITY IN UNIVERSE	1,997
PERCENT	19.970%
STANDARD ERROR	
PROJECTED QUANTITY	150
PERCENT	1.498%
CONFIDENCE LIMITS	
80% CONFIDENCE LEVEL	
LOWER LIMIT - QUANTITY	1,805
PERCENT	18.050%
UPPER LIMIT - QUANTITY	2,202
PERCENT	22.020%
90% CONFIDENCE LEVEL	
LOWER LIMIT - QUANTITY	1,754
PERCENT	17.540%
UPPER LIMIT - QUANTITY	2,259
PERCENT	22.590%
95% CONFIDENCE LEVEL	
LOWER LIMIT - QUANTITY	1,710
PERCENT	17.100%
UPPER LIMIT - QUANTITY	2,310
PERCENT	23.100%

Example 6. Repeat Example 5 where no information is available regarding the proportion of documents not containing the proper signature.

Solution. Here, the user should enter 50% ($\hat{p} = .5$) in the Anticipated Error Rate box. The resulting computer output is shown below. The necessary sample size (highlighted) is now $n = 991$, approximately 50% larger than the previous sample size of 666.

```

                                DEPARTMENT OF HEALTH & HUMAN SERVICES
                                OIG - OFFICE OF AUDIT SERVICES
Date: 5/19/2010                Sample Size Determination                Time: 12:33

                                Confidence Level
                                80%      90%      95%      99%
Sample Size      466      725      987      1,580

Anticipated Rate of Occurrence: 50%
Desired Precision Range: 6%
Universe Size: 10,000

```

Discussion. Example 6 illustrates how using $\hat{p} = .5$ produces a very large value of n . **The user should be encouraged to use even a rough guess for the value of \hat{p} .** Using $\hat{p} = .5$ is a very conservative procedure because with a sample of size $n = 987$, quite likely the resulting confidence interval will have a width considerably less than the desired precision range of 6%. To illustrate, the computer output below was obtained when the sample of 989 documents produced 247 not containing the proper signature. Here, $\hat{p} = 247/989 = .250$ and the confidence interval width (using the highlighted values in the following computer output) is 5.21%. This value is less than 6%, but the user did have the guarantee that this value would be no more than 6%.

Department of Health and Human Services
OIG - Office of Audit Services
Date: 5/19/2010 Single Stage Attribute Appraisal Time: 12:43
AUDIT/REVIEW: Example

UNIVERSE SIZE	10,000
SAMPLE SIZE	987
CHARACTERISTIC(S) OF INTEREST	
QUANTITY IDENTIFIED IN SAMPLE	247
PROJECTED QUANTITY IN UNIVERSE	2,503
PERCENT	25.025%
STANDARD ERROR	
PROJECTED QUANTITY	131
PERCENT	1.310%
CONFIDENCE LIMITS	
80% CONFIDENCE LEVEL	
LOWER LIMIT - QUANTITY	2,334
PERCENT	23.340%
UPPER LIMIT - QUANTITY	2,678
PERCENT	26.780%
90% CONFIDENCE LEVEL	
LOWER LIMIT - QUANTITY	2,288
PERCENT	22.880%
UPPER LIMIT - QUANTITY	2,727
PERCENT	27.270%
95% CONFIDENCE LEVEL	
LOWER LIMIT - QUANTITY	2,249
PERCENT	22.490%
UPPER LIMIT - QUANTITY	2,770
PERCENT	27.700%

FORMULAS

In the discussion to follow, a sample item having the attribute of interest will be referred to as an item “in error.” Consequently, the universe proportion, p , will be the “error rate.”

Consider the case where the specified confidence level is 95%. The upper limit of the 95% confidence interval for the universe total is, say, k_2 , where k_2 is the largest value of k for which

$$\sum_{i=0}^x \frac{\binom{k}{i} \binom{N-k}{n-i}}{\binom{N}{n}} > .025$$

where N = universe size

n = sample size

k = total number of universe items in error

x = number of sample items in error

$.025 = [1 - (\text{confidence level})]/2$

NOTE: Here, the “confidence level” is expressed as .95.

The lower limit of the 95% confidence interval is, say, k_1 , where k_1 is the smallest value of k for which

$$\sum_{i=x}^n \frac{\binom{k}{i} \binom{N-k}{n-i}}{\binom{N}{n}} > .025$$

The resulting 95% confidence interval for the total number of universe items in error is k_1 to k_2 .

The procedure used to derive this confidence interval can be found in John P. Buonaccorsi (1987), "A Note on Confidence Intervals for Proportions in Finite Populations," *The American Statistician*, Vol. 41, No. 3, pp. 215-218.

Suppose that the universe size is $N = 10,000$, the anticipated rate of occurrence (i.e., error rate) is 20%, and the desired precision range is 6%. Since $(10,000)(.06)$ is 600, we know that $k_2 = k_1 + 600$; that is, the upper confidence limit must be 600 more than the lower limit. The anticipated rate of occurrence is used to specify the number of sample items that contain the characteristic of interest. Here, it would be 20% of n , where n is the sample size determined by this program.

For example, suppose that $n = 300$ and $(300)(.20) = 60$ (call this x). If the values, $N = 10,000$, $n = 300$, and $x = 60$ are used as input to the **Unrestricted Attribute Appraisal** program, the resulting 95% confidence interval for the universe proportion (p) has a lower limit of .1569 [i.e., $k_1 = (10,000)(.1569) = 1,569$] and an upper limit of .2490 [i.e., $k_2 = (10,000)(.2490) = 2,490$]. But this is not a satisfactory value of n since $k_2 - k_1 = 2,490 - 1,569 = 921$, which must equal 600 according to the previous discussion.

Summary of program procedure. For a specified confidence level of 95%, this program searches for the value of n that produces a confidence interval (k_1 to k_2) such that k_1 and k_2 satisfy the preceding two inequalities and $k_2 - k_1 = 600$, where, in general, 600 is equal to $N \cdot$ (desired precision range). For the preceding example, if $n = 666$, then $(666)(.20) \approx 133$. If the values $N = 10,000$, $n = 666$, and $x = 133$ are used as input to the **Unrestricted Attribute Appraisal** module, the resulting 95% confidence interval for the universe proportion (p) has a lower limit of .1710 [i.e., $k_1 = (10,000)(.1710) = 1,710$] and an upper limit of .2310 [i.e.,

$k_2 = (10,000)(.2310) = 2,310$. This is satisfactory, since $k_2 - k_1 = 600$ and the difference of the two proportions is .06 (i.e., 6%).