

# Statistics Handouts for Graphing Calculator

## Chapter 4 Counting

To find arrangement of items when the order does not matter use the Combination:

1. Type n (bigger number)
2. Hit Math key
3. Select Prob
4. Select nCr
5. Type r (smaller number)

To find arrangement of items when the order does matter use the Permutation:

1. Type n (bigger number)
2. Hit Math key
3. Select Prob
4. Select nPr
5. Type r (smaller number)

## Chapter 5 Binomial Distribution

To find the probability:

1. Exactly or equal  $2^{\text{nd}}$  Vars binomialpdf (n, p, x)
2. At most or less than or equal  $2^{\text{nd}}$  Vars binomialcdf (n, p, x)
3. Less than  $2^{\text{nd}}$  Vars binomialcdf (n, p, x-1)
4. At least or greater than or equal  $2^{\text{nd}}$  Vars 1-binomialcdf (n, p, x-1)
5. Greater than  $2^{\text{nd}}$  Vars 1-binomialcdf (n, p, x)

## Chapter 6 Normal Distribution

To find the probability, percentage or area when  $n=1$ :  $2^{\text{nd}}$  Vars normalcdf (lower, upper, mean, standard deviation)

To find the probability, percentage or area when  $n>1$ :  $2^{\text{nd}}$  Vars normalcdf (lower, upper, mean, standard deviation/ $\sqrt{n}$ )

To find the number or the score:  $2^{\text{nd}}$  Vars invnorm (left area, mean, standard deviation)

## Chapter 7 Confidence Interval

**To find Confidence Interval for proportion (single population):**

STAT

TESTS

1-PropZInt

**To find Confidence Interval for mean (single population):**

1. If population standard deviation  $\sigma$  given STAT TESTS Zinterval
2. If sample standard deviation  $s$  given STAT TESTS Tinterval

## Chapter 8 Hypothesis Testing

**To test a claim for proportion (single population):**

STAT

TESTS

1-PropZTest

**To test a claim for mean (single population):**

1. If population standard deviation  $\sigma$  given STAT TESTS ZTest
2. If sample standard deviation  $s$  given STAT TESTS TTest

## Chapter 9 Inferences from Two Samples

**To find Confidence Interval for the difference of proportions (dual population):**

STAT

TESTS

2-PropZInt

**To find Confidence Interval for the difference of means (dual population):**

1. If population standard deviations  $\sigma_1, \sigma_2$  given STAT TESTS 2-SampZinterval
2. If sample standard deviations  $s_1, s_2$  given STAT TESTS 2-SampTinterval

**To find Confidence Interval for dependent pairs  $\mu_d$  (dual population):**

1. Put the data in L1 and L2
2. In L3 type L1 - L2
3. STAT
4. TESTS

5. TInterval

**To test the claim about the dependent pairs  $\mu_d$  (dual population):**

1. Put the data in L1 and L2
2. In L3 type L1 - L2
3. STAT
4. TESTS
5. TTest

**To test the claim for the difference of proportions (dual population):**

STAT

TESTS

2-PropZTest

**To test the claim for the difference of means (dual population):**

1. If population standard deviations  $\sigma_1, \sigma_2$  given STAT TESTS 2-SampZTest
2. If sample standard deviations  $s_1, s_2$  given STAT TESTS 2-SampTTest

## **Chapter 10 Correlation**

**To test the claim for correlation:**

1. Put the data in L1 and L2
2. STAT
3. TESTS
4. LinRegTtest

## **Chapter 11 Goodness of Fit**

**To test the claim for good fit:**

1. Enter observed frequency in L1
2. Enter expected frequency in L2
3. STAT
4. TESTS
5.  $\chi^2$  GOF-Test